


(五) 集合: 关系 (Relation)


魏恒峰


hfwei@nju.edu.cn

2021 年 04 月 08 日










I'm so excited.



离散数学学得好不好，
一个重要的衡量标准就是是否完成了这种转变



I'm so excited.



The Relational Data Model

A Relational Model of Data for Large Shared Data Banks


E. F. CODD

IBM Research Laboratory, San Jose, California

Future users of large data banks must be protected from having to know how the data is organized in the machine (the internal representation). A prompting service which supplies such information is not a satisfactory solution. Activities of users at terminals and most application programs should remain unaffected when the internal representation of data is changed and even when some aspects of the external representation are changed. Changes in data representation will often be needed as a result of changes in query, update, and report traffic and natural growth in the types of stored information.

Existing noninferential, formatted data systems provide users with tree-structured files or slightly more general network models of the data. In Section 1, inadequacies of these models are discussed. A model based on n -ary relations, a normal form for data base relations, and the concept of a universal data sublanguage are introduced. In Section 2, certain operations on relations (other than logical inference) are discussed and applied to the problems of redundancy and consistency in the user's model.

Codd@CACM'1970
(Turing Award'1981)



Edgar F. Codd (1923 – 2003)

The Relational Data Model — 如何靠“关系”赢得图灵奖?

A Relational Model of Data for Large Shared Data Banks


E. F. CODD

IBM Research Laboratory, San Jose, California

Future users of large data banks must be protected from having to know how the data is organized in the machine (the internal representation). A prompting service which supplies such information is not a satisfactory solution. Activities of users at terminals and most application programs should remain unaffected when the internal representation of data is changed and even when some aspects of the external representation are changed. Changes in data representation will often be needed as a result of changes in query, update, and report traffic and natural growth in the types of stored information.

Existing noninferential, formatted data systems provide users with tree-structured files or slightly more general network models of the data. In Section 1, inadequacies of these models are discussed. A model based on n -ary relations, a normal form for data base relations, and the concept of a universal data sublanguage are introduced. In Section 2, certain operations on relations (other than logical inference) are discussed and applied to the problems of redundancy and consistency in the user's model.

Codd@CACM'1970
(Turing Award'1981)



Edgar F. Codd (1923 – 2003)

\mathbb{R} : 实数集

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

$$\forall a \in X. (a, a) \in R$$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R)$$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R) \quad (\text{对称性})$$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R) \quad (\text{对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R)$$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R) \quad (\text{对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$

\mathbb{R} : 实数集

“Near” 关系: $|a - b| < 1$

$$R = \{(a, b) \mid |a - b| < 1\}$$

$$(0, 0.618) \in R \quad (-0.618, 0.618) \notin R$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R) \quad (\text{对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$

自反性 + 反对称性 = 相容关系

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R =$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$

$$\forall a \in X. (a, a) \in R$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b)$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R)$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$

X 上的整除关系

$$R = \{(1, 2), \dots, (4, 12), \dots, (12, 60), \dots, (4, 60), \dots, (60, 60)\}$$


$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$


$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$

自反性 + 反对称性 + 传递性 = 偏序关系

$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$



$$X = \{1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60\}$$



“偏序”严格刻画了人类对于“序”的直观理解

\mathbb{N} ：自然数集

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

\mathbb{N} ：自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R$$

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$\forall a \in X. (a, a) \in R$ (自反性)

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b)$$

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R)$$

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$

$$\forall a, b \in X. ((a, b) \in R \vee (b, a) \in R)$$

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$

$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$

$$\forall a, b \in X. ((a, b) \in R \vee (b, a) \in R) \quad (\text{连接性})$$

\mathbb{N} : 自然数集

$$\leq = \{(a, b) \mid a \leq b\}$$

$$\forall a \in X. (a, a) \in R \quad (\text{自反性})$$


$$\forall a, b \in X. ((a, b) \in R \wedge (b, a) \in R \rightarrow a = b) \quad (\text{反对称性})$$

$$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R) \quad (\text{传递性})$$


$$\forall a, b \in X. ((a, b) \in R \vee (b, a) \in R) \quad (\text{连接性})$$

自反性 + 反对称性 + 传递性 + 连接性 = **全序关系**


考虑无向图中的**顶点**集合





考虑无向图中的顶点集合




顶点间的“可达 (Reachability) 关系”: $R = \{(a, b) \mid a \sim b\}$




$$\forall a \in X. (a, a) \in R$$




$\forall a \in X. (a, a) \in R$ (自反性)




$\forall a \in X. (a, a) \in R$ (自反性)

$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R)$



$\forall a \in X. (a, a) \in R$ (自反性)


$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R)$ (对称性)



$\forall a \in X. (a, a) \in R$ (自反性)

$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R)$ (对称性)


$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R)$



$\forall a \in X. (a, a) \in R$ (自反性)

$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R)$ (对称性)

$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R)$ (传递性)




$\forall a \in X. (a, a) \in R$ (自反性)

$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R)$ (对称性)

$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R)$ (传递性)

自反性 + 对称性 + 传递性 = 等价关系



$\forall a \in X. (a, a) \in R$ (自反性)

$\forall a, b \in X. ((a, b) \in R \rightarrow (b, a) \in R)$ (对称性)

$\forall a, b, c \in X. ((a, b) \in R \wedge (b, c) \in R \rightarrow (a, c) \in R)$ (传递性)

自反性 + 对称性 + 传递性 = 等价关系

“可达关系”将顶点划分成相互独立的“连通分量”




Definition (有序对**公理** (Ordered Pairs))

$$(a, b) = (c, d) \iff a = c \wedge b = d$$


Definition (Ordered Pairs (Norbert Wiener; 1914))

$$(a, b) \triangleq \left\{ \{\{a\}, \emptyset\}, \{\{b\}\} \right\}$$



Definition (Ordered Pairs (Norbert Wiener; 1914))

$$(a, b) \triangleq \left\{ \{\{a\}, \emptyset\}, \{\{b\}\} \right\}$$



Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Definition (Ordered Pairs (Kazimierz Kuratowski; 1921))

$$(a, b) \triangleq \{\{a\}, \{a, b\}\}$$



Definition (Ordered Pairs (Kazimierz Kuratowski; 1921))

$$(a, b) \triangleq \{\{a\}, \{a, b\}\}$$



Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

$$\left(\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\} \right) \iff (a = c \wedge b = d)$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

$$\left(\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\} \right) \iff (a = c \wedge b = d)$$

$$\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\}$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

$$\left(\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\} \right) \iff (a = c \wedge b = d)$$

$$\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\}$$

$$\iff (\{a\} = \{c\} \vee \{a\} = \{c, d\}) \wedge (\{a, b\} = \{c\} \vee \{a, b\} = \{c, d\})$$

Theorem

$$(a, b) = (c, d) \iff a = c \wedge b = d$$

$$\left(\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\} \right) \iff (a = c \wedge b = d)$$

$$\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\}$$

$$\iff (\{a\} = \{c\} \vee \{a\} = \{c, d\}) \wedge (\{a, b\} = \{c\} \vee \{a, b\} = \{c, d\})$$

$$\iff (\{a\} = \{c\} \wedge \{a, b\} = \{c\}) \vee$$

$$(\{a\} = \{c\} \wedge \{a, b\} = \{c, d\}) \vee$$

$$(\{a\} = \{c, d\} \wedge \{a, b\} = \{c\}) \vee$$

$$(\{a\} = \{c, d\} \wedge \{a, b\} = \{c, d\})$$

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

$$(x_1, x_2, \dots, x_{n-1}, x_n) \triangleq ((x_1, x_2, \dots, x_{n-1}), x_n)$$

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

$$(x_1, x_2, \dots, x_{n-1}, x_n) \triangleq ((x_1, x_2, \dots, x_{n-1}), x_n)$$

Theorem

$$(x_1, \dots, x_n) = (y_1, \dots, y_n) \iff x_1 = y_1 \wedge \dots x_n = y_n$$

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

$$(x_1, x_2, \dots, x_{n-1}, x_n) \triangleq ((x_1, x_2, \dots, x_{n-1}), x_n)$$

Theorem

$$(x_1, \dots, x_n) = (y_1, \dots, y_n) \iff x_1 = y_1 \wedge \dots x_n = y_n$$

By mathematical induction.

Definition (n -元组 (n-ary tuples))

$$(x, y, z) \triangleq ((x, y), z)$$

$$(x_1, x_2, \dots, x_{n-1}, x_n) \triangleq ((x_1, x_2, \dots, x_{n-1}), x_n)$$

Theorem

$$(x_1, \dots, x_n) = (y_1, \dots, y_n) \iff x_1 = y_1 \wedge \dots x_n = y_n$$

By mathematical induction.

多数情况下, 我们仅处理“二元关系”, 因此也仅使用“有序对”

Definition (笛卡尔积 (Cartesian Products))

The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$

Definition (笛卡尔积 (Cartesian Products))

The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$


$$X^2 \triangleq X \times X$$


Definition (笛卡尔积 (Cartesian Products))

The *Cartesian product* $A \times B$ of A and B is defined as

$$A \times B \triangleq \{(a, b) \mid a \in A \wedge b \in B\}$$


$$X^2 \triangleq X \times X$$





$$\mathbb{Z}^2 \triangleq \mathbb{Z} \times \mathbb{Z}$$

Ranks = {2, ..., 10, J, Q, K, A}



Suits = {♠, ♥, ♣, ♦}

$$X \times \emptyset = \emptyset \times X$$

$$X \times \emptyset = \emptyset \times X$$

$$X \times Y \neq Y \times X$$

$$X \times \emptyset = \emptyset \times X$$

$$X \times Y \neq Y \times X$$

$$(X \times Y) \times Z \neq X \times (Y \times Z)$$

$$X \times \emptyset = \emptyset \times X$$

$$X \times Y \neq Y \times X$$

$$(X \times Y) \times Z \neq X \times (Y \times Z)$$

$$A = \{1\} \quad (A \times A) \times A \neq A \times (A \times A)$$

Theorem (分配律 (Distributivity))

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$A \times (B \cup C) = (A \times B) \cup (A \times C)$$


$$A \times (B \setminus C) = (A \times B) \setminus (A \times C)$$

Theorem (分配律 (Distributivity))

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$A \times (B \setminus C) = (A \times B) \setminus (A \times C)$$



$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

对任意有序对 (a, b) ,

$$(a, b) \in A \times (B \cap C) \quad (1)$$

(6)

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

对任意有序对 (a, b) ,

$$(a, b) \in A \times (B \cap C) \quad (1)$$

$$\iff a \in A \wedge b \in (B \cap C) \quad (2)$$

(6)

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

对任意有序对 (a, b) ,

$$(a, b) \in A \times (B \cap C) \quad (1)$$

$$\iff a \in A \wedge b \in (B \cap C) \quad (2)$$

$$\iff a \in A \wedge b \in B \wedge b \in C \quad (3)$$

(6)

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

对任意有序对 (a, b) ,

$$(a, b) \in A \times (B \cap C) \quad (1)$$

$$\iff a \in A \wedge b \in (B \cap C) \quad (2)$$

$$\iff a \in A \wedge b \in B \wedge b \in C \quad (3)$$

$$\iff (a \in A \wedge b \in B) \wedge (a \in A \wedge b \in C) \quad (4)$$

(6)

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

对任意有序对 (a, b) ,

$$(a, b) \in A \times (B \cap C) \quad (1)$$

$$\iff a \in A \wedge b \in (B \cap C) \quad (2)$$

$$\iff a \in A \wedge b \in B \wedge b \in C \quad (3)$$

$$\iff (a \in A \wedge b \in B) \wedge (a \in A \wedge b \in C) \quad (4)$$

$$\iff (a, b) \in A \times B \wedge (a, b) \in A \times C \quad (5)$$

$$(6)$$

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

对任意有序对 (a, b) ,

$$(a, b) \in A \times (B \cap C) \quad (1)$$

$$\iff a \in A \wedge b \in (B \cap C) \quad (2)$$

$$\iff a \in A \wedge b \in B \wedge b \in C \quad (3)$$

$$\iff (a \in A \wedge b \in B) \wedge (a \in A \wedge b \in C) \quad (4)$$

$$\iff (a, b) \in A \times B \wedge (a, b) \in A \times C \quad (5)$$

$$\iff (a, b) \in (A \times B) \cap (A \times C) \quad (6)$$

Definition (n -元笛卡尔积 (n -ary Cartesian Product))

$$X_1 \times X_2 \times X_3 \triangleq (X_1 \times X_2) \times X_3$$

Definition (n -元笛卡尔积 (n -ary Cartesian Product))

$$X_1 \times X_2 \times X_3 \triangleq (X_1 \times X_2) \times X_3$$

$$X_1 \times X_2 \times \cdots \times X_n \triangleq (X_1 \times X_2 \times \cdots \times X_{n-1}) \times X_n$$

Definition (n -元笛卡尔积 (n -ary Cartesian Product))

$$X_1 \times X_2 \times X_3 \triangleq (X_1 \times X_2) \times X_3$$

$$X_1 \times X_2 \times \cdots \times X_n \triangleq (X_1 \times X_2 \times \cdots \times X_{n-1}) \times X_n$$

$$X^n \triangleq \underbrace{X \times \cdots \times X}_n$$

Definition (n -元笛卡尔积 (n -ary Cartesian Product))

$$X_1 \times X_2 \times X_3 \triangleq (X_1 \times X_2) \times X_3$$

$$X_1 \times X_2 \times \cdots \times X_n \triangleq (X_1 \times X_2 \times \cdots \times X_{n-1}) \times X_n$$

$$X^n \triangleq \underbrace{X \times \cdots \times X}_n$$

多数情况下，我们仅处理“二元关系”，因此也仅使用“二元笛卡尔积”

Definition (关系 (Relations))

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Definition (关系 (Relations))

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

If $A = B$, R is called a relation **on** A .

Definition (关系 (Relations))

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

If $A = B$, R is called a relation **on** A .

Definition (Notations)

$$(a, b) \in R \quad aRb$$

$$(a, b) \notin R \quad a\overline{R}b$$

Definition (Relations)

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Examples

Definition (Relations)

A *relation* R from A to B is a **subset** of $A \times B$:

$$R \subseteq A \times B$$

Examples

- Both $A \times B$ and \emptyset are relations from A to B .

Definition (Relations)

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Examples

- Both $A \times B$ and \emptyset are relations from A to B .



$$< = \{(a, b) \in \mathbb{R} \times \mathbb{R} \mid a \text{ is less than } b\}$$

Definition (Relations)

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Examples

- ▶ Both $A \times B$ and \emptyset are relations from A to B .
- ▶
 $< = \{(a, b) \in \mathbb{R} \times \mathbb{R} \mid a \text{ is less than } b\}$
- ▶
 $D = \{(a, b) \in \mathbb{N} \times \mathbb{N} \mid \exists q \in \mathbb{N} : a \cdot q = b\}$

Definition (Relations)

A *relation* R from A to B is a subset of $A \times B$:

$$R \subseteq A \times B$$

Examples


- ▶ Both $A \times B$ and \emptyset are relations from A to B .
- ▶
 $< = \{(a, b) \in \mathbb{R} \times \mathbb{R} \mid a \text{ is less than } b\}$
- ▶
 $D = \{(a, b) \in \mathbb{N} \times \mathbb{N} \mid \exists q \in \mathbb{N} : a \cdot q = b\}$
- ▶ P : the set of people
 - $M = \{(a, b) \in P \times P \mid a \text{ is the mother of } b\}$
 - $B = \{(a, b) \in P \times P \mid a \text{ is the brother of } b\}$

Important Relations:

Equivalence Relations

Ordering Relations (next class)

Functions (next class)



Outline:

3 Definitions

5 Operations

7 Properties

3 Definitions

$\text{dom}(R)$ $\text{ran}(R)$ $\text{fld}(R)$

Definition (定义域 (Domain))

$$\text{dom}(R) = \{a \mid \exists b. (a, b) \in R\}$$

Definition (定义域 (Domain))

$$\text{dom}(R) = \{a \mid \exists b. (a, b) \in R\}$$

Definition (值域 (Range))

$$\text{ran}(R) = \{b \mid \exists a : (a, b) \in R\}$$

Definition (定义域 (Domain))

$$\text{dom}(R) = \{a \mid \exists b. (a, b) \in R\}$$

Definition (值域 (Range))

$$\text{ran}(R) = \{b \mid \exists a : (a, b) \in R\}$$

Definition (域 (Field))

$$\text{fld}(R) = \text{dom}(R) \cup \text{ran}(R)$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$\text{dom}(R) = \mathbb{R} \quad \text{ran}(R) = \mathbb{R} \quad \text{fld}(R) = \mathbb{R}$$

$$R = \{(x, y) \mid x^2 + y^2 = 1\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$R = \{(x, y) \mid x^2 + y^2 = 1\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$\text{dom}(R) = [1, 1] \quad \text{ran}(R) = [-1, 1] \quad \text{fld}(R) = [-1, 1]$$

Theorem

$$dom(R) \subseteq \bigcup \bigcup R \quad ran(R) \subseteq \bigcup \bigcup R$$

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{7}$$

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{7}$$

$$\implies \exists b. (a, b) \in R \tag{8}$$

(12)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{7}$$

$$\implies \exists b. (a, b) \in R \tag{8}$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \tag{9}$$

(12)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{7}$$

$$\implies \exists b. (a, b) \in R \tag{8}$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \tag{9}$$

$$\implies \exists b. \{a, b\} \in \bigcup R \tag{10}$$

(12)

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{7}$$

$$\implies \exists b. (a, b) \in R \tag{8}$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \tag{9}$$

$$\implies \exists b. \{a, b\} \in \bigcup R \tag{10}$$

$$\implies \exists b. a \in \bigcup \bigcup R \tag{11}$$

$$(12)$$

Theorem

$$\text{dom}(R) \subseteq \bigcup \bigcup R \quad \text{ran}(R) \subseteq \bigcup \bigcup R$$

对任意 a ,

$$a \in \text{dom}(R) \tag{7}$$

$$\implies \exists b. (a, b) \in R \tag{8}$$

$$\implies \exists b. \{\{a\}, \{a, b\}\} \in R \tag{9}$$

$$\implies \exists b. \{a, b\} \in \bigcup R \tag{10}$$

$$\implies \exists b. a \in \bigcup \bigcup R \tag{11}$$

$$\implies a \in \bigcup \bigcup R \tag{12}$$

5 Operations

R^{-1} $R|_X$ $R[X]$ $R^{-1}[Y]$ $R \circ S$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} =$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} =$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = \{(x, y) \mid y = x^2 \wedge x > 0\}$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = \{(x, y) \mid y = x^2 \wedge x > 0\}$$

$$\leq = \{(x, y) \mid x \leq y\} \subseteq \mathbb{R} \times \mathbb{R} \quad \leq^{-1} =$$

Definition (逆 (Inverse))

The *inverse* of R is the **relation**

$$R^{-1} = \{(a, b) \mid (b, a) \in R\}$$

$$R = \{(x, y) \mid x = y\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = R$$

$$R = \{(x, y) \mid y = \sqrt{x}\} \subseteq \mathbb{R} \times \mathbb{R} \quad R^{-1} = \{(x, y) \mid y = x^2 \wedge x > 0\}$$

$$\leq = \{(x, y) \mid x \leq y\} \subseteq \mathbb{R} \times \mathbb{R} \quad \leq^{-1} = \geq \triangleq \{(x, y) \mid x \geq y\}$$

Theorem

$$(R^{-1})^{-1} = R$$

Theorem

$$(R^{-1})^{-1} = R$$

对任意 (a, b) ,

$$(a, b) \in (R^{-1})^{-1} \quad (1)$$

(3)

Theorem

$$(R^{-1})^{-1} = R$$

对任意 (a, b) ,

$$(a, b) \in (R^{-1})^{-1} \quad (1)$$

$$\iff (b, a) \in R^{-1} \quad (2)$$

(3)

Theorem

$$(R^{-1})^{-1} = R$$

对任意 (a, b) ,

$$(a, b) \in (R^{-1})^{-1} \quad (1)$$

$$\iff (b, a) \in R^{-1} \quad (2)$$

$$\iff (a, b) \in R \quad (3)$$

Theorem (关系的逆)

R, S 均为关系

$$(R \cup S)^{-1} = R^{-1} \cup S^{-1}$$

$$(R \cap S)^{-1} = R^{-1} \cap S^{-1}$$

$$(R \setminus S)^{-1} = R^{-1} \setminus S^{-1}$$

Definition (左限制 (Left-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq X$. The *left-restriction* relation of R to S over X and Y is

$$R|_S = \{(x, y) \in R \mid x \in S\}$$

Definition (左限制 (Left-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq X$. The *left-restriction* relation of R to S over X and Y is

$$R|_S = \{(x, y) \in R \mid x \in S\}$$

Definition (右限制 (Right-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq Y$. The *right-restriction* relation of R to S over X and Y is

$$R|^S = \{(x, y) \in R \mid y \in S\}$$

Definition (左限制 (Left-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq X$. The *left-restriction* relation of R to S over X and Y is

$$R|_S = \{(x, y) \in R \mid x \in S\}$$

Definition (右限制 (Right-Restriction))

Suppose $R \subseteq X \times Y$ and $S \subseteq Y$. The *right-restriction* relation of R to S over X and Y is

$$R|^S = \{(x, y) \in R \mid y \in S\}$$

Definition (限制 (Restriction))

Suppose $R \subseteq X \times X$ and $S \subseteq X$. The *restriction* relation of R to S over X is

$$R|_S = \{(x, y) \in R \mid x \in S \wedge y \in S\}$$

$$R = \{(x, y) \mid x^2 + y^2 = 1\} \subseteq \mathbb{R} \times \mathbb{R}$$

$$R = \{(x, y) \mid x^2 + y^2 = 1\} \subseteq \mathbb{R} \times \mathbb{R}$$

$R|_{\mathbb{R}^+}$ (left restriction, restriction)

$$R = \{(x, y) \mid x^2 + y^2 = 1\} \subseteq \mathbb{R} \times \mathbb{R}$$

$R|_{\mathbb{R}^+}$ (left restriction, restriction)

$R|^{\mathbb{R}^+}$ (right restriction)

Definition (像 (Image))

The *image* of X under R is the set

$$R[X] = \{b \in \text{ran}(R) \mid \exists a \in X. (a, b) \in R\}$$

Definition (像 (Image))

The *image* of X under R is the set

$$R[X] = \{b \in \text{ran}(R) \mid \exists a \in X. (a, b) \in R\}$$

$$R[a] \triangleq R[\{a\}] = \{b \mid (a, b) \in R\}$$

Definition (逆像 (Inverse Image))

The *inverse image* of Y under R is the set

$$R^{-1}[Y] = \{a \in \text{dom}(R) \mid \exists b \in Y. (a, b) \in R\}$$

Definition (逆像 (Inverse Image))

The *inverse image* of Y under R is the set

$$R^{-1}[Y] = \{a \in \text{dom}(R) \mid \exists b \in Y. (a, b) \in R\}$$

$$R^{-1}[b] \triangleq R^{-1}[\{b\}] = \{a \mid (a, b) \in R\}$$

$$R \subseteq A \times B \quad X \subseteq A \quad Y \subseteq B$$

$$R \subseteq A \times B \quad X \subseteq A \quad Y \subseteq B$$

$$R^{-1}[R[X]] \textcolor{red}{?} X$$

$$R[R^{-1}[Y]] \textcolor{red}{?} Y$$

$$R \subseteq A \times B \quad X \subseteq A \quad Y \subseteq B$$

$$R^{-1}[R[X]] \textcolor{red}{?} X$$

$$R[R^{-1}[Y]] \textcolor{red}{?} Y$$



Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \cup X_2. (a, b) \in R$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \cup X_2. (a, b) \in R$$

$$\iff \exists a \in X_1. (a, b) \in R \vee \exists a \in X_2. (a, b) \in R$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \cup X_2. (a, b) \in R$$

$$\iff \exists a \in X_1. (a, b) \in R \vee \exists a \in X_2. (a, b) \in R$$

$$\iff b \in R[X_1] \vee b \in R[X_2]$$

Theorem

$$R[X_1 \cup X_2] = R[X_1] \cup R[X_2]$$

$$R[X_1 \cap X_2] \subseteq R[X_1] \cap R[X_2]$$

$$R[X_1 \setminus X_2] \supseteq R[X_1] \setminus R[X_2]$$

对任意 b ,

$$b \in R[X_1 \cup X_2]$$

$$\iff \exists a \in X_1 \text{ } \textcolor{red}{\cup} \text{ } X_2. (a, b) \in R$$

$$\iff \exists a \in X_1. (a, b) \in R \vee \exists a \in X_2. (a, b) \in R$$

$$\iff b \in R[X_1] \vee b \in R[X_2]$$

$$\iff b \in R[X_1] \cup R[X_2]$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S =$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R =$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R =$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\}$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R =$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S =$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S = \{(2, 2), (2, 3)\}$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$

$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S = \{(2, 2), (2, 3)\} \quad (S \circ S) \circ S =$$

Definition (复合 (Composition; $R \circ S$, $R; S$))

The *composition* of relations $R \subseteq X \times Y$ and $S \subseteq Y \times Z$ is the **relation**

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$


$$R = \{(1, 2), (3, 1)\} \quad S = \{(1, 3), (2, 2), (2, 3)\}$$

$$R \circ S = \{(1, 1), (2, 1)\}$$


$$S \circ R = \{(1, 2), (1, 3), (3, 3)\}$$

$$R^{(2)} \triangleq R \circ R = \{(3, 2)\} \quad (R \circ R) \circ R = \emptyset$$

$$S^{(2)} \triangleq S \circ S = \{(2, 2), (2, 3)\} \quad (S \circ S) \circ S = \{(2, 2), (2, 3)\}$$



$$|R \circ S| =$$



$$|R \circ S| = 7$$

$\leq \circ \leq =$

$\leq \circ \leq = \leq$

$\leq \circ \leq = \leq$ $\leq \circ \geq =$

$$\leq \circ \leq = \leq$$

$$\leq \circ \geq = \mathbb{R} \times \mathbb{R}$$

$$\leq \circ \leq = \leq$$

$$\leq \circ \geq = \mathbb{R} \times \mathbb{R}$$

$$\forall a, b \in \mathbb{R}. (a, b) \in \leq \circ \geq$$

$$\leq \circ \leq = \leq$$

$$\leq \circ \geq = \mathbb{R} \times \mathbb{R}$$

$\forall a, b \in \mathbb{R}. (a, b) \in \leq \circ \geq$

$$(a, |a| + |b|) \quad (|a| + |b|, b)$$

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

(5)

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

(5)

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

$$\iff \exists c. (b, \textcolor{blue}{c}) \in R \wedge (\textcolor{blue}{c}, a) \in S \quad (3)$$

(5)

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

$$\iff \exists c. (b, \textcolor{blue}{c}) \in R \wedge (\textcolor{blue}{c}, a) \in S \quad (3)$$

$$\iff \exists c. (\textcolor{red}{c}, b) \in R^{-1} \wedge (a, \textcolor{red}{c}) \in S^{-1} \quad (4)$$

(5)

Theorem

$$(R \circ S)^{-1} = S^{-1} \circ R^{-1}$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S)^{-1} \quad (1)$$

$$\iff (b, a) \in R \circ S \quad (2)$$

$$\iff \exists c. (b, \textcolor{blue}{c}) \in R \wedge (\textcolor{blue}{c}, a) \in S \quad (3)$$

$$\iff \exists c. (\textcolor{red}{c}, b) \in R^{-1} \wedge (a, \textcolor{red}{c}) \in S^{-1} \quad (4)$$

$$\iff (a, b) \in S^{-1} \circ R^{-1} \quad (5)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. \left((a, c) \in T \wedge (c, b) \in R \circ S \right) \quad (2)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

$$\iff \exists d. \exists c. ((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R) \quad (4)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

$$\iff \exists d. \exists c. ((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R) \quad (4)$$

$$\iff \exists d. ((\exists c. (a, c) \in T \wedge (c, d) \in S) \wedge (d, b) \in R) \quad (5)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

$$\iff \exists d. \exists c. ((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R) \quad (4)$$

$$\iff \exists d. ((\exists c. (a, c) \in T \wedge (c, d) \in S) \wedge (d, b) \in R) \quad (5)$$

$$\iff \exists d. ((a, d) \in S \circ T \wedge (d, b) \in R) \quad (6)$$

Theorem

$$(R \circ S) \circ T = R \circ (S \circ T)$$

对任意 (a, b) ,

$$(a, b) \in (R \circ S) \circ T \quad (1)$$

$$\iff \exists c. ((a, c) \in T \wedge (c, b) \in R \circ S) \quad (2)$$

$$\iff \exists c. ((a, c) \in T \wedge (\exists d : (c, d) \in S \wedge (d, b) \in R)) \quad (3)$$

$$\iff \exists d. \exists c. ((a, c) \in T \wedge (c, d) \in S \wedge (d, b) \in R) \quad (4)$$

$$\iff \exists d. ((\exists c. (a, c) \in T \wedge (c, d) \in S) \wedge (d, b) \in R) \quad (5)$$

$$\iff \exists d. ((a, d) \in S \circ T \wedge (d, b) \in R) \quad (6)$$

$$\iff (a, b) \in R \circ (S \circ T)$$



帮我照顾好我七舅姥爷和我外甥女

燕小六: “帮我照顾好我七舅姥爷和我外甥女”

“舅姥爷”: 姥姥/外婆的兄弟

“舅姥爷”：姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

“舅姥爷”: 姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

“舅姥爷”: 姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

$$G = B \circ (M \circ M)$$

“舅姥爷”: 姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

$$G = B \circ (M \circ M)$$

$$G = B \circ (M \circ M) = (B \circ M) \circ M$$

“舅姥爷”: 姥姥/外婆的兄弟

$$G = \{(a, b) : a \text{ 是 } b \text{ 的舅姥爷}\}$$

$$B = \{(a, b) \mid a \text{ is the brother of } b\}$$

$$M = \{(a, b) \mid a \text{ is the mother of } b\}$$

$$G = B \circ (M \circ M)$$

$$G = B \circ (M \circ M) = (B \circ M) \circ M$$

“舅姥爷”: 妈妈的舅舅

Theorem (关系的复合)

$$(X \cup Y) \circ Z = (X \circ Z) \cup (Y \circ Z)$$

$$(X \cap Y) \circ Z \subseteq (X \circ Z) \cap (Y \circ Z)$$


7 Properties

$$R \subseteq X \times X$$

Definition (自反的 (Reflexive))

$R \subseteq X \times X$ is *reflexive* if


$$\forall a \in X : (a, a) \in R$$



Definition (自反的 (Reflexive))

$R \subseteq X \times X$ is *reflexive* if

$$\forall a \in X : (a, a) \in R$$




$\leq \subseteq \mathbb{R} \times \mathbb{R}$ is reflexive

Definition (自反的 (Reflexive))

$R \subseteq X \times X$ is *reflexive* if

$$\forall a \in X : (a, a) \in R$$



$\leq \subseteq \mathbb{R} \times \mathbb{R}$ is reflexive

三角形上的全等关系是自反的

Definition (反自反 (Irreflexive))

$R \subseteq X \times X$ is *irreflexive* if

$$\forall a \in X. (a, a) \notin R$$

Definition (反自反 (Irreflexive))

$R \subseteq X \times X$ is *irreflexive* if

$$\forall a \in X. (a, a) \notin R$$

$< \subseteq \mathbb{R} \times \mathbb{R}$ is irreflexive

Definition (反自反 (Irreflexive))

$R \subseteq X \times X$ is *irreflexive* if

$$\forall a \in X. (a, a) \notin R$$

$< \subseteq \mathbb{R} \times \mathbb{R}$ is irreflexive

$> \subseteq \mathbb{R} \times \mathbb{R}$ is irreflexive

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 3), (2, 1), (2, 2), (3, 3)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 3), (2, 1), (2, 2), (3, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 3), (2, 1), (2, 2), (3, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$\{(1, 2), (2, 2), (2, 3), (3, 1)\}$$

Definition (对称 (Symmetric))

$R \subseteq X \times X$ is *symmetric* if

$$\forall a, b \in X. aRb \rightarrow bRa$$



Definition (对称 (Symmetric))

$R \subseteq X \times X$ is *symmetric* if

$$\forall a, b \in X. aRb \rightarrow bRa$$



$$\forall a, b \in X. aRb \leftrightarrow bRa$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$

$$\{(1, 1), (2, 2), (3, 3)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$

$$\{(1, 1), (2, 2), (3, 3)\}$$

$$\{(1, 2), (2, 1), (2, 3)\}$$

Definition (反对称 (AntiSymmetric))

$R \subseteq X \times X$ is *antisymmetric* if

$$\forall a, b \in X. (aRb \wedge bRa) \rightarrow a = b$$

Definition (反对称 (AntiSymmetric))

$R \subseteq X \times X$ is *antisymmetric* if

$$\forall a, b \in X. (aRb \wedge bRa) \rightarrow a = b$$

\geq is antisymmetric

Definition (反对称 (AntiSymmetric))

$R \subseteq X \times X$ is *antisymmetric* if

$$\forall a, b \in X. (aRb \wedge bRa) \rightarrow a = b$$

\geq is antisymmetric

| is antisymmetric

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$

$$\{(1, 1), (2, 2), (3, 3)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (3, 1), (3, 3)\}$$

$$\{(1, 2), (2, 3), (2, 2), (3, 1)\}$$


$$\{(1, 1), (2, 2), (3, 3)\}$$

$$\{(1, 2), (2, 1), (2, 3)\}$$

Definition (传递的 (Transitive))

$R \subseteq X \times X$ is *transitive* if

$$\forall a, b, c \in X. (aRb \wedge bRc \rightarrow aRc)$$



$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$\{(1, 3)\}$$

$$A = \{1, 2, 3\} \quad R \subseteq A \times A$$

$$\{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\}$$

$$\{(1, 2), (2, 3), (3, 1)\}$$

$$\{(1, 3)\}$$

\emptyset

Definition (连接的 (Connex))

$R \subseteq X \times X$ is *connex* if

$$\forall a, b \in X. (aRb \vee bRa)$$

Definition (连接的 (Connex))

$R \subseteq X \times X$ is *connex* if

$$\forall a, b \in X. (aRb \vee bRa)$$

Definition (三分的 (Trichotomous))

$R \subseteq X \times X$ is *trichotomous* if

$$\forall a, b \in X. (\text{exactly one of } aRb, bRa, \text{ or } a = b \text{ holds})$$

Theorem

R is reflexive $\iff I \subseteq R$

$$I = \{(a, a) \in A \times A \mid a \in A\}$$

Theorem

$$R \text{ is reflexive} \iff I \subseteq R$$

$$I = \{(a, a) \in A \times A \mid a \in A\}$$

Theorem

$$R \text{ is symmetric} \iff R^{-1} = R$$

Theorem

R is transitive $\iff R \circ R \subseteq R$

Theorem

R is transitive $\iff R \circ R \subseteq R$

$$R = \{(1, 2), (2, 3), (1, 3), (4, 4)\}$$

Theorem

$$R \text{ is transitive} \iff R \circ R \subseteq R$$

$$R = \{(1, 2), (2, 3), (1, 3), (4, 4)\}$$

对任意 (a, b) ,

$$(a, b) \in R \circ R \tag{1}$$

$$\implies \exists c. (a, c) \in R \wedge (b, c) \in R \tag{2}$$

(3)

Theorem

$$R \text{ is transitive} \iff R \circ R \subseteq R$$

$$R = \{(1, 2), (2, 3), (1, 3), (4, 4)\}$$

对任意 (a, b) ,

$$(a, b) \in R \circ R \tag{1}$$

$$\implies \exists c. (a, c) \in R \wedge (b, c) \in R \tag{2}$$

$$\implies (a, b) \in R \tag{3}$$

Theorem

$$R \text{ is transitive} \iff R \circ R \subseteq R$$

$$R = \{(1, 2), (2, 3), (1, 3), (4, 4)\}$$

对任意 (a, b) ,

$$(a, b) \in R \circ R \tag{1}$$

$$\implies \exists c. (a, c) \in R \wedge (b, c) \in R \tag{2}$$

$$\implies (a, b) \in R \tag{3}$$

对任意 a, b, c

$$(a, b) \in R \wedge (b, c) \in R$$

Theorem

$$R \text{ is transitive} \iff R \circ R \subseteq R$$

$$R = \{(1, 2), (2, 3), (1, 3), (4, 4)\}$$

对任意 (a, b) ,

$$(a, b) \in R \circ R \tag{1}$$

$$\implies \exists c. (a, c) \in R \wedge (b, c) \in R \tag{2}$$

$$\implies (a, b) \in R \tag{3}$$

对任意 a, b, c

$$(a, b) \in R \wedge (b, c) \in R \implies (a, c) \in R$$

Theorem

R is symmetric and transitive $\iff R = R^{-1} \circ R$

Equivalence Relations

Definition (Equivalence Relation)

$R \subseteq X \times X$ is an *equivalence relation* on X iff R is

- ▶ reflexive: $\forall a \in X. aRa$
- ▶ symmetric: $\forall a, b \in X. (aRb \leftrightarrow bRa)$
- ▶ transitive: $\forall a, b, c \in X. (aRb \wedge bRc \rightarrow aRc)$

Definition (Equivalence Relation)

$R \subseteq X \times X$ is an *equivalence relation* on X iff R is

- ▶ reflexive: $\forall a \in X. aRa$
- ▶ symmetric: $\forall a, b \in X. (aRb \leftrightarrow bRa)$
- ▶ transitive: $\forall a, b, c \in X. (aRb \wedge bRc \rightarrow aRc)$

$$= \in \mathbb{R} \times \mathbb{R}$$

Definition (Equivalence Relation)

$R \subseteq X \times X$ is an *equivalence relation* on X iff R is

- ▶ reflexive: $\forall a \in X. aRa$
- ▶ symmetric: $\forall a, b \in X. (aRb \leftrightarrow bRa)$
- ▶ transitive: $\forall a, b, c \in X. (aRb \wedge bRc \rightarrow aRc)$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

Definition (Equivalence Relation)

$R \subseteq X \times X$ is an *equivalence relation* on X iff R is

- ▶ reflexive: $\forall a \in X. aRa$
- ▶ symmetric: $\forall a, b \in X. (aRb \leftrightarrow bRa)$
- ▶ transitive: $\forall a, b, c \in X. (aRb \wedge bRc \rightarrow aRc)$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

三角形的相似关系

Definition (Equivalence Relation)

$R \subseteq X \times X$ is an *equivalence relation* on X iff R is

- ▶ reflexive: $\forall a \in X. aRa$
- ▶ symmetric: $\forall a, b \in X. (aRb \leftrightarrow bRa)$
- ▶ transitive: $\forall a, b, c \in X. (aRb \wedge bRc \rightarrow aRc)$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

三角形的相似关系

$$R = \{(x, y) \mid x \equiv y \pmod{k}\} \subseteq \mathbb{Z} \times \mathbb{Z}$$

Definition (Equivalence Relation)

$R \subseteq X \times X$ is an *equivalence relation* on X iff R is

- ▶ reflexive: $\forall a \in X. aRa$
- ▶ symmetric: $\forall a, b \in X. (aRb \leftrightarrow bRa)$
- ▶ transitive: $\forall a, b, c \in X. (aRb \wedge bRc \rightarrow aRc)$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$


三角形的相似关系

$$R = \{(x, y) \mid x \equiv y \pmod{k}\} \subseteq \mathbb{Z} \times \mathbb{Z}$$


Why are equivalence relations important?

Equivalence Relations as Abstractions

Equivalence Relations as Abstractions




Equivalence Relations as Abstractions



“全国人民代表大会各省代表团”


Equivalence Relations as Abstractions



“全国人民代表大会各省代表团”

Equivalence Relation \iff Partition

Partition



“不空、不漏、不重”

Definition (划分 (Partition))

A family of sets $\Pi = \{A_\alpha \mid \alpha \in I\}$ is a *partition* of X if

(i) (不空)

$$\forall \alpha \in I. A_\alpha \neq \emptyset$$

(ii) (不漏)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

(iii) (不重)

$$\forall \alpha, \beta \in I. A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

Definition (划分 (Partition))

A family of sets $\Pi = \{A_\alpha \mid \alpha \in I\}$ is a *partition* of X if

(i) (不空)

$$\forall \alpha \in I. A_\alpha \neq \emptyset$$

$$(\forall \alpha \in I. \exists x \in X. x \in A_\alpha)$$

(ii) (不漏)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

(iii) (不重)

$$\forall \alpha, \beta \in I. A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

Definition (划分 (Partition))

A family of sets $\Pi = \{A_\alpha \mid \alpha \in I\}$ is a *partition* of X if

(i) (不空)

$$\forall \alpha \in I. A_\alpha \neq \emptyset$$

$$(\forall \alpha \in I. \exists x \in X. x \in A_\alpha)$$

(ii) (不漏)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

$$(\forall x \in X. \exists \alpha \in I. x \in A_\alpha)$$

(iii) (不重)

$$\forall \alpha, \beta \in I. A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

Definition (划分 (Partition))

A family of sets $\Pi = \{A_\alpha \mid \alpha \in I\}$ is a *partition* of X if

(i) (不空)

$$\forall \alpha \in I. A_\alpha \neq \emptyset$$

$$(\forall \alpha \in I. \exists x \in X. x \in A_\alpha)$$

(ii) (不漏)

$$\bigcup_{\alpha \in I} A_\alpha = X$$

$$(\forall x \in X. \exists \alpha \in I. x \in A_\alpha)$$

(iii) (不重)

$$\forall \alpha, \beta \in I. A_\alpha \cap A_\beta = \emptyset \vee A_\alpha = A_\beta$$

$$(\forall \alpha, \beta \in I. A_\alpha \cap A_\beta \neq \emptyset \implies A_\alpha = A_\beta)$$

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Definition (等价类 (Equivalence Class))

The *equivalence class* of a modulo R is a set:

$$[a]_R = \{b \in X : aRb\}$$

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Definition (等价类 (Equivalence Class))

The *equivalence class* of a modulo R is a set:

$$[a]_R = \{b \in X : aRb\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Definition (等价类 (Equivalence Class))

The *equivalence class* of a modulo R is a set:

$$[a]_R = \{b \in X : aRb\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Definition (等价类 (Equivalence Class))

The *equivalence class* of a modulo R is a set:

$$[a]_R = \{b \in X : aRb\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

三角形的相似关系

Equivalence Relation $R \subseteq X \times X \implies$ Partition Π of X

Definition (等价类 (Equivalence Class))

The *equivalence class* of a modulo R is a set:

$$[a]_R = \{b \in X : aRb\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

三角形的相似关系

$$R = \{(x, y) \mid x \equiv y \pmod{k}\} \subseteq \mathbb{Z} \times \mathbb{Z}$$

Definition (商集 (Quotient Set))

The *quotient set* of X by R (X modulo R) is a **set**:

$$X/R = \{[a]_R \mid a \in X\}$$

Definition (商集 (Quotient Set))

The *quotient set* of X by R (X modulo R) is a **set**:

$$X/R = \{[a]_R \mid a \in X\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

Definition (商集 (Quotient Set))

The *quotient set* of X by R (X modulo R) is a **set**:

$$X/R = \{[a]_R \mid a \in X\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

Definition (商集 (Quotient Set))

The *quotient set* of X by R (X modulo R) is a **set**:

$$X/R = \{[a]_R \mid a \in X\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

三角形的相似关系

Definition (商集 (Quotient Set))

The *quotient set* of X by R (X modulo R) is a **set**:

$$X/R = \{[a]_R \mid a \in X\}$$

$$= \in \mathbb{R} \times \mathbb{R}$$

$$\parallel \in \mathbb{L} \times \mathbb{L}$$

三角形的相似关系

$$R = \{(x, y) \mid x \equiv y \pmod{k}\} \subseteq \mathbb{Z} \times \mathbb{Z}$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X. [a]_R \neq \emptyset$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X. [a]_R \neq \emptyset$$

$$\forall a \in X. \exists b \in X. a \in [b]_R$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X. [a]_R \neq \emptyset$$

$$\forall a \in X. \exists b \in X. a \in [b]_R$$

Theorem

$$\forall a \in X, b \in X. [a]_R \cap [b]_R = \emptyset \vee [a]_R = [b]_R$$

Theorem

$X/R = \{[a]_R \mid a \in X\}$ is a partition of X .

$$\forall a \in X. [a]_R \neq \emptyset$$

$$\forall a \in X. \exists b \in X. a \in [b]_R$$

Theorem

$$\forall a \in X, b \in X. [a]_R \cap [b]_R = \emptyset \vee [a]_R = [b]_R$$

$$\forall a \in X, b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

不妨设 $x \in [a]_R \wedge [b]_R$

$$x \in [a]_R \wedge [b]_R \quad (1)$$

(3)

(4)

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

不妨设 $x \in [a]_R \wedge [b]_R$

$$x \in [a]_R \wedge [b]_R \quad (1)$$

$$\implies aRx \wedge xRb \quad (2)$$

$$(3)$$

$$(4)$$

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

不妨设 $x \in [a]_R \wedge [b]_R$

$$x \in [a]_R \wedge [b]_R \quad (1)$$

$$\implies aRx \wedge xRb \quad (2)$$

$$\implies aRb \quad (3)$$

(4)

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

不妨设 $x \in [a]_R \wedge [b]_R$ 对于任意 x ,

$$x \in [a]_R \wedge [b]_R \quad (1) \qquad x \in [a]_R \quad (1)$$

$$\implies aRx \wedge xRb \quad (2)$$

$$\implies aRb \quad (3)$$

(4)

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

不妨设 $x \in [a]_R \wedge [b]_R$ 对于任意 x ,

$$x \in [a]_R \wedge [b]_R \quad (1) \qquad x \in [a]_R \quad (1)$$

$$\implies aRx \wedge xRb \quad (2) \qquad \iff xRa \quad (2)$$

$$\implies aRb \quad (3) \qquad (4)$$

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

不妨设 $x \in [a]_R \wedge [b]_R$

对于任意 x ,

$$x \in [a]_R \wedge [b]_R \quad (1)$$

$$x \in [a]_R \quad (1)$$

$$\implies aRx \wedge xRb \quad (2)$$

$$\iff xRa \quad (2)$$

$$\implies aRb \quad (3)$$

$$\iff xRb \quad (3)$$

$$(4)$$

$$\forall a \in X. b \in X. [a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$$

不妨设 $x \in [a]_R \wedge [b]_R$

对于任意 x ,

$$x \in [a]_R \wedge [b]_R \quad (1)$$

$$\implies aRx \wedge xRb \quad (2)$$

$$\implies aRb \quad (3)$$

$$x \in [a]_R \quad (1)$$

$$\iff xRa \quad (2)$$

$$\iff xRb \quad (3)$$

$$\iff x \in [b]_R \quad (4)$$

Theorem

$$\forall a, b \in X. ([a]_R = [b]_R \leftrightarrow aRb)$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi. a \in S \wedge b \in S$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi. a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi. a \in S \wedge b \in S\}$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi. a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi. a \in S \wedge b \in S\}$$

Theorem

R is an equivalence relation on X.

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi. a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi. a \in S \wedge b \in S\}$$

Theorem

R is an equivalence relation on X.

$$\forall x \in X. xRx$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi. a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi. a \in S \wedge b \in S\}$$

Theorem

R is an equivalence relation on X.

$$\forall x \in X. xRx$$

$$\forall x, y \in X. xRy \rightarrow yRx$$

Partition Π of $X \implies$ Equivalence Relation $R \subseteq X \times X$

Definition

$$(a, b) \in R \iff \exists S \in \Pi. a \in S \wedge b \in S$$

$$R = \{(a, b) \in X \times X \mid \exists S \in \Pi. a \in S \wedge b \in S\}$$


Theorem

R is an equivalence relation on X.

$$\forall x \in X. xRx$$

$$\forall x, y \in X. xRy \rightarrow yRx$$

$$\forall x, y, z \in X. xRy \wedge yRz \rightarrow xRz$$



Equivalence Relation \iff Partition

Definition

$$\sim \subseteq \mathbb{N} \times \mathbb{N}$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_N c$$

Definition

$$\sim \subseteq \mathbb{N} \times \mathbb{N}$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_N c$$

Theorem

\sim is an equivalence relation.

Definition

$$\sim \subseteq \mathbb{N} \times \mathbb{N}$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_N c$$

Theorem

\sim is an equivalence relation.

Q : What is $\mathbb{N} \times \mathbb{N} / \sim$?

Definition

$$\sim \subseteq \mathbb{N} \times \mathbb{N}$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_N c$$

Theorem

\sim is an equivalence relation.

Q : What is $\mathbb{N} \times \mathbb{N} / \sim$?

Definition (\mathbb{Z})

$$\mathbb{Z} \triangleq \mathbb{N} \times \mathbb{N} / \sim$$

Definition

$$\sim \subseteq \mathbb{N} \times \mathbb{N}$$

$$(a, b) \sim (c, d) \iff a +_{\mathbb{N}} d = b +_N c$$

Theorem


\sim is an equivalence relation.

Q : What is $\mathbb{N} \times \mathbb{N} / \sim$?

Definition (\mathbb{Z})

$$\mathbb{Z} \triangleq \mathbb{N} \times \mathbb{N} / \sim$$

$$[(1, 3)]_{\sim} = \{(0, 2), (1, 3), (2, 4), (3, 5), \dots\} \triangleq -2 \in \mathbb{Z}$$



$$\mathbb{Z} \triangleq \mathbb{N} \times \mathbb{N} / \sim$$

Definition ($+_{\mathbb{Z}}$)

$$[(m_1, n_1)] +_{\mathbb{Z}} [(m_2, n_2)] = [m_1 +_{\mathbb{N}} m_2, n_1 +_{\mathbb{N}} n_2]$$

Definition ($+_{\mathbb{Z}}$)

$$[(m_1, n_1)] +_{\mathbb{Z}} [(m_2, n_2)] = [m_1 +_{\mathbb{N}} m_2, n_1 +_{\mathbb{N}} n_2]$$

Definition ($\cdot_{\mathbb{Z}}$)

$$\begin{aligned} & [(m_1, n_1)] \cdot_{\mathbb{Z}} [(m_2, n_2)] \\ &= [m_1 \cdot_{\mathbb{N}} m_2 +_{\mathbb{N}} n_1 \cdot_{\mathbb{N}} n_2, m_1 \cdot_{\mathbb{N}} n_2 +_{\mathbb{N}} n_1 \cdot_{\mathbb{N}} m_2] \end{aligned}$$

Definition

$$\sim \subseteq (\mathbb{Z} \times \mathbb{Z} \setminus \{0_{\mathbb{Z}}\})^2$$

$$(a, b) \sim (c, d) \iff a \cdot_{\mathbb{Z}} d = b \cdot_{\mathbb{Z}} c$$


Definition

$$\sim \subseteq (\mathbb{Z} \times \mathbb{Z} \setminus \{0_{\mathbb{Z}}\})^2$$

$$(a, b) \sim (c, d) \iff a \cdot_{\mathbb{Z}} d = b \cdot_{\mathbb{Z}} c$$

Definition (\mathbb{Q})


$$\mathbb{Q} \triangleq \mathbb{Z} \times \mathbb{Z} / \sim$$



$$\mathbb{Q} \triangleq \mathbb{Z} \times \mathbb{Z} / \sim$$


如何用有理数定义实数?

如何用有理数定义实数?



Dedekind Cut (戴德金分割)

Thank You!



Office 926

hfwei@nju.edu.cn