

(十二) 图论: 对偶 (Duality: Matching, Network Flow, and Connectivity)

魏恒峰

hfwei@nju.edu.cn

2021 年 05 月 27 日





let's get
married
today




The Marriage Problem (1935)



If there is a finite set of **girls**, each of whom knows several **boys**,
under what conditions can all the girls marry boys in such a way that
each girl marries **a boy** that she knows?



The Marriage Problem (1935)

If there is a finite set of **girls**, each of whom knows several **boys**,
under what conditions can all the girls marry boys in such a way that
each girl marries **a boy** that she knows?



Philip Hall (1904 ~ 1982)






Definition (X -Perfect Matching (X -Saturating Matching))


Let $G = (X, Y, E)$ be a bipartite graph.

An X -perfect matching of G is a **matching** which covers each vertex in X .

$$|X| \leq |Y|$$



$$|X| \leq |Y|$$



$$\forall W \subseteq X. |W| \leq |N(W)|$$

Theorem (Hall Theorem; 1935)

Let $G = (X, Y, E)$ be a bipartite graph. There is a X -perfect matching of G iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$

By induction on the number $|X|$ of vertices in X .

By induction on the number $|X|$ of vertices in X .

Basis Step: $|X| = 1$. $|X| \leq |N_G(X)|$.

By induction on the number $|X|$ of vertices in X .

Basis Step: $|X| = 1$. $|X| \leq |N_G(X)|$. I am married!

Induction Hypothesis: Suppose that it holds if $|X| < m$.

By induction on the number $|X|$ of vertices in X .

Basis Step: $|X| = 1$. $|X| \leq |N_G(X)|$. I am married!

Induction Hypothesis: Suppose that it holds if $|X| < m$.

Induction Step: Consider the case $|X| = m$.

Consider the case $|X| = m$.

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

Consider the case $|X| = m$.

- ▶ CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

Consider the case $|X| = m$.

- CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

There is a $(X - \{x\})$ -perfect matching in G' .

Consider the case $|X| = m$.

- CASE I: Every $k < m$ girls in X know $\geq k + 1$ boys in Y .

Take any girl x and marry her to any boy y she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for G' .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

There is a $(X - \{x\})$ -perfect matching in G' .

Therefore, there is a $(X - \{x\})$ -perfect matching in G .

- ▶ CASE II: There is a set of $k < m$ girls in X who know k boys in Y .

Theorem (Hall Theorem; 1935)

Let $G = (X, Y, E)$ be a bipartite graph. There is a X -perfect matching of G iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$

By contradiction.

By contradiction.

Suppose that there is *no* X -perfect matching.

By contradiction.


Suppose that there is *no* X -perfect matching.

We show that Hall's Condition is violated for *some* $S \subseteq X$.

By contradiction.

Suppose that there is *no* X -perfect matching.


We show that Hall's Condition is violated for *some* $S \subseteq X$.



By contradiction.

Suppose that there is *no* X -perfect matching.

We show that Hall's Condition is violated for *some* $S \subseteq X$.




Let M be a *maximum* matching.

By contradiction.

Suppose that there is *no* X -perfect matching.

We show that Hall's Condition is violated for *some* $S \subseteq X$.




Let M be a *maximum* matching.

Let $u \in X$ be a vertex of X not saturated by M .

By contradiction.

Suppose that there is *no* X -perfect matching.


We show that Hall's Condition is violated for *some* $S \subseteq X$.




Let M be a *maximum* matching.

Let $u \in X$ be a vertex of X not saturated by M .

Consider all *M -alternating paths* starting from u .




$T \triangleq$ the set of vertices in Y reachable from u by M -alternating paths.



$T \triangleq$ the set of vertices in Y reachable from u by M -alternating paths.

$S \triangleq$ the set of vertices in X reachable from u by M -alternating paths.




$T \triangleq$ the set of vertices in Y reachable from u by M -alternating paths.

$S \triangleq$ the set of vertices in X reachable from u by M -alternating paths.

We will show that

$$T = N(S) \wedge |T| = |S - \{u\}|$$



$T \triangleq$ the set of vertices in Y reachable from u by M -alternating paths.


$S \triangleq$ the set of vertices in X reachable from u by M -alternating paths.

We will show that


$$T = N(S) \wedge |T| = |S - \{u\}|$$

$$|N(S)| = |T| = |S| - 1 < |S|$$

$$|T| = |S - \{u\}|$$




$$|T| = |S - \{u\}|$$




We show that there is a bijection from T to $S - \{u\}$.

$$|T| = |S - \{u\}|$$




We show that there is a bijection from T to $S - \{u\}$.
 M matches T with $S - \{u\}$.

$$T = N(S)$$




$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$


$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T$


$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u$$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u \implies s \in S - \{u\}$$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u \implies s \in S - \{u\} \implies \{s, y\} \notin M$$

$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

By contradiction: $N(S) \not\subseteq T \implies \exists y \in Y - T. y \in N(s)$

$$\exists y \in Y - T. \exists s \in S. \{s, y\} \in E$$

$$s \neq u \implies s \in S - \{u\} \implies \{s, y\} \notin M \implies y \in T$$

Theorem (Hall Theorem; 1935)

Let $G = (X, Y, E)$ be a bipartite graph. There is a X -perfect matching of G iff


$$\forall W \subseteq X. |W| \leq |N_G(W)|$$


algorithm

Definitions (Vertex Cover (点覆盖))

A **vertex cover** of a graph G is a set $Q \subseteq V(G)$ that **covers** all edges.

$$\forall e \in E(G). e \cap Q \neq \emptyset$$







examples


Theorem (König (1931), Egerváry (1931))

If G is a bipartite graph, then the maximum size of a matching in G equals the minimum size of a vertex cover of G






$$R = Q \cap X \quad T = Q \cap Y$$



$$R = Q \cap X \quad T = Q \cap Y$$

$H \triangleq (R \cup (Y - T))$ -induced subgraph of G

$H' \triangleq (T \cup (X - R))$ -induced subgraph of G




$$R = Q \cap X \quad T = Q \cap Y$$

$H \triangleq (R \cup (Y - T))$ -induced subgraph of G

$H' \triangleq (T \cup (X - R))$ -induced subgraph of G

G has no edges from $X - R$ to $Y - T$.



$$R = Q \cap X \quad T = Q \cap Y$$


$H \triangleq (R \cup (Y - T))$ -induced subgraph of G

$H' \triangleq (T \cup (X - R))$ -induced subgraph of G


G has no edges from $X - R$ to $Y - T$.

H has a R -perfect matching and H' has a T -perfect matching.

H has a R -perfect matching and H' has a T -perfect matching.




H has a R -perfect matching and H' has a T -perfect matching.



By contradiction.


H has a R -perfect matching and H' has a T -perfect matching.



By contradiction.

$$\exists S \subseteq R. |N_H(S)| < |S|$$

H has a R -perfect matching and H' has a T -perfect matching.




By contradiction.

$$\exists S \subseteq R. |N_H(S)| < |S|$$

$T \cup (R - S + N_H(S))$ is a smaller vertex cover than Q

Thank
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn