

# (十二) 图论: 匹配与网络流

## (Matching and Network Flow)

魏恒峰

hfwei@nju.edu.cn

2021 年 05 月 27 日





### 3 Theorems + 1 Algorithm

To **maximize** the size of a mathematical structure  $\mathcal{S}$  in  $G$



To maximize the size of a mathematical structure  $\mathcal{S}$  in  $G$



### Theorem

$\mathcal{S}$  is maximum iff  $G$  does not contain  $\mathcal{S}$ -augmenting objects.

To **maximize** the size of a mathematical structure  $\mathcal{S}$  in  $G$



### Theorem

$\mathcal{S}$  is maximum *iff*  $G$  does not contain  $\mathcal{S}$ -augmenting objects.

### Algorithm

Repeatedly finding  $\mathcal{S}$ -augmenting objects until no more ones exist.

To **maximize** the size of a mathematical structure  $\mathcal{S}$  in  $G$



To **maximize** the size of a mathematical structure  $\mathcal{S}$  in  $G$



To **minimize** the size of its **dual** mathematical structure  $\mathcal{S}'$  in  $G$

To **maximize** the size of a mathematical structure  $\mathcal{S}$  in  $G$



To **minimize** the size of its **dual** mathematical structure  $\mathcal{S}'$  in  $G$

Theorem (Weak Duality Theorem)

*The size of a maximum  $\mathcal{S} \leq$  The size of a minimum  $\mathcal{S}'$*



To **maximize** the size of a mathematical structure  $\mathcal{S}$  in  $G$



To **minimize** the size of its **dual** mathematical structure  $\mathcal{S}'$  in  $G$

Theorem (Weak Duality Theorem)

*The size of a maximum  $\mathcal{S} \leq$  The size of a minimum  $\mathcal{S}'$*

Theorem (Strong Duality Theorem)

*The size of a maximum  $\mathcal{S} =$  The size of a minimum  $\mathcal{S}'$*

let's get  
married  
today




## The Marriage Problem (Philip Hall, 1935)


If there is a finite set of **girls**, each of whom knows several **boys**,  
**under what conditions** can all the girls marry boys in such a way that  
**each girl** marries **a boy** that she knows?


## The Marriage Problem (Philip Hall, 1935)

If there is a finite set of **girls**, each of whom knows several **boys**,  
**under what conditions** can all the girls marry boys in such a way that  
**each girl** marries **a boy** that she knows?




Philip Hall (1904 ~ 1982)





### Definition (Matching (匹配))

A **matching** in a graph  $G$  is a set of edges with **no shared endpoints**.




### Definition ( $X$ -Perfect Matching ( $X$ -Saturating Matching))

Let  $G = (X, Y, E)$  be a bipartite graph.


An  $X$ -perfect matching of  $G$  is a matching which covers each vertex in  $X$ .

$$|X| \leq |Y|$$





$$|X| \leq |Y|$$



$$\forall W \subseteq X. |W| \leq |N(W)|$$

## Theorem (Hall Theorem; 1935)

Let  $G = (X, Y, E)$  be a bipartite graph. There is an  $X$ -perfect matching of  $G$  iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$

By induction on the number  $|X|$  of vertices in  $X$ .

By induction on the number  $|X|$  of vertices in  $X$ .

Basis Step:  $|X| = 1$ .  $|X| \leq |N_G(X)|$ .

By induction on the number  $|X|$  of vertices in  $X$ .

Basis Step:  $|X| = 1$ .  $|X| \leq |N_G(X)|$ . I am married!

Induction Hypothesis: Suppose that it holds if  $|X| < m$ .

By induction on the number  $|X|$  of vertices in  $X$ .

Basis Step:  $|X| = 1$ .  $|X| \leq |N_G(X)|$ . I am married!

Induction Hypothesis: Suppose that it holds if  $|X| < m$ .

Induction Step: Consider the case  $|X| = m$ .

Consider the case  $|X| = m$ .

Consider the case  $|X| = m$ .

- ▶ CASE I: Every  $k < m$  girls in  $X$  know  $\geq k + 1$  boys in  $Y$ .



Consider the case  $|X| = m$ .

- ▶ CASE I: Every  $k < m$  girls in  $X$  know  $\geq k + 1$  boys in  $Y$ .

Take any girl  $x$  and marry her to any boy  $y$  she knows.

Consider the case  $|X| = m$ .

- ▶ CASE I: Every  $k < m$  girls in  $X$  know  $\geq k + 1$  boys in  $Y$ .

Take any girl  $x$  and marry her to any boy  $y$  she knows.

$$G' = G - \{x, y\}$$

Consider the case  $|X| = m$ .

- ▶ CASE I: Every  $k < m$  girls in  $X$  know  $\geq k + 1$  boys in  $Y$ .

Take any girl  $x$  and marry her to any boy  $y$  she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for  $G'$ .

Consider the case  $|X| = m$ .

- ▶ CASE I: Every  $k < m$  girls in  $X$  know  $\geq k + 1$  boys in  $Y$ .

Take any girl  $x$  and marry her to any boy  $y$  she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for  $G'$ .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

Consider the case  $|X| = m$ .

- CASE I: Every  $k < m$  girls in  $X$  know  $\geq k + 1$  boys in  $Y$ .

Take any girl  $x$  and marry her to any boy  $y$  she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for  $G'$ .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

There is a  $(X - \{x\})$ -perfect matching in  $G'$ .

Consider the case  $|X| = m$ .

- CASE I: Every  $k < m$  girls in  $X$  know  $\geq k + 1$  boys in  $Y$ .

Take any girl  $x$  and marry her to any boy  $y$  she knows.

$$G' = G - \{x, y\}$$

The Hall's Condition still holds for  $G'$ .

$$\forall W \subseteq X - \{x\}. |W| \leq |N_{G'}(W)|$$

There is a  $(X - \{x\})$ -perfect matching in  $G'$ .

Therefore, there is a  $(X - \{x\})$ -perfect matching in  $G$ .

- ▶ CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

- ▶ CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

There  $k$  girls can be married **by induction** to the  $k$  boys.



- CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

There  $k$  girls can be married **by induction** to the  $k$  boys.

$$G' = G - \{\text{these } k \text{ girls}\} - \{\text{these } k \text{ boys}\}$$

- CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

There  $k$  girls can be married **by induction** to the  $k$  boys.

$$G' = G - \{\text{these } k \text{ girls}\} - \{\text{these } k \text{ boys}\}$$

$G'$  satisfies the Hall's Condition.

- ▶ CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

There  $k$  girls can be married **by induction** to the  $k$  boys.

$$G' = G - \{\text{these } k \text{ girls}\} - \{\text{these } k \text{ boys}\}$$

$G'$  satisfies the Hall's Condition.

**By contradiction:**

Suppose that in  $G'$ , there are  $l \leq m - k$  girls who know  $< l$  boys.

- CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

There  $k$  girls can be married **by induction** to the  $k$  boys.

$$G' = G - \{\text{these } k \text{ girls}\} - \{\text{these } k \text{ boys}\}$$

$G'$  satisfies the Hall's Condition.

**By contradiction:**

Suppose that in  $G'$ , there are  $l \leq m - k$  girls who know  $< l$  boys.

Then, in  $G$ , there are  $k + l$  girls who know  $< k + l$  boys.

- CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

There  $k$  girls can be married **by induction** to the  $k$  boys.

$$G' = G - \{\text{these } k \text{ girls}\} - \{\text{these } k \text{ boys}\}$$

$G'$  satisfies the Hall's Condition.

**By contradiction:**

Suppose that in  $G'$ , there are  $l \leq m - k$  girls who know  $< l$  boys.

Then, in  $G$ , there are  $k + l$  girls who know  $< k + l$  boys.

There is a  $(X - \{\text{these } k \text{ girls}\})$ -perfer matching in  $G'$ .

- CASE II: There is a set of  $k < m$  girls in  $X$  who know  $k$  boys in  $Y$ .

There  $k$  girls can be married **by induction** to the  $k$  boys.

$$G' = G - \{\text{these } k \text{ girls}\} - \{\text{these } k \text{ boys}\}$$

$G'$  satisfies the Hall's Condition.

**By contradiction:**

Suppose that in  $G'$ , there are  $l \leq m - k$  girls who know  $< l$  boys.

Then, in  $G$ , there are  $k + l$  girls who know  $< k + l$  boys.

There is a  $(X - \{\text{these } k \text{ girls}\})$ -perfer matching in  $G'$ .

Therefore, there is a  $(X - \{x\})$ -perfer matching in  $G$ .

### Theorem (Hall Theorem; 1935)

Let  $G = (X, Y, E)$  be a bipartite graph. There is a  $X$ -perfect matching of  $G$  *iff*

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$

## Definition ( $M$ -alternating Paths)

Let  $M$  be a matching. An  $M$ -alternating path is a path that alternates between edges in  $M$  and edges not in  $M$ .





### Definition ( $M$ -alternating Paths)

Let  $M$  be a matching. An  $M$ -alternating path is a path that alternates between edges in  $M$  and edges not in  $M$ .



### Definition ( $M$ -augmenting Paths)

An  $M$ -augmenting path is an  $M$ -alternating path whose endpoints are unsaturated by  $M$ .

By contradiction.

By contradiction.

Suppose that there is *no*  $X$ -perfect matching.

By contradiction.


Suppose that there is *no*  $X$ -perfect matching.

We show that Hall's Condition is violated for *some*  $S \subseteq X$ .

By contradiction.

Suppose that there is *no*  $X$ -perfect matching.


We show that Hall's Condition is violated for *some*  $S \subseteq X$ .



By contradiction.

Suppose that there is *no*  $X$ -perfect matching.

We show that Hall's Condition is violated for *some*  $S \subseteq X$ .




Let  $M$  be a *maximum* matching.

By contradiction.

Suppose that there is *no*  $X$ -perfect matching.

We show that Hall's Condition is violated for *some*  $S \subseteq X$ .




Let  $M$  be a *maximum* matching.

Let  $u \in X$  be a vertex of  $X$  not saturated by  $M$ .

By contradiction.

Suppose that there is *no*  $X$ -perfect matching.

We show that Hall's Condition is violated for *some*  $S \subseteq X$ .




Let  $M$  be a *maximum* matching.


Let  $u \in X$  be a vertex of  $X$  not saturated by  $M$ .

Consider all  *$M$ -alternating paths* starting from  $u$ .






$T \triangleq$  the set of vertices in  $Y$  reachable from  $u$  by  $M$ -alternating paths.



$T \triangleq$  the set of vertices in  $Y$  reachable from  $u$  by  $M$ -alternating paths.

$S \triangleq$  the set of vertices in  $X$  reachable from  $u$  by  $M$ -alternating paths.




$T \triangleq$  the set of vertices in  $Y$  reachable from  $u$  by  $M$ -alternating paths.

$S \triangleq$  the set of vertices in  $X$  reachable from  $u$  by  $M$ -alternating paths.

We will show that

$$T = N(S) \wedge |T| = |S - \{u\}|$$



$T \triangleq$  the set of vertices in  $Y$  reachable from  $u$  by  $M$ -alternating paths.


$S \triangleq$  the set of vertices in  $X$  reachable from  $u$  by  $M$ -alternating paths.

We will show that


$$T = N(S) \wedge |T| = |S - \{u\}|$$

$$|N(S)| = |T| = |S| - 1 < |S|$$

$$|T| = |S - \{u\}|$$




$$|T| = |S - \{u\}|$$



We show that there is a bijection from  $T$  to  $S - \{u\}$ .


$$|T| = |S - \{u\}|$$



We show that there is a bijection from  $T$  to  $S - \{u\}$ .


$M$  matches  $T$  with  $S - \{u\}$ .

$$T = N(S)$$






$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$


$$T = N(S)$$



$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

We need to show that  $N(S) \subseteq T$

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

We need to show that  $N(S) \subseteq T$

Consider the neighbors of  $x \in S$ :

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$

We need to show that  $N(S) \subseteq T$

Consider the neighbors of  $x \in S$ :  $x = u$ ,

$$T = N(S)$$




$$|T| = |S - \{u\}| \implies T \subseteq N(S)$$



We need to show that  $N(S) \subseteq T$




Consider the neighbors of  $x \in S$ :  $x = u$ ,  $x \in S - \{u\}$

## Definitions (Vertex Cover (点覆盖))

A **vertex cover** of a graph  $G$  is a set  $Q \subseteq V(G)$  that **covers** all edges.










## Theorem (Weak Duality Theorem (弱对偶定理))

*Let  $G$  be a graph. The maximum size of a matching in  $G$   
 $\leq$  the minimum size of a vertex cover of  $G$ .*



## Theorem (Weak Duality Theorem (弱对偶定理))

Let  $G$  be a graph. The maximum size of a matching in  $G$   
 $\leq$  the minimum size of a vertex cover of  $G$ .




## Theorem (König (1931), Egerváry (1931))


Let  $G$  be a *bipartite* graph. The maximum size of a matching in  $G$   
*equals* the minimum size of a vertex cover of  $G$

Given a vertex cover  $Q$ , we construct a matching  $M$  of the same size.

Given a vertex cover  $Q$ , we construct a matching  $M$  of the same size.




Given a vertex cover  $Q$ , we construct a matching  $M$  of the same size.



$$R = Q \cap X \quad T = Q \cap Y$$

Given a vertex cover  $Q$ , we construct a matching  $M$  of the same size.




$$R = Q \cap X \quad T = Q \cap Y$$

$H \triangleq (R \cup (Y - T))$ -induced subgraph of  $G$

$H' \triangleq (T \cup (X - R))$ -induced subgraph of  $G$

Given a vertex cover  $Q$ , we construct a matching  $M$  of the same size.



$$R = Q \cap X \quad T = Q \cap Y$$


$H \triangleq (R \cup (Y - T))$ -induced subgraph of  $G$

$H' \triangleq (T \cup (X - R))$ -induced subgraph of  $G$

$G$  has no edges from  $X - R$  to  $Y - T$ .



Given a vertex cover  $Q$ , we construct a matching  $M$  of the same size.



$$R = Q \cap X \quad T = Q \cap Y$$


$H \triangleq (R \cup (Y - T))$ -induced subgraph of  $G$

$H' \triangleq (T \cup (X - R))$ -induced subgraph of  $G$


$G$  has no edges from  $X - R$  to  $Y - T$ .

$H$  has a  $R$ -perfect matching and  $H'$  has a  $T$ -perfect matching.

$H$  has a  $R$ -perfect matching and  $H'$  has a  $T$ -perfect matching.




$H$  has a  $R$ -perfect matching and  $H'$  has a  $T$ -perfect matching.



By contradiction.


$H$  has a  $R$ -perfect matching and  $H'$  has a  $T$ -perfect matching.



By contradiction.

$$\exists S \subseteq R. |N_H(S)| < |S|$$

$H$  has a  $R$ -perfect matching and  $H'$  has a  $T$ -perfect matching.



By contradiction.


$$\exists S \subseteq R. |N_H(S)| < |S|$$

$T \cup (R - S + N_H(S))$  is a smaller vertex cover than  $Q$

## Definition (Network (网络))


A **network** is a **digraph** with

- ▶ a distinguished **source vertex**  $s$ ,
- ▶ a distinguished **sink vertex**  $t$ ,
- ▶ a **capacity**  $c(e) \geq 0$  on each edge  $e$



## Definition (Flow (流))

A **flow**  $f$  is a **function** that assigns a value  $f(e)$  to each edge  $e$ .



## Definition (Feasible Flow (可行流))


A flow  $f$  is **feasible** if it satisfies

Capacity Constraints:

$$\forall e \in E. 0 \leq f(e) \leq c(e)$$


Flow Conservation:

$$\forall v \in V. f^+(v) = f^-(v)$$






$$f^+(v) = \sum_{(v,w) \in E} f(v,w) \quad f^-(v) = \sum_{(u,v) \in E} f(u,v)$$




$$f^+(v) = \sum_{(v,w) \in E} f(v,w) \quad f^-(v) = \sum_{(u,v) \in E} f(u,v)$$





$$f^+(U) = \sum_{u \in U, v \in \overline{U}, (u,v) \in E} f(u,v)$$

$$f^+(v) = \sum_{(v,w) \in E} f(v,w) \quad f^-(v) = \sum_{(u,v) \in E} f(u,v)$$




$$f^+(U) = \sum_{u \in U, v \in \bar{U}, (u,v) \in E} f(u,v) \quad f^-(U) = \sum_{v \in \bar{U}, u \in U, (v,u) \in E} f(v,u)$$






$$\forall U \subseteq (V - \{s, t\}). f^+(U) = f^-(U)$$



$$\forall U \subseteq (V - \{s, t\}). f^+(U) = f^-(U)$$

$$s \in U \wedge t \notin U \implies f^+(U) - f^-(U) =$$




$$\forall U \subseteq (V - \{s, t\}). f^+(U) = f^-(U)$$

$$s \in U \wedge t \notin U \implies f^+(U) - f^-(U) = f^+(s)$$

## Definition (Value (值))

The **value**  $\text{val}(f)$  of a **flow**  $f$  is

$$\text{val}(f) = f^-(t) = f^+(s).$$







## Definition (Value (值))

The **value**  $\text{val}(f)$  of a **flow**  $f$  is



$$\text{val}(f) = f^-(t) = f^+(s).$$



## Definition (Value (值))

The **value**  $\text{val}(f)$  of a **flow**  $f$  is



$$\text{val}(f) = f^-(t) = f^+(s).$$





## Definition (Maximum Flow (最大流))

A **maximum flow** is a **feasible flow** of maximum **value**.

$$s - x - v - t$$



$$s - x - v - t$$



### Definition ( $f$ -augmenting Paths (增广路径))



When  $f$  is a feasible flow, an  **$f$ -augmenting path** is a  $s \sim t$  path  $P$  in the underlying graph such that for each edge  $e \in E(P)$ ,

- (a) if  $P$  follows  $e$  in the forward direction, then  $f(e) < c(e)$ ;
- (b) if  $P$  follows  $e$  in the backward direction, then  $f(e) > 0$ .

## Definition ( $f$ -augmenting Paths)

Let  $P$  be an  $f$ -augmenting path.



$$\epsilon(e) = \begin{cases} c(e) - f(e) & \text{if } e \text{ is forward on } P \\ f(e) & \text{if } e \text{ is backward on } P \end{cases}$$



## Definition ( $f$ -augmenting Paths)


Let  $P$  be an  $f$ -augmenting path.



$$\epsilon(e) = \begin{cases} c(e) - f(e) & \text{if } e \text{ is forward on } P \\ f(e) & \text{if } e \text{ is backward on } P \end{cases}$$



An  $f$ -augmenting path leads to a flow with **larger** value.

$$\min_{e \in E(P)} \epsilon(e)$$









## Definition (Source/Sink Cut (割))

In a network, a **source/sink cut**  $[S, T]$  consists of the edges **from** a **source set**  $S$  **to** a **sink set**  $T$ , where


$$(T = V - S) \wedge (s \in S) \wedge (t \in T)$$






Definition (Capacity of Cut (割的容量))


$$\text{cap}(S, T) = \sum_{u \in S, v \in T, uv \in E} c(u, v)$$






Definition (Minimum Cut (最小割))

A **minimum cut** is a **cut** of minimum value.



Definition (Minimum Cut (最小割))


A **minimum cut** is a **cut** of minimum value.



## Theorem (Weak Duality (弱对偶定理))

Let  $f$  be any feasible *flow* and  $[S, T]$  be any source/sink *cut*.


$$\text{val}(f) \leq \text{cap}(S, T).$$



## Theorem (Weak Duality (弱对偶定理))

Let  $f$  be any feasible *flow* and  $[S, T]$  be any source/sink *cut*.

$$\text{val}(f) \leq \text{cap}(S, T).$$




$$\text{val}(f) = f^+(S) - f^-(S)$$

## Theorem (Weak Duality (弱对偶定理))

Let  $f$  be any feasible *flow* and  $[S, T]$  be any source/sink *cut*.

$$\text{val}(f) \leq \text{cap}(S, T).$$




$$\text{val}(f) = f^+(S) - f^-(S) \leq f^+(S)$$



## Theorem (Weak Duality (弱对偶定理))

Let  $f$  be any feasible *flow* and  $[S, T]$  be any source/sink *cut*.

$$\text{val}(f) \leq \text{cap}(S, T).$$



$$\text{val}(f) = f^+(S) - f^-(S) \leq f^+(S) \leq \text{cap}(S, T)$$

## Lemma

$$\max_f \text{val}(f) \leq \min_{[S,T]} \text{cap}(S, T)$$

## Lemma

$$\max_f \text{val}(f) \leq \min_{[S,T]} \text{cap}(S, T)$$

What if  $\text{val}(f) = \text{cap}(S, T)$  for some flow  $f$  and some cut  $[S, T]$ ?

## Lemma

$$\max_f \text{val}(f) \leq \min_{[S,T]} \text{cap}(S, T)$$

What if  $\text{val}(f) = \text{cap}(S, T)$  for some flow  $f$  and some cut  $[S, T]$ ?

$f$  is maximum and  $[S, T]$  is minimum


$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$f^-(S) = 0 \wedge f^+(S) = \text{cap}(S, T)$$


$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$f^-(S) = 0 \wedge f^+(S) = \text{cap}(S, T)$$



$$\text{val}(f) = f^+(S) - f^-(S) = f^+(S) = \text{cap}(S, T)$$

$$f^-(S) = 0 \wedge f^+(S) = \text{cap}(S, T)$$





## Theorem (Max-flow Min-cut Theorem (Ford and Fulkerson; 1956))

$$\max_f \text{val}(f) = \min_{[S,T]} \text{cap}(S,T)$$


*(Strong Duality)*





L. R. Ford Jr. (1927 ~ 2017)



D. R. Fulkerson (1924 ~ 1976)





## Theorem

*A feasible flow  $f$  is maximum iff there are no  $f$ -augmenting paths.*

## Theorem


*A feasible flow  $f$  is maximum iff there are no  $f$ -augmenting paths.*

We construct a cut  $[S, T]$  with  $\text{val}(f) = \text{cap}(S, T)$ .

## Theorem

*A feasible flow  $f$  is maximum iff there are no  $f$ -augmenting paths.*


We construct a cut  $[S, T]$  with  $\text{val}(f) = \text{cap}(S, T)$ .



$S \triangleq \{\text{the vertices reachable from } s \text{ along } \textcolor{red}{\text{partial}} \text{ } f\text{-augmenting paths}\}$


$S \triangleq \{\text{the vertices reachable from } s \text{ along partial } f\text{-augmenting paths}\}$

$$T \triangleq V - S$$



$S \triangleq \{\text{the vertices reachable from } s \text{ along partial } f\text{-augmenting paths}\}$

$$T \triangleq V - S$$




$$f^{-1}(S) = 0 \wedge f^{+}(S) = \text{cap}(S, T)$$



$S \triangleq \{\text{the vertices reachable from } s \text{ along partial } f\text{-augmenting paths}\}$

$$T \triangleq V - S$$



$$f^{-1}(S) = 0 \wedge f^{+}(S) = \text{cap}(S, T)$$


$$\text{val}(f) = f^{+}(S) - f^{-}(S) = f^{+}(S) = \text{cap}(S, T)$$

## The Ford-Fulkerson Method

Repeatedly finding  $f$ -augmenting paths until no more ones exist.


## The Ford-Fulkerson Method

Repeatedly finding  $f$ -augmenting paths until no more ones exist.



## The Ford-Fulkerson Method

Repeatedly finding  $f$ -augmenting paths until no more ones exist.




## The Edmonds-Karp Algorithm

Using **BFS** (Breadth-first Search) to find  $f$ -augmenting paths.


## The Edmonds-Karp Algorithm

Using **BFS** (Breadth-first Search) to find  $f$ -augmenting paths.



## The Edmonds-Karp Algorithm

Using **BFS** (Breadth-first Search) to find  $f$ -augmenting paths.



## Theorem (Hall Theorem; 1935)

There is an  *$X$ -perfect matching* of  $G$  iff


$$\forall W \subseteq X. |W| \leq |N_G(W)|$$



Theorem (Hall Theorem; 1935)

There is an *X*-perfect matching of  $G$  iff

$$\forall W \subseteq X. |W| \leq |N_G(W)|$$




$$\forall x \in X. c(s, x) = 1 \quad \forall y \in Y. c(y, t) = 1 \quad \forall x \in X, y \in Y. c(x, y) = \infty$$

Theorem (Hall Theorem; 1935)

There is an  $X$ -perfect matching of  $G$  iff


$$\forall W \subseteq X. |W| \leq |N_G(W)|$$




$$\forall x \in X. c(s, x) = 1 \quad \forall y \in Y. c(y, t) = 1 \quad \forall x \in X, y \in Y. c(x, y) = \infty$$

We need to show that  $\max_f \text{val}(f) = |X|$ .

We need to show that  $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) = |X|$ .




We need to show that  $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) = |X|$ .



$$\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) \leq |X|$$


We need to show that  $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) = |X|$ .



$$\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) \leq |X|$$


Therefore, we need to show that  $\min_{[S, \bar{S}]} \text{cap}(S, \bar{S}) \geq |X|$ .

Let  $[S, \bar{S}]$  be a cut. We need to show that  $\text{cap}(S, \bar{S}) \geq |X|$ .



$$N(S \cap X) \not\subseteq (S \cap Y)$$


Let  $[S, \bar{S}]$  be a cut. We need to show that  $\text{cap}(S, \bar{S}) \geq |X|$ .



$$N(S \cap X) \not\subseteq (S \cap Y)$$


$$\text{cap}(S, \bar{S}) = \infty \geq |X|$$

$$N(S \cap X) \subseteq (S \cap Y)$$






$$N(S \cap X) \subseteq (S \cap Y)$$




$$\text{cap}(S, \bar{S}) = \sum_{u \in S, v \in \bar{S}} c(u, v)$$

$$N(S \cap X) \subseteq (S \cap Y)$$




$$\begin{aligned} \text{cap}(S, \bar{S}) &= \sum_{u \in S, v \in \bar{S}} c(u, v) \\ &= \sum_{v \in \bar{S} \cap X} c(s, v) + \sum_{u \in S \cap Y} c(u, t) \end{aligned}$$

$$N(S \cap X) \subseteq (S \cap Y)$$




$$\begin{aligned}
 \text{cap}(S, \bar{S}) &= \sum_{u \in S, v \in \bar{S}} c(u, v) \\
 &= \sum_{v \in \bar{S} \cap X} c(s, v) + \sum_{u \in S \cap Y} c(u, t) \\
 &= |X| - |S \cap X| + |S \cap Y|
 \end{aligned}$$

$$N(S \cap X) \subseteq (S \cap Y)$$



$$\begin{aligned}
 \text{cap}(S, \bar{S}) &= \sum_{u \in S, v \in \bar{S}} c(u, v) \\
 &= \sum_{v \in \bar{S} \cap X} c(s, v) + \sum_{u \in S \cap Y} c(u, t) \\
 &= |X| - |S \cap X| + |S \cap Y| \\
 &\geq |X| - |S \cap X| + |N(S \cap X)|
 \end{aligned}$$

$$N(S \cap X) \subseteq (S \cap Y)$$



$$\begin{aligned}
 \text{cap}(S, \bar{S}) &= \sum_{u \in S, v \in \bar{S}} c(u, v) \\
 &= \sum_{v \in \bar{S} \cap X} c(s, v) + \sum_{u \in S \cap Y} c(u, t) \\
 &= |X| - |S \cap X| + |S \cap Y| \\
 &\geq |X| - |S \cap X| + |N(S \cap X)| \geq |X|
 \end{aligned}$$

## Theorem (König (1931), Egerváry (1931))

*If  $G$  is a bipartite graph, then the maximum size of a matching in  $G$  equals the minimum size of a vertex cover of  $G$*


Theorem (König (1931), Egerváry (1931))

*If  $G$  is a bipartite graph, then the maximum size of a matching in  $G$  equals the minimum size of a vertex cover of  $G$*



Thank  
You!





Office 302

Mailbox: H016

hfwei@nju.edu.cn