(十五) 离散数学: 复习 (Review)

魏恒峰

hfwei@nju.edu.cn

2021 年 06 月 17 日

REVIEW

Back to basics

LOGICAL/
MATHEMATICAL

$$\vdash \qquad \models$$

Theorem

$$\Sigma \vdash \alpha \Longleftrightarrow \Sigma \models \alpha$$

$$\rightarrow \qquad \Longrightarrow$$

$$\leftrightarrow \qquad \Longleftrightarrow$$

$$\rightarrow \qquad \Longrightarrow$$

$$\leftrightarrow \qquad \Longleftrightarrow$$

"$\rightarrow$" and "$\leftrightarrow$" are used in a single formula.

" $\Longrightarrow$ " and " $\Longleftrightarrow$ " are used to connect two formulas.

$$\rightarrow \qquad \Longrightarrow$$

$$\leftrightarrow \qquad \Longleftrightarrow$$

"$\rightarrow$" and "$\leftrightarrow$" are used in a single formula.

" $\Longrightarrow$ " and " $\Longleftrightarrow$ " are used to connect two formulas.

$$\begin{aligned}
& x \in A \setminus B \\
\Longleftrightarrow\ & x \in A \wedge x \notin B \\
\Longleftrightarrow\ & x \in A \wedge (x \in U \wedge x \notin B) \\
\Longleftrightarrow\ & x \in A \wedge x \in \overline{B} \\
\Longleftrightarrow\ & x \in A \cap \overline{B}
\end{aligned}$$

$$
\begin{aligned}
p \oplus q &\triangleq (p \vee q) \wedge \neg(p \wedge q) \\
&= (p \wedge \neg q) \vee (\neg q \wedge q)
\end{aligned}
$$

p	q	$p \oplus q$
0	0	0
0	1	1
1	0	1
1	1	0

$$
\begin{aligned}
p \oplus q &\triangleq (p \vee q) \wedge \neg(p \wedge q) \\
&= (p \wedge \neg q) \vee (\neg q \wedge q)
\end{aligned}
$$

p	q	$p \oplus q$
0	0	0
0	1	1
1	0	1
1	1	0

$$p \oplus q = q \oplus r$$

$$(p \oplus q) \oplus r = p \oplus (q \oplus r)$$

Theorem

$\sqrt{2}$ is irrational.

The First Crisis in Mathematics

Theorem (Pigeonhole Principle)

If n objects are placed in r boxes, where $r < n$, then at least one of the boxes contains ≥ 2 ($\geq \lceil \frac{n}{r} \rceil$) object.

Numbers

Consider the numbers $1, 2, \ldots, 2n$, and take any $n+1$ of them. There are two among these $n+1$ numbers which are relatively prime.

Numbers

Consider the numbers $1, 2, \ldots, 2n$, and take any $n+1$ of them. There are two among these $n+1$ numbers which are relatively prime.

There must be two numbers which are only 1 apart.

Numbers

Consider the numbers $1, 2, \ldots, 2n$, and take any $n+1$ of them. There are two among these $n+1$ numbers such as one divides the other.

Numbers

Consider the numbers $1, 2, \ldots, 2n$, and take any $n+1$ of them. There are two among these $n+1$ numbers such as one divides the other.

$$a = 2^k m, \quad (1 \leq m \leq 2n-1 \text{ is odd})$$

There $n+1$ numbers have only n different odd parts.

There must be two numbers with the same odd part.

Hand-shaking

If there are $n > 1$ people who can shake hands with one another, there are two people who shake hands with the same number of people.

Hand-shaking

If there are $n > 1$ people who can shake hands with one another, there are two people who shake hands with the same number of people.

$$0 \sim n - 1$$

Hand-shaking

If there are $n > 1$ people who can shake hands with one another, there are two people who shake hands with the same number of people.

$$0 \sim n - 1$$

Either the '0' hole or the 'n − 1' hole or both must be empty.

Sums

Suppose we are given n integers $a_1, a_2, \ldots, a_n$.
Then there is a set of consecutive numbers $a_{k+1}, a_{k+2}, \ldots, a_l$ whose sum $\sum_{i=k+1}^{l} a_i$ is a multiple of n.

Sums

Suppose we are given n integers $a_1, a_2, \ldots, a_n$.
Then there is a set of consecutive numbers $a_{k+1}, a_{k+2}, \ldots, a_l$ whose sum $\sum_{i=k+1}^{l} a_i$ is a multiple of n.

$$A_i = \sum_{k=1}^{k=i} a_i$$

Sums

Suppose we are given n integers $a_1, a_2, \ldots, a_n$.
Then there is a set of consecutive numbers $a_{k+1}, a_{k+2}, \ldots, a_l$ whose sum $\sum_{i=k+1}^{l} a_i$ is a multiple of n.

$$A_i = \sum_{k=1}^{k=i} a_i$$

$$A_0, A_1, A_2, \ldots, A_n$$

Sums

Suppose we are given n integers $a_1, a_2, \ldots, a_n$.
Then there is a set of consecutive numbers $a_{k+1}, a_{k+2}, \ldots, a_l$ whose sum $\sum_{i=k+1}^{l} a_i$ is a multiple of n.

$$A_i = \sum_{k=1}^{k=i} a_i$$

$$A_0, A_1, A_2, \ldots, A_n$$

$$\exists 0 \leq i < j \leq n.\ A_i = A_j \mod n$$

Sums

Suppose we are given n integers $a_1, a_2, \ldots, a_n$.
Then there is a set of consecutive numbers $a_{k+1}, a_{k+2}, \ldots, a_l$ whose sum $\sum_{i=k+1}^{l} a_i$ is a multiple of n.

$$A_i = \sum_{k=1}^{k=i} a_i$$

$$A_0, A_1, A_2, \ldots, A_n$$

$$\exists 0 \leq i < j \leq n.\ A_i = A_j \mod n$$

$$A_j - A_i = a_{i+1} + \cdots + a_j = 0 \mod n$$

Championship Match

“胡司令” (胡荣华) 要安排一次长达 77 天的象棋练习赛。

他想每天至少要有一场比赛, 但是总共不超过 132 场比赛。

请证明, 无论如何安排, 他都要在连续的若干天内恰好完成 21 场比赛。

Championship Match

“胡司令” (胡荣华) 要安排一次长达 77 天的象棋练习赛。

他想每天至少要有一场比赛, 但是总共不超过 132 场比赛。

请证明, 无论如何安排, 他都要在连续的若干天内恰好完成 21 场比赛。

Let a_i denote the number of games he plays up through the i-th day.

Championship Match

“胡司令” (胡荣华) 要安排一次长达 77 天的象棋练习赛。

他想每天至少要有一场比赛, 但是总共不超过 132 场比赛。

请证明, 无论如何安排, 他都要在连续的若干天内恰好完成 21 场比赛。

Let a_i denote the number of games he plays up through the i-th day.

$$a_1, a_2, \ldots, a_{76}, a_{77}, a_1 + 21, a_2 + 21, \ldots, a_{76} + 21, a_{77} + 21$$

Championship Match

“胡司令” (胡荣华) 要安排一次长达 77 天的象棋练习赛。

他想每天至少要有一场比赛, 但是总共不超过 132 场比赛。

请证明, 无论如何安排, 他都要在连续的若干天内恰好完成 21 场比赛。

Let a_i denote the number of games he plays up through the i-th day.

$$a_1, a_2, \ldots, a_{76}, a_{77}, a_1 + 21, a_2 + 21, \ldots, a_{76} + 21, a_{77} + 21$$

There must be ≥ 2 elements having the same value.

Championship Match

“胡司令” (胡荣华) 要安排一次长达 77 天的象棋练习赛。

他想每天至少要有一场比赛, 但是总共不超过 132 场比赛。

请证明, 无论如何安排, 他都要在连续的若干天内恰好完成 21 场比赛。

Let a_i denote the number of games he plays up through the i-th day.

$$a_1, a_2, \ldots, a_{76}, a_{77}, a_1 + 21, a_2 + 21, \ldots, a_{76} + 21, a_{77} + 21$$

There must be ≥ 2 elements having the same value.

It must be $a_i + 21 = a_j$.

Sequences

In any sequence $a_1, a_2, \ldots, a_{mn+1}$ of $mn+1$ distinct numbers, there exists an increasing subsequence

$$a_{i_1} < a_{i_2} < \cdots < a_{i_{m+1}} \quad (i_1 < i_2 < \cdots < i_{m+1})$$

of length $m+1$, or a decreasing subsequence

$$a_{j_1} > a_{j_2} > \cdots > a_{j_{n+1}} \quad (j_1 > i_2 < \cdots > j_{n+1})$$

of length $n+1$, or both.

Paul Erdős (1913 ∼ 1996)

Chapter 28 of “Proofs from THE Book”

$$|A \cup B| = |A| + |B| - |A \cap B|$$

$$|A \cup B| = |A| + |B| - |A \cap B|$$

$$\begin{aligned} |A \cup B \cup C| = {} & |A| + |B| + |C| \\ & - |A \cap B| - |A \cap C| - |B \cap C| \\ & + |A \cap B \cap C| \end{aligned}$$

Theorem (Inclusion-Exclusion Principle)

$$\left|\bigcup_{i=1}^{n} A_i\right| = \sum_{i=1}^{n} |A_i| - \sum_{1\leqslant i<j\leqslant n} |A_i \cap A_j|$$

$$+ \sum_{1\leqslant i<j<k\leqslant n} |A_i \cap A_j \cap A_k|$$

$$- \cdots$$

$$+ (-1)^{n-1} |A_1 \cap \cdots \cap A_n| .$$

Theorem (Inclusion-Exclusion Principle)

$$\left|\bigcup_{i=1}^{n} A_i\right| = \sum_{i=1}^{n} |A_i| - \sum_{1\leqslant i<j\leqslant n} |A_i \cap A_j|$$

$$+ \sum_{1\leqslant i<j<k\leqslant n} |A_i \cap A_j \cap A_k|$$

$$- \cdots$$

$$+ (-1)^{n-1} |A_1 \cap \cdots \cap A_n| .$$

$$\left|\bigcap_{i=1}^{n} \bar{A}_i\right| = \left|S - \bigcup_{i=1}^{n} A_i\right| = |S| - \sum_{i=1}^{n} |A_i| + \sum_{1\leqslant i<j\leqslant n} |A_i \cap A_j|$$

$$- \cdots + (-1)^n |A_1 \cap \cdots \cap A_n|.$$

Counting Integers

How many integers in $1, \ldots, 100$ are not divisible by 2, 3 or 5?

Counting Integers

How many integers in $1, \ldots, 100$ are not divisible by 2, 3 or 5?

$$100 - (50 + 33 + 20) + (16 + 10 + 6) - 3 = 26.$$

Counting Derangements (错排)

Suppose there is a deck of n cards numbered from 1 to n. Suppose a card numbered i is in the correct position if it is the i-th card in the deck. How many ways can the cards be shuffled without any cards being in the correct position?

Counting Derangements (错排)

Suppose there is a deck of n cards numbered from 1 to n. Suppose a card numbered i is in the correct position if it is the i-th card in the deck. How many ways can the cards be shuffled without any cards being in the correct position?

A_m : all of the orderings of cards with the m-th card correct

Counting Derangements (错排)

Suppose there is a deck of n cards numbered from 1 to n. Suppose a card numbered i is in the correct position if it is the i-th card in the deck. How many ways can the cards be shuffled without any cards being in the correct position?

A_m : all of the orderings of cards with the m-th card correct

$$\left|\bigcap_{i=1}^{n} \overline{A_i}\right| = \left|S - \bigcup_{i=1}^{n} A_i\right| = n! - \sum_{i=1}^{n} |A_i| + \sum_{1 \leqslant i < j \leqslant n} |A_i \cap A_j|$$

$$- \cdots + (-1)^n |A_1 \cap \cdots \cap A_n|.$$

Counting Derangements (错排)

Suppose there is a deck of n cards numbered from 1 to n. Suppose a card numbered i is in the correct position if it is the i-th card in the deck. How many ways can the cards be shuffled without any cards being in the correct position?

A_m : all of the orderings of cards with the m-th card correct

$$\left|\bigcap_{i=1}^{n} \overline{A_i}\right| = \left|S - \bigcup_{i=1}^{n} A_i\right| = n! - \sum_{i=1}^{n} |A_i| + \sum_{1 \leqslant i < j \leqslant n} |A_i \cap A_j|$$

$$- \cdots + (-1)^n |A_1 \cap \cdots \cap A_n|.$$

$$S_k \triangleq \sum_{1 \leq i_1 < i_2 < \cdots < i_k \leq n} |A_{i_1} \cap A_{i_2} \cap \cdots \cap A_{i_k}| =$$

Counting Derangements (错排)

Suppose there is a deck of n cards numbered from 1 to n. Suppose a card numbered i is in the correct position if it is the i-th card in the deck. How many ways can the cards be shuffled without any cards being in the correct position?

A_m : all of the orderings of cards with the m-th card correct

$$\left|\bigcap_{i=1}^{n} \overline{A_i}\right| = \left|S - \bigcup_{i=1}^{n} A_i\right| = n! - \sum_{i=1}^{n} |A_i| + \sum_{1 \leqslant i < j \leqslant n} |A_i \cap A_j|$$

$$- \cdots + (-1)^n |A_1 \cap \cdots \cap A_n|.$$

$$S_k \triangleq \sum_{1 \leq i_1 < i_2 < \cdots < i_k \leq n} |A_{i_1} \cap A_{i_2} \cap \cdots \cap A_{i_k}| = \binom{n}{k}(n-k)! = \frac{n!}{k!}$$

$$S_k = \frac{n!}{k!}$$

$$\left|\bigcap_{i=1}^{n} \overline{A_i}\right| = n! - \frac{n!}{1!} + \frac{n!}{2!} - \cdots + (-1)^n \frac{n!}{n!}$$

$$= n! \sum_{k=0}^{n} \frac{(-1)^k}{k!}$$

$$S_k = \frac{n!}{k!}$$

$$\left|\bigcap_{i=1}^{n} \overline{A_i}\right| = n! - \frac{n!}{1!} + \frac{n!}{2!} - \cdots + (-1)^n \frac{n!}{n!}$$

$$= n! \sum_{k=0}^{n} \frac{(-1)^k}{k!}$$

$$n \to \infty \implies \sum_{k=0}^{n} \frac{(-1)^k}{k!} \to e^{-1} \approx 0.368$$

$$\boxed{a_n = f(a_{n-1}, a_{n-2}, \ldots, a_{n-t}) + g(n)}$$

$$\boxed{a_n = f(a_{n-1}, a_{n-2}, \ldots, a_{n-t}) + g(n)}$$

recurrence type	typical example
first-order	
linear	$a_n = na_{n-1} - 1$
nonlinear	$a_n = 1/(1 + a_{n-1})$
second-order	
linear	$a_n = a_{n-1} + 2a_{n-2}$
nonlinear	$a_n = a_{n-1}a_{n-2} + \sqrt{a_{n-2}}$
variable coefficients	$a_n = na_{n-1} + (n-1)a_{n-2} + 1$
tth order	$a_n = f(a_{n-1}, a_{n-2}, \ldots, a_{n-t})$
full-history	$a_n = n + a_{n-1} + a_{n-2} \ldots + a_1$
divide-and-conquer	$a_n = a_{\lfloor n/2 \rfloor} + a_{\lceil n/2 \rceil} + n$

Table 2.1 Classification of recurrences

Homogeneous Linear Recurrence Relations with Constant Coefficients

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_t a_{n-t}$$

Homogeneous Linear Recurrence Relations with Constant Coefficients

$$a_n = c_1 a_{n-1} + c_2 a_{n-2} + \cdots + c_t a_{n-t}$$

https://www.bilibili.com/video/BV1Cf4y187Cu?share_source=copy_web

$$R \subseteq A \times A$$

$$\begin{cases} R^0 = I_A \\ R^{n+1} = R \circ R^n \end{cases}$$

Representing Relations as Matrices/Digraphs

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1,1), (1,2), (2,1), (2,2), (2,3), (2,4), (3,4), (4,1)\}$$

Representing Relations as Matrices/Digraphs

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1,1), (1,2), (2,1), (2,2), (2,3), (2,4), (3,4), (4,1)\}$$

$$R^2 \qquad R^3$$

Representing Relations as Matrices/Digraphs

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1,1), (1,2), (2,1), (2,2), (2,3), (2,4), (3,4), (4,1)\}$$

$$R^2 \qquad R^3$$

$$R^+ = \bigcup_{i=1}^{\infty} R \qquad R^* = \bigcup_{i=0}^{\infty} R$$

Definition (Reflexive Closure (自反闭包))

The reflexive closure $\mathsf{cl}_{\mathrm{ref}}(R)$ of a relation $R \subseteq X \times X$ is the smallest reflexive relation on X that contains R.

Definition (Reflexive Closure (自反闭包))

The reflexive closure $\mathsf{cl}_{\text{ref}}(R)$ of a relation $R \subseteq X \times X$ is the smallest reflexive relation on X that contains R.

$$\mathsf{cl}_{\text{ref}}(R) = R \cup I_X$$

Definition (Symmetric Closure (对称闭包))

The symmetric closure $\mathsf{cl}_{\mathrm{sym}}(R)$ of a relation $R \subseteq X \times X$ is the smallest symmetric relation on X that contains R.

Definition (Symmetric Closure (对称闭包))

The symmetric closure $\mathsf{cl}_{\mathrm{sym}}(R)$ of a relation $R \subseteq X \times X$ is the smallest symmetric relation on X that contains R.

$$\mathsf{cl}_{\mathrm{sym}}(R) = R \cup R^{-1}$$

Definition (Transitive Closure (传递闭包))

The transitive closure $\mathsf{cl}_{\mathrm{trn}}(R)$ of a relation $R \subseteq X \times X$ is the smallest transitive relation on X that contains R.

Definition (Transitive Closure (传递闭包))

The transitive closure $\mathsf{cl}_{\mathrm{trn}}(R)$ of a relation $R \subseteq X \times X$ is the smallest transitive relation on X that contains R.

$$\mathsf{cl}_{\mathrm{trn}}(R) = R^{+}$$

Definition (Transitive Closure (传递闭包))

The transitive closure $\mathsf{cl}_{\mathrm{trn}}(R)$ of a relation $R \subseteq X \times X$ is the smallest transitive relation on X that contains R.

$$\mathsf{cl}_{\mathrm{trn}}(R) = R^+$$

- R^+ contains R
- R^+ is transitive
- R^+ is minimal

Definition (Transitive Closure (传递闭包))

The transitive closure $\mathsf{cl}_{\mathrm{trn}}(R)$ of a relation $R \subseteq X \times X$ is the smallest transitive relation on X that contains R.

$$\mathsf{cl}_{\mathrm{trn}}(R) = R^+$$

- R^+ contains R
- R^+ is transitive
- R^+ is minimal

If T is any transitive relation containing R, then $R^+ \subset T$.

Definition (Transitive Closure (传递闭包))

The transitive closure $\mathsf{cl}_{\mathrm{trn}}(R)$ of a relation $R \subseteq X \times X$ is the smallest transitive relation on X that contains R.

$$\mathsf{cl}_{\mathrm{trn}}(R) = R^+$$

- R^+ contains R
- R^+ is transitive
- R^+ is minimal

If T is any transitive relation containing R, then $R^+ \subset T$.

By induction on i, we can show that $R^i \subseteq T$.

$$f(x)$$

Injection (one-to-one; 1-1)

Surjection

Bijection (one-to-one correspondence)

Definition (Characteristic Function (特征函数) of a Subset)

For a given subset $A \subseteq X$,

$$\chi_A : X \to \{0, 1\}$$

$$\chi_A(x) = 1 \iff x \in A.$$

Definition (Characteristic Function (特征函数) of a Subset)

For a given subset $A \subseteq X$,

$$\chi_A : X \to \{0, 1\}$$

$$\chi_A(x) = 1 \iff x \in A.$$

$$\chi_A : X \to \{0, 1\} \quad vs. \quad \mathcal{P}(X)$$

Definition (Natural Function)

Let $R \subseteq A \times A$ be an equivalence relation. The following function f

$$f : A \to A/R$$

$$f : a \mapsto R(a)$$

is called the natural function on A.

Asymptotic Growth Rates of Functions

https://www.bilibili.com/video/BV175411T7ph?share_source=copy_web

Ordering

Definition (Order Isomorphism (同构))

Given two posets $(S, \leq_S)$ and $(T, \leq_T)$, an order isomorphism from $(S, \leq_S)$ to $(T, \leq_T)$ is a bijection from S to T such that

$$\forall x, y \in S.\ x \leq_S y \leftrightarrow f(x) \leq_T f(y).$$

Definition (Order Isomorphism (同构))

Given two posets $(S, \leq_S)$ and $(T, \leq_T)$, an order isomorphism from $(S, \leq_S)$ to $(T, \leq_T)$ is a bijection from S to T such that

$$\forall x, y \in S.\ x \leq_S y \leftrightarrow f(x) \leq_T f(y).$$

$$(\mathbb{R}, \leq) \xrightarrow[f\colon x \mapsto -x]{f\colon \mathbb{R} \to \mathbb{R}} (\mathbb{R}, \geq)$$

Definition (Order Isomorphism (同构))

Given two posets $(S, \leq_S)$ and $(T, \leq_T)$, an order isomorphism from $(S, \leq_S)$ to $(T, \leq_T)$ is a bijection from S to T such that

$$\forall x, y \in S.\ x \leq_S y \leftrightarrow f(x) \leq_T f(y).$$

$$(\mathbb{R}, \leq) \xrightarrow[f\colon x \mapsto -x]{f\colon \mathbb{R} \to \mathbb{R}} (\mathbb{R}, \geq)$$

Definition (Order Automorphism (自同构))

An order isomorphism from a poset to itself is an order automorphism.

Definition (Rooted Tree (有根树))

A rooted tree is a tree where one vertex has been designated the root.

Definition (Rooted Tree (有根树))

A rooted tree is a tree where one vertex has been designated the root.

Definition (Directed Rooted Tree (有向有根树))

A directed rooted tree is a rooted tree where all edges directed away from or towards the root.

Definition

Parent, Child; Sibling; Ancestor, Descendant

Definition

Parent, Child; Sibling; Ancestor, Descendant

Definition (k-ary Trees (k-叉树))

A k-ary tree is a rooted tree in which each vertex has $\leq k$ children.

2-ary trees are often called binary trees.

Definition

Parent, Child; Sibling; Ancestor, Descendant

Definition (k-ary Trees (k-叉树))

A k-ary tree is a rooted tree in which each vertex has $\leq k$ children.

2-ary trees are often called binary trees.

Definition (Complete k-Tree (完全 k-叉树))

A complete k-tree is a k-ary tree in which each vertex, other than leaves, has $= k$ children.

Depth-First Search (DFS)

Pre-order (前序) Traversal: $F, B, A, D, C, E, G, I, H$

In-order (中序) Traversal: $A, B, C, D, E, F, G, H, I$

Post-order (后序) Traversal: $A, C, E, D, B, H, I, G, F$

Prefix Expression (前缀表达式): $+ * A - BC + DE$

Prefix Expression (前缀表达式): $+ * A - BC + DE$

Infix Expression (中缀表达式): $A * (B - C) + (D + E)$

Prefix Expression (前缀表达式): $+ * A - BC + DE$

Infix Expression (中缀表达式): $A * (B - C) + (D + E)$

Postfix Expression (后缀表达式): $ABC - *DE++$

Breadth-First Search (BFS): $F, B, G, A, D, I, C, E, H$

David A. Huffman (1925 ∼ 1999)

$C[1 \ldots n]$	a	b	c	d	e	f
$F[1 \ldots n]$	45	13	12	16	9	5
Fixed Length Code	000	001	010	011	100	101
Variable Length Code	0	101	100	111	1101	1100

Prefix code (前缀码): No code is a prefix of some other code

The Encoding Problem

To find the optimal binary prefix code for C and F.

Let E be a binary prefix code for C and F. The length $L(E)$ is

$$L(E) = \sum_{c \in C} f_c \cdot l_E(c)$$

$C[1\ldots n]$	a	b	c	d	e	f
$F[1\ldots n]$	45	13	12	16	9	5

Definition (Dual Graph (对偶图))

The dual graph of a plane graph G is a graph G'

- G' has a vertex for each face of G;
- G' has an edge for each pair of faces in G that are separated from each other by an edge, and a self-loop when the same face appears on both sides of an edge.

e_1
e_2
v_1
v_2
e_{n-1}
e_n

The dual graph G' depends on the choice of embedding of the graph G.

Theorem

G *is a bipartite graph* $\iff \chi(G) = 2 \iff G$ *has no odd cycles.*

25
20
15
10
5
1
2
3

Definition (Chromatic Polynomial (色多项式; 非严格定义))

The chromatic polynomial $P(G, k)$ counts the number of colorings of graph G as a function of the number k of colors.

Definition (Chromatic Polynomial (色多项式; 非严格定义))

The chromatic polynomial $P(G,k)$ counts the number of colorings of graph G as a function of the number k of colors.

Triangle K_3	$x(x-1)(x-2)$
Complete graph K_n	$x(x-1)(x-2)\cdots(x-(n-1))$
Edgeless graph $\overline{K}_n$	x^n
Path graph P_n	$x(x-1)^{n-1}$
Any tree on n vertices	$x(x-1)^{n-1}$
Cycle C_n	$(x-1)^n+(-1)^n(x-1)$
Petersen graph	$x(x-1)(x-2)\left(x^7-12x^6+67x^5-230x^4+529x^3-814x^2+775x-352\right)$

Theorem (Recurrence for Chromatic Polynomial)

Given a graph G and an edge $e \in E(G)$, then

$$P(G,k) = P(G-e,k) - P(G/e,k)$$

G/e : 边的收缩

$$P(G,k) = P(G - e, k) - P(G/e, k)$$

$$P(G,k) = P(G-e,k) - P(G/e,k)$$

$$P(G-e,k) = P(G/e,k) + P(G,k)$$

In $G - \{u,v\}$, $\mathrm{Color}(u) = \mathrm{Color}(v)$ or $\mathrm{Color}(u) \neq \mathrm{Color}(v)$.

$$P(G,k) = P(G-e,k) - P(G/e,k)$$

$$P(G,k) = P(G-e,k) - P(G/e,k)$$

$$\begin{aligned} P(C_4,k) &= P(P_4,k) - P(K_3,k) \\ &= k(k-1)^3 - k(k-1)(k-2) \\ &= k(k-1)(k^2-3k+3) \\ &= (k-1)^4 + (-1)^4(k-1) \end{aligned}$$

Cyclic Notation (轮换表示法) & Transposition (对换)

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\begin{aligned}
\sigma &= (1\ 4)(2\ 3\ 6)(5) \\
&= (1\ 4)(2\ 3\ 6) \\
&= (2\ 3\ 6)(1\ 4) \\
&= (2\ 3\ 6)(4\ 1) \\
&= (3\ 6\ 2)(4\ 1) \\
&= (3\ 6)(6\ 2)(4\ 1)
\end{aligned}$$

Cyclic Notation (轮换表示法) & Transposition (对换)

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 3 & 6 & 1 & 5 & 2 \end{pmatrix}$$

$$\begin{aligned} \sigma &= (1\ 4)(2\ 3\ 6)(5) \\ &= (1\ 4)(2\ 3\ 6) \\ &= (2\ 3\ 6)(1\ 4) \\ &= (2\ 3\ 6)(4\ 1) \\ &= (3\ 6\ 2)(4\ 1) \\ &= (3\ 6)(6\ 2)(4\ 1) \end{aligned}$$

$$(i_1\ i_2\ \ldots\ i_r) = (i_1\ i_2)(i_2\ i_3)\ldots(i_{r-2}\ i_{r-1})(i_{r-1}\ i_r)$$

By induction on the length r.

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 7 & 3 & 6 & 2 & 5 & 4 & 1 \end{pmatrix} = (1\ 7)(2\ 3)(3\ 6)(6\ 4)$$

$$= (1\ 7)(3\ 6)(2\ 5)(6\ 4)(4\ 5)(2\ 5)$$

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 7 & 3 & 6 & 2 & 5 & 4 & 1 \end{pmatrix} = (1\ 7)(2\ 3)(3\ 6)(6\ 4)$$

$$= (1\ 7)(3\ 6)(2\ 5)(6\ 4)(4\ 5)(2\ 5)$$

Theorem (Parity (奇偶性) of Permutations)

将一个置换表示成若干对换的乘积, 所用对换个数的奇偶性是唯一的。

Definition (Even/Odd Permutations (偶置换/奇置换))

可表示为偶数个对换的乘积的置换称为偶置换; 否则, 称为奇置换。

Definition (Even/Odd Permutations (偶置换/奇置换))

可表示为偶数个对换的乘积的置换称为偶置换; 否则, 称为奇置换。

Definition (Alternating Group (交错群; A_n))

由 S_n 的全体偶置换构成的子群称为 n 次交错群。

Definition (Even/Odd Permutations (偶置换/奇置换))

可表示为偶数个对换的乘积的置换称为偶置换; 否则, 称为奇置换。

Definition (Alternating Group (交错群; A_n))

由 S_n 的全体偶置换构成的子群称为 n 次交错群。

$$A_3 = \{(1), (1\ 2\ 3), (1\ 3\ 2)\}$$

$$\mathrm{sgn} : S_n \to \{1, -1\}$$

$$\mathrm{sgn} : S_n \to \{1, -1\}$$

$$\mathrm{sgn}(\sigma_1\sigma_2) = \mathrm{sgn}(\sigma_1)\mathrm{sgn}(\sigma_2)$$

$$\mathrm{sgn} : S_n \to \{1, -1\}$$

$$\mathrm{sgn}(\sigma_1\sigma_2) = \mathrm{sgn}(\sigma_1)\mathrm{sgn}(\sigma_2)$$

$$S_n/A_n \cong \{1, -1\}$$

$$A_3 = \{(1), (1\ 2\ 3), (1\ 3\ 2)\}$$

$$(1\ 2)A_3 = \{(1\ 2), (2\ 3), (1\ 3)\}$$

Thank You!

Office 302

Mailbox: H016

hfwei@nju.edu.cn