


(八) 集合: 无穷 (Infinity)


魏恒峰

hfwei@nju.edu.cn


2021 年 04 月 29 日








Georg Cantor (1845 – 1918)



Georg Cantor (1845 – 1918)



Leopold Kronecker
(1823 – 1891)




Georg Cantor (1845 – 1918)



Leopold Kronecker
(1823 – 1891)



Henri Poincaré
(1854 – 1912)



Georg Cantor (1845 – 1918)




Leopold Kronecker
(1823 – 1891)



Henri Poincaré
(1854 – 1912)



Ludwig Wittgenstein
(1889 – 1951)



Georg Cantor (1845 – 1918)



David Hilbert (1862 – 1943)



Leopold Kronecker
(1823 – 1891)



Henri Poincaré
(1854 – 1912)



Ludwig Wittgenstein
(1889 – 1951)

*From his paradise that Cantor with us unfolded, we hold our
breath in awe; knowing, we shall not be expelled.*

— *David Hilbert*

“没有人能把我们从 Cantor 创造的乐园中驱逐出去”

*From his paradise that Cantor with us unfolded, we hold our
breath in awe; knowing, we shall not be expelled.*

— *David Hilbert*

“没有人能把我们从 Cantor 创造的乐园中驱逐出去”








“das wesen der mathematik liegt in ihrer freiheit”



“das wesen der mathematik liegt in ihrer freiheit”

“The essence of mathematics lies in its freedom”


Before Cantor









公理：“整体大于部分”



Galileo Galilei (1564 – 1642)



“关于两门新科学的对话” (1638)



Galileo Galilei (1564 – 1642)

“关于两门新科学的对话” (1638)

“用我们有限的心智来讨论无限 …”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”



吓得我吃了一鲸

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”



吓得我吃了一鲸

说到底，“等于”、“大于”和“小于”诸性质不能用于无限，而只能用于有限的数量。
— Galileo Galilei

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”



吓得我吃了一鲸

说到底，“等于”、“大于”和“小于”诸性质不能用于无限，而只能用于有限的数量。
— Galileo Galilei

无穷数是不可能的。
— Gottfried Wilhelm Leibniz

这些证明一开始就期望那些数要具有有穷数的一切性质，或者甚至于把有穷数的性质强加于无穷。

相反，这些无穷数，如果它们能够以任何形式被理解的话，倒是由于它们与有穷数的对应，它们必须具有完全新的数量特征。

这些性质完全依赖于事物的本性，… 而并非来自我们的主观任意性或我们的偏见。

— Georg Cantor (1885)

Definition (Dedekind-infinite & Dedekind-finite (Dedekind, 1888))


A set A is *Dedekind-infinite* if there is a bijective function from A onto some proper subset B of A .

A set is *Dedekind-finite* if it is not Dedekind-infinite.

Definition (Dedekind-infinite & Dedekind-finite (Dedekind, 1888))

A set A is *Dedekind-infinite* if there is a bijective function from A onto some proper subset B of A .

A set is *Dedekind-finite* if it is not Dedekind-infinite.



This is a **theorem** in our theory of infinity.



We have not defined “finite” and “infinite”!

Comparing Sets

Comparing Sets



Comparing Sets



Function



Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

$\overline{\overline{A}}$ (two *abstractions*)

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

$\overline{\overline{A}}$ (two *abstractions*)

Abstract from elements: $\{1, 2, 3\}$ vs. $\{a, b, c\}$

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

$\overline{\overline{A}}$ (two *abstractions*)

Abstract from elements: $\{1, 2, 3\}$ vs. $\{a, b, c\}$

Abstract from order: $\{1, 2, 3, \dots\}$ vs. $\{1, 3, 5, \dots, 2, 4, 6, \dots\}$

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Q : Is “ \approx ” an equivalence relation?

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Q : Is “ \approx ” an equivalence relation?

Theorem ()

For any sets A, B, C :

- (a) $A \approx B$
- (b) $A \approx B \implies B \approx A$
- (c) $A \approx B \wedge B \approx C \implies A \approx C$

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Q : Is “ \approx ” an equivalence relation?

Theorem (The “Equivalence Concept” of Equipotent)

For any sets A, B, C :

- (a) $A \approx B$
- (b) $A \approx B \implies B \approx A$
- (c) $A \approx B \wedge B \approx C \implies A \approx C$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n.$$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n.$$

$$|X| = |\{0, 1, \dots, n-1\}|$$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n.$$

$$|X| = |\{0, 1, \dots, n-1\}|$$

Theorem

*Let A be a finite set. There is a **unique** $n \in \mathbb{N}$ such that $A \approx \{0, 1, \dots, n-1\}$.*

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem

\mathbb{N} is infinite.

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem

\mathbb{N} is infinite.

By Contradiction.

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

$$g \triangleq f|_{\{0,1,\dots,n\}} : \{0, 1, \dots, n\} \rightarrow \{0, 1, \dots, n-1\}$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

$$g \triangleq f|_{\{0,1,\dots,n\}} : \{0, 1, \dots, n\} \rightarrow \{0, 1, \dots, n-1\}$$

By the Pigeonhole Principle : g is not 1-1

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem

\mathbb{N} is infinite. (So are \mathbb{Z} , \mathbb{Q} , \mathbb{R} .)

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

$$g \triangleq f|_{\{0,1,\dots,n\}} : \{0, 1, \dots, n\} \rightarrow \{0, 1, \dots, n-1\}$$

By the Pigeonhole Principle : g is not 1-1 $\implies f$ is not 1-1

Definition (Infinite)

For any set X ,

Countably Infinite

$$|X| = |\mathbb{N}| \triangleq \aleph_0$$

Countable

(finite \vee countably infinite)

Uncountable


(\neg countable)

(infinite) \wedge (\neg (countably infinite))

\aleph_0


Theorem (\mathbb{Z} is Countable.)

$$|\mathbb{Z}| = |\mathbb{N}|$$



Theorem (\mathbb{Z} is Countable.)

$$|\mathbb{Z}| = |\mathbb{N}|$$



0 1 -1 2 -2 ...

Theorem (\mathbb{Q} is Countable. (Cantor 1873-11; Published in 1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$

Theorem (\mathbb{Q} is Countable. (Cantor 1873-11; Published in 1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$

$$|\mathbb{Q}| = |\mathbb{N}|$$


$$q \in \mathbb{Q}^+ : a/b \ (a, b \in \mathbb{N}^+)$$

Theorem (\mathbb{Q} is Countable. (Cantor 1873-11; Published in 1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$


$$|\mathbb{Q}| = |\mathbb{N}|$$

$$q \in \mathbb{Q}^+ : a/b \ (a, b \in \mathbb{N}^+)$$




Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)

$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$



Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)


$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$



$$\pi : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$$

Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)

$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$




$$\pi : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$$

$$\pi(k_1, k_2) = \frac{1}{2}(k_1 + k_2)(k_1 + k_2 + 1) + k_2$$

Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)

$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$



Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

*The Cartesian product of **finitely many** countable sets is countable.*

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

*The Cartesian product of **finitely many** countable sets is countable.*

$$\mathbb{N}^n \quad \text{vs.} \quad \mathbb{N}^{\mathbb{N}}$$

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

*The Cartesian product of **finitely many** countable sets is countable.*

$$\mathbb{N}^n \quad \text{vs.} \quad \mathbb{N}^{\mathbb{N}}$$

$$\pi^{(n)} : \mathbb{N}^n \rightarrow \mathbb{N}$$

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

The Cartesian product of *finitely many* countable sets is countable.

$$\mathbb{N}^n \quad \text{vs.} \quad \mathbb{N}^{\mathbb{N}}$$

$$\pi^{(n)} : \mathbb{N}^n \rightarrow \mathbb{N}$$

$$\pi^{(n)}(k_1, \dots, k_{n-1}, k_n) = \pi(\pi^{(n-1)}(k_1, \dots, k_{n-1}), k_n)$$

Theorem

*Any **finite** union of countable sets is countable.*

Theorem

Any *finite* union of countable sets is countable.

$$A = \{a_n \mid n \in \mathbb{N}\} \quad B = \{b_n \mid n \in \mathbb{N}\} \quad C = \{c_n \mid n \in \mathbb{N}\}$$

Theorem

Any *finite* union of countable sets is countable.

$$A = \{a_n \mid n \in \mathbb{N}\} \quad B = \{b_n \mid n \in \mathbb{N}\} \quad C = \{c_n \mid n \in \mathbb{N}\}$$


$$a_0 \quad b_0 \quad c_0 \quad a_1 \quad b_1 \quad c_1 \cdots$$

Theorem

*The union of **countably many** countable sets is countable.*

Theorem


The union of *countably many* countable sets is countable.



Counting by Diagonals.

Theorem

The union of *countably many* countable sets is countable.



Counting by Diagonals.

We need Axiom of (Countable) Choice!

Beyond

\aleph_0

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

**VERY
IMPORTANT**

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$



Different “Sizes” of Infinity

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$



Different “Sizes” of Infinity

Cantor’s Diagonal Argument (1890)

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} \mathbb{N}$$

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} \mathbb{N}$$

3.14159...
1.41421...
1.73205...
2.23606...
2.71828...
0.14285...



3.43625...



2.32514...

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} \mathbb{N}$$

3.14159...
1.41421...
1.73205...
2.23606...
2.71828...
0.14285...



3.43625...



2.32514...

By Diagonal Argument.

$$\mathfrak{c} \triangleq |\mathbb{R}|$$

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$f(x) = \tan \frac{(2x - 1)\pi}{2}$$

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$f(x) = \tan \frac{(2x - 1)\pi}{2}$$

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots)$$

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$f(x) = \tan \frac{(2x - 1)\pi}{2}$$

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$



$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots)$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$



Was Cantor Surprised?

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^n|$$

“Je le vois, mais je ne le crois pas !”

“I see it, but I don't believe it !”

— Cantor's letter to Dedekind (1877).

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^n|$$

“Je le vois, mais je ne le crois pas !”

“I see it, but I don't believe it !”

— Cantor's letter to Dedekind (1877).

Q : Then, what is “dimension”?

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^n|$$

“Je le vois, mais je ne le crois pas !”

“I see it, but I don't believe it !”

— Cantor's letter to Dedekind (1877).

Q : Then, what is “dimension”?

Theorem (Brouwer (Topological Invariance of Dimension))

*There is no **continuous** bijections between \mathbb{R}^m and \mathbb{R}^n for $m \neq n$.*

Beyond



Theorem (Cantor's Theorem (1891))

$$|A| \neq |\mathcal{P}(A)|$$

Theorem (Cantor's Theorem (1891))

$$|A| \neq |\mathcal{P}(A)|$$

Theorem (Cantor Theorem)

If $f : A \rightarrow \mathcal{P}(A)$, then f is not onto.

Theorem (Cantor Theorem)

$$|A| < |\mathcal{P}(A)|$$

Theorem (Cantor Theorem)

$$|A| < |\mathcal{P}(A)|$$


$$A \quad \mathcal{P}(A) \quad \mathcal{P}(\mathcal{P}(A)) \quad \dots$$

Theorem (Cantor Theorem)

$$|A| < |\mathcal{P}(A)|$$

$$A \quad \mathcal{P}(A) \quad \mathcal{P}(\mathcal{P}(A)) \quad \dots$$

There is no largest infinity.



Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

Q : What about onto function $f : A \rightarrow B$?

Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

Q : What about onto function $f : A \rightarrow B$?

$$|B| \leq |A|$$

Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

Q : What about onto function $f : A \rightarrow B$?

$|B| \leq |A|$ (Axiom of Choice)

Definition ($|A| < |B|$)

$$|A| < |B| \iff |A| \leq |B| \wedge |A| \neq |B|$$

Definition ($|A| < |B|$)

$$|A| < |B| \iff |A| \leq |B| \wedge |A| \neq |B|$$

$$|\mathbb{N}| < |\mathbb{R}|$$

$$|X| < |2^X|$$

$$|\mathbb{N}| < |2^{\mathbb{N}}|$$

Definition (Countable Revisited)

X is countable:

$$(\exists n \in \mathbb{N} : |X| = n) \vee |X| = |\mathbb{N}|$$

Definition (Countable Revisited)

X is countable:

$$(\exists n \in \mathbb{N} : |X| = n) \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable)

X is countable iff

$$|X| \leq |\mathbb{N}|.$$

Definition (Countable Revisited)

X is countable:

$$(\exists n \in \mathbb{N} : |X| = n) \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable)

X is countable iff

$$|X| \leq |\mathbb{N}|.$$

X is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

Definition (Countable Revisited)

X is countable:

$$(\exists n \in \mathbb{N} : |X| = n) \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable)

X is countable iff

$$|X| \leq |\mathbb{N}|.$$

X is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

Subsets of Countable Set

Every subset B of a countable set A is countable.

Definition (Countable Revisited)

X is countable:

$$(\exists n \in \mathbb{N} : |X| = n) \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable)

X is countable iff

$$|X| \leq |\mathbb{N}|.$$

X is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

Subsets of Countable Set

Every subset B of a countable set A is countable.

$$f : A \xrightarrow{1-1} \mathbb{N}$$

Definition (Countable Revisited)

X is countable:

$$(\exists n \in \mathbb{N} : |X| = n) \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable)

X is countable iff

$$|X| \leq |\mathbb{N}|.$$

X is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

Subsets of Countable Set

Every subset B of a countable set A is countable.

$$f : A \xrightarrow{1-1} \mathbb{N} \quad g = f|_B$$

Slope

(a) The set of all lines with rational slopes

Slope

(a) The set of all lines with rational slopes

$$(\mathbb{Q}, \mathbb{R})$$

Slope

(a) The set of all lines with rational slopes

$$(\mathbb{Q}, \mathbb{R})$$

$$|\mathbb{R}| \leq |\mathbb{Q} \times \mathbb{R}| \leq |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}|$$

Q : Is “ \leq ” a partial order?

Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder–Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$


$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$




Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$




Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder–Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$



Schröder–Bernstein
theorem @ wiki

Q : Is “ \leq ” a total order?

Q : Is “ \leq ” a total order?

Theorem (PCC)

Principle of Cardinal Comparability (PCC) \iff Axiom of Choice

Theorem

$$|\mathbb{R}| = |\mathcal{P}(\mathbb{N})|$$

Theorem

$$|\mathbb{R}| = |\mathcal{P}(\mathbb{N})|$$

$$|\mathbb{R}| \leq |\mathcal{P}(\mathbb{N})| \quad |\mathcal{P}(\mathbb{N})| \leq |\mathbb{R}|$$

Theorem

$$|\mathbb{R}| = |\mathcal{P}(\mathbb{N})|$$

$$|\mathbb{R}| \leq |\mathcal{P}(\mathbb{N})| \quad |\mathcal{P}(\mathbb{N})| \leq |\mathbb{R}|$$

$$\mathfrak{c} \triangleq |\mathbb{R}| = |\mathcal{P}(\mathbb{N})| = |2^{\mathbb{N}}| \triangleq 2^{\aleph_0}$$

Theorem

$$|\mathbb{R}| = |\mathcal{P}(\mathbb{N})|$$

$$|\mathbb{R}| \leq |\mathcal{P}(\mathbb{N})| \quad |\mathcal{P}(\mathbb{N})| \leq |\mathbb{R}|$$

$$\mathfrak{c} \triangleq |\mathbb{R}| = |\mathcal{P}(\mathbb{N})| = |2^{\mathbb{N}}| \triangleq 2^{\aleph_0}$$

$$\mathfrak{c} = 2^{\aleph_0}$$

Continuum Hypothesis (CH)

$$\nexists A : \aleph_0 < |A| < \mathfrak{c}$$



👉 Dangerous Knowledge (22:20; BBC 2007)




👉 Dangerous Knowledge (22:20; BBC 2007)

Independence from ZFC:

Kurt Gödel (1940) CH cannot be disproved from ZF.

Paul Cohen (1964) CH cannot be proven from the ZFC axioms.

Thank
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn