

# Homework

魏恒峰

hfwei@nju.edu.cn

2018 年 02 月 28 日



## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

## Definition (Well-ordered Set (SM Definition 14.1))

An ordered set  $S$  is said to be *well-ordered* if every non-empty subset of  $S$  has a first element.

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

## Definition (Well-ordered Set (SM Definition 14.1))

An ordered set  $S$  is said to be *well-ordered* if every non-empty subset of  $S$  has a first element.

## Definition (Isomorphic)

Two ordered sets  $A$  and  $B$  are said to be *isomorphic*, written  $A \simeq B$ , if  $\exists f : A \xrightarrow[\text{onto}]{1-1} B$  which preserves the order relations

$$\forall a, a' \in A : a \prec a' \iff f(a) \prec f(a')$$

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Remark: What if “similarity mapping”?

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Remark: What if “similarity mapping”?

### Definition (Similarity Mapping)

A function  $f : A \xrightarrow{1-1} B$  is called a *similarity mapping* from  $A$  to  $B$  if  $f$  preserves the order relations

$$\forall a, a' \in A : a \prec a' \iff f(a) \prec f(a')$$

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Remark: What if “similarity mapping”?

### Definition (Similarity Mapping)

A function  $f : A \xrightarrow{1-1} B$  is called a *similarity mapping* from  $A$  to  $B$  if  $f$  preserves the order relations

$$\forall a, a' \in A : a \prec a' \iff f(a) \prec f(a')$$

Counterexample for “similarity mapping”:

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Remark: What if “similarity mapping”?

### Definition (Similarity Mapping)

A function  $f : A \xrightarrow{1-1} B$  is called a *similarity mapping* from  $A$  to  $B$  if  $f$  preserves the order relations

$$\forall a, a' \in A : a \prec a' \iff f(a) \prec f(a')$$

Counterexample for “similarity mapping”:

$$A = B = \mathbb{N} \quad f : a \mapsto a \quad f' : a \mapsto a + 1$$



## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .



## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .



## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

## Lemma

$X$  is a well-ordered set.  $f : X \rightarrow X$  is a similarity mapping.

Then  $\forall x \in X : f(x) \geq x$ .

### SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

### Lemma

$X$  is a well-ordered set.  $f : X \rightarrow X$  is a similarity mapping.

Then  $\forall x \in X : f(x) \geq x$ .

### Lemma

$X$  is a well-ordered set.

Then  $X$  has only one *automorphism*, i.e.,  $id_X$ .

## Lemma

*$X$  is a well-ordered set.  $f : X \rightarrow X$  is a similarity mapping.  
Then  $\forall x \in X : f(x) \geq x$ .*

Proof.

## Lemma

*$X$  is a well-ordered set.  $f : X \rightarrow X$  is a similarity mapping.  
Then  $\forall x \in X : f(x) \geq x$ .*

## Proof.

By contradiction.

## Lemma

*$X$  is a well-ordered set.  $f : X \rightarrow X$  is a similarity mapping.  
Then  $\forall x \in X : f(x) \geq x$ .*

## Proof.

By contradiction.

$$Y = \{x \in X \mid f(x) < x\} \neq \emptyset$$



## Lemma

*$X$  is a well-ordered set.  $f : X \rightarrow X$  is a similarity mapping.  
Then  $\forall x \in X : f(x) \geq x$ .*

## Proof.

By contradiction.

$$Y = \{x \in X \mid f(x) < x\} \neq \emptyset$$

Well-ordered:  $x = \min Y \implies f(x) < x$

## Lemma

*$X$  is a well-ordered set.  $f : X \rightarrow X$  is a similarity mapping.  
Then  $\forall x \in X : f(x) \geq x$ .*

## Proof.

By contradiction.

$$Y = \{x \in X \mid f(x) < x\} \neq \emptyset$$

**Well-ordered:**  $x = \min Y \implies f(x) < x$

**Similarity:**  $f(f(x)) < f(x) \implies f(x) \in Y$



## Lemma

$X$  is a well-ordered set.

Then  $X$  has only one *automorphism*, i.e.,  $\text{id}_X$ .

Proof.

## Lemma

$X$  is a well-ordered set.

Then  $X$  has only one *automorphism*, i.e.,  $\text{id}_X$ .

## Proof.

Suppose  $f : X \rightarrow X$  is an automorphism.

## Lemma

$X$  is a well-ordered set.

Then  $X$  has only one *automorphism*, i.e.,  $id_X$ .

## Proof.

Suppose  $f : X \rightarrow X$  is an automorphism.

$f$  is a similarity from  $X$  to  $X \implies \forall x \in X : f(x) \geq x$

## Lemma

$X$  is a well-ordered set.

Then  $X$  has only one *automorphism*, i.e.,  $id_X$ .

## Proof.

Suppose  $f : X \rightarrow X$  is an automorphism.

$f$  is a similarity from  $X$  to  $X \implies \forall x \in X : f(x) \geq x$

$f^{-1}$  is a similarity from  $X$  to  $X \implies \forall x \in X : f^{-1}(x) \geq x$

## Lemma

$X$  is a well-ordered set.

Then  $X$  has only one *automorphism*, i.e.,  $id_X$ .

## Proof.

Suppose  $f : X \rightarrow X$  is an automorphism.

$f$  is a similarity from  $X$  to  $X \implies \forall x \in X : f(x) \geq x$

$f^{-1}$  is a similarity from  $X$  to  $X \implies \forall x \in X : f^{-1}(x) \geq x$

$f$  is a similarity from  $X$  to  $X \implies x = f(f^{-1}(x)) \geq f(x)$

## Lemma

$X$  is a well-ordered set.

Then  $X$  has only one *automorphism*, i.e.,  $id_X$ .

## Proof.

Suppose  $f : X \rightarrow X$  is an automorphism.

$f$  is a similarity from  $X$  to  $X \implies \forall x \in X : f(x) \geq x$

$f^{-1}$  is a similarity from  $X$  to  $X \implies \forall x \in X : f^{-1}(x) \geq x$

$f$  is a similarity from  $X$  to  $X \implies x = f(f^{-1}(x)) \geq f(x)$

$$id_X : f(x) = x$$





## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Proof.

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Proof.

By contradiction.

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Proof.

By contradiction.

$\exists$  isomorphic mappings  $f, g : A \rightarrow B$

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Proof.

By contradiction.

$\exists$  isomorphic mappings  $f, g : A \rightarrow B$

$h = g^{-1} \circ f$  is an automorphism on  $A \implies h = id_A$

## SM Problem 14.62

Suppose  $A$  and  $B$  are *well-ordered isomorphic* sets.

Show that there is only one *isomorphic mapping*  $f : A \rightarrow B$ .

Proof.

By contradiction.

$\exists$  isomorphic mappings  $f, g : A \rightarrow B$

$h = g^{-1} \circ f$  is an automorphism on  $A \implies h = id_A$

$$f = g \circ h = g \circ id_A = g$$



## Boolean Algebra

(i) 等势且有穷的布尔代数均同构。

Proof.

## Boolean Algebra

(i) 等势且有穷的布尔代数均同构。

Proof.

$$|B_1| = |B_2| = m$$

## Boolean Algebra

(i) 等势且有穷的布尔代数均同构。

Proof.

$$|B_1| = |B_2| = m \quad f : B_1 \xrightarrow[\text{onto}]{1-1} B_2$$



## Boolean Algebra

(i) 等势且有穷的布尔代数均同构。

Proof.

$$|B_1| = |B_2| = m \quad f : B_1 \xrightarrow[\text{onto}]{1-1} B_2$$

### Theorem (Representation Theorem)

*B is a finite Boolean algebra.  $A_B$  is the set of atoms of B.*

$$f : B \xrightarrow[\text{onto}]{1-1} \mathcal{P}(A_B)$$

*$x = a_1 + a_2 + \cdots + a_r \mapsto \{a_1, a_2, \cdots, a_r\}$  is an isomorphism.*

## Boolean Algebra

(i) 等势且有穷的布尔代数均同构。

Proof.

$$|B_1| = |B_2| = m \quad f : B_1 \xleftrightarrow[\text{onto}]{1-1} B_2$$

### Theorem (Representation Theorem)

$B$  is a finite Boolean algebra.  $A_B$  is the set of atoms of  $B$ .

$$f : B \xleftrightarrow[\text{onto}]{1-1} \mathcal{P}(A_B)$$

$x = a_1 + a_2 + \cdots + a_r \mapsto \{a_1, a_2, \cdots, a_r\}$  is an isomorphism.

$$|A_{B_1}| = |A_{B_2}| = n = \log_2 m \quad A_{B_1} = \{b_1, \cdots, b_n\} \quad A_{B_2} = \{b_2, \cdots, b_n\}$$



## Boolean Algebra

(ii) 举例说明等势但无穷的布尔代数不一定同构。

## Boolean Algebra

(ii) 举例说明等势但无穷的布尔代数不一定同构。

Not Exactly an Answer.

## Boolean Algebra

(ii) 举例说明等势但无穷的布尔代数不一定同构。

Not Exactly an Answer.

并非任何 Boolean Algebra 皆同构于某幂集代数。

有穷-余有穷 (finite co-finite) 代数  $F(\mathbb{N})$

$$F(\mathbb{N}) = \{X \in \mathcal{P}(\mathbb{N}) \mid X \text{ is finite} \vee \mathbb{N} \setminus X \text{ is finite}\}$$

## Boolean Algebra

(ii) 举例说明等势但无穷的布尔代数不一定同构。

Not Exactly an Answer.

并非任何 Boolean Algebra 皆同构于某幂集代数。

有穷-余有穷 (finite co-finite) 代数  $F(\mathbb{N})$

$$F(\mathbb{N}) = \{X \in \mathcal{P}(\mathbb{N}) \mid X \text{ is finite} \vee \mathbb{N} \setminus X \text{ is finite}\}$$

$$|F(\mathbb{N})| = \aleph_0$$

## Boolean Algebra

(ii) 举例说明等势但无穷的布尔代数不一定同构。

Not Exactly an Answer.

并非任何 Boolean Algebra 皆同构于某幂集代数。

有穷-余有穷 (finite co-finite) 代数  $F(\mathbb{N})$

$$F(\mathbb{N}) = \{X \in \mathcal{P}(\mathbb{N}) \mid X \text{ is finite} \vee \mathbb{N} \setminus X \text{ is finite}\}$$

$$|F(\mathbb{N})| = \aleph_0$$

$$f : F(\mathbb{N}) \xrightarrow{1-1} \mathbb{N} \quad \{a_1, a_2, \dots, a_n\} = p_{a_1} p_{a_2} \cdots p_{a_n}$$



## Theorem (Representation Theorem)

- (i) 任何有穷 *Boolean Algebra* 同构于某幂集代数。
- (ii) 有穷 *Boolean Algebra* 之势呈形  $2^n$ 。
- (iii) 两个等势的有穷 *Boolean Algebra* 是同构的。
- (iv) 并非任何 *Boolean Algebra* 皆同构于某幂集代数。



Thank  
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn