


# Finite and Infinite

魏恒峰


hfwei@nju.edu.cn

2018 年 02 月 26 日






Georg Cantor (1845 – 1918)



Georg Cantor (1845 – 1918)



Leopold Kronecker  
(1823 – 1891)




Georg Cantor (1845 – 1918)



Leopold Kronecker  
(1823 – 1891)



Henri Poincaré  
(1854 – 1912)



## Georg Cantor (1845 – 1918)



Leopold Kronecker  
(1823 – 1891)




Henri Poincaré  
(1854 – 1912)



## Ludwig Wittgenstein

◀ ◻ ▶ ◀ ◻ ▶ (1889–1951) ≡ ↺ 🔍 ↻



Georg Cantor (1845 – 1918)



David Hilbert (1862 – 1943)



Leopold Kronecker  
(1823 – 1891)



Henri Poincaré  
(1854 – 1912)



Ludwig Wittgenstein

◀ ◻ ▶ ◀ ◻ ▶ (1889 – 1951) ≡ 🔍 ↺ ↻

*From his paradise that Cantor with us unfolded, we hold our  
breath in awe; knowing, we shall not be expelled.*

— *David Hilbert*

“没有人能把我们从 Cantor 创造的乐园中驱逐出去”

*From his paradise that Cantor with us unfolded, we hold our  
breath in awe; knowing, we shall not be expelled.*

— *David Hilbert*

“没有人能把我们从 Cantor 创造的乐园中驱逐出去”










*“das wesen der mathematik liegt in ihrer freiheit”*




*“das wesen der mathematik liegt in ihrer freiheit”*

*“The essence of mathematics lies in its freedom”*



Galileo Galilei (1564 – 1642)



《关于两门新科学的对话》(1638)

“用我们有限的心智来讨论无限 …”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

“用我们有限的心智来讨论无限 …”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2|, S_1 \subsetneq S_2$$

“用我们有限的心智来讨论无限 …”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2|, S_1 \subsetneq S_2$$

“部分等于全体”

“用我们有限的心智来讨论无限 ……”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2|, S_1 \subsetneq S_2$$

“部分等于全体”



吓得我吃了一鲸



“用我们有限的心智来讨论无限 …”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2|, S_1 \subsetneq S_2$$

“部分等于全体”



吓得我吃了一鲸

说到底，“等于”、“大于”和“小于”诸性质不能用于无限，而  
只能用于有限的数量。  
— Galileo Galilei

“用我们有限的心智来讨论无限 …”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2|, S_1 \subsetneq S_2$$

“部分等于全体”



吓得我吃了一鲸

说到底，“等于”、“大于”和“小于”诸性质不能用于无限，而  
只能用于有限的数量。  
— Galileo Galilei

无穷数是不可能的。  
— Gottfried Wilhelm Leibniz

这些证明一开始就期望那些数要具有有穷数的一切性质，或者甚至于把有穷数的性质强加于无穷。

相反，这些无穷数，如果它们能够以任何形式被理解的话，倒是由于它们与有穷数的对应，它们必须具有完全新的数量特征。

这些性质完全依赖于事物的本性，... 而并非来自我们的主观任意性或我们的偏见。

— Georg Cantor (1885)

这些证明一开始就期望那些数要具有有穷数的一切性质，或者甚至于把有穷数的性质强加于无穷。

相反，这些无穷数，如果它们能够以任何形式被理解的话，倒是由于它们与有穷数的对应，它们必须具有完全新的数量特征。

这些性质完全依赖于事物的本性， $\cdots$  而并非来自我们的主观任意性或我们的偏见。

— Georg Cantor (1885)

### Definition (Dedekind-infinite & Dedekind-finite (Dedekind, 1888))

A set  $A$  is **Dedekind-infinite** if there is a bijective function from  $A$  onto some proper subset  $B$  of  $A$ .

A set is **Dedekind-finite** if it is not Dedekind-infinite.



## Comparing Sets



## Comparing Sets





## Comparing Sets



Function





Definition ( $|A| = |B|$  ( $A \approx B$ ) (1878))

Two sets of  $A$  and  $B$  are *equipotent* if there exists a *bijection* from  $A$  to  $B$ .

Definition ( $|A| = |B|$  ( $A \approx B$ ) (1878))

Two sets of  $A$  and  $B$  are *equipotent* if there exists a *bijection* from  $A$  to  $B$ .

“ $\approx$ ” is an equivalence relation.

Definition ( $|A| = |B|$  ( $A \approx B$ ) (1878))

Two sets of  $A$  and  $B$  are *equipotent* if there exists a *bijection* from  $A$  to  $B$ .

“ $\approx$ ” is an equivalence relation.

$\overline{A}$  (two *abstractions*)

Definition ( $|A| = |B|$  ( $A \approx B$ ) (1878))

Two sets of  $A$  and  $B$  are *equipotent* if there exists a *bijection* from  $A$  to  $B$ .

“ $\approx$ ” is an equivalence relation.

$\overline{A}$  (two *abstractions*)

$\{1, 2, 3\}$  vs.  $\{a, b, c\}$

Definition ( $|A| = |B|$  ( $A \approx B$ ) (1878))

Two sets of  $A$  and  $B$  are *equipotent* if there exists a *bijection* from  $A$  to  $B$ .

“ $\approx$ ” is an equivalence relation.

$\overline{A}$  (two *abstractions*)

$\{1, 2, 3\}$  vs.  $\{a, b, c\}$

$\{1, 2, 3, \dots\}$  vs.  $\{1, 3, 5, \dots, 2, 4, 6, \dots\}$

## Definition (Finite and Infinite)

For any set  $X$ ,

Finite

$$\exists n \in \mathbb{N} : |X| = n \quad (0 \in \mathbb{N})$$

Infinite ( $\neg$  finite)

$$\forall n \in \mathbb{N} : |X| \neq n$$

## Definition (Finite and Infinite)

For any set  $X$ ,

Countably Infinite

$$|X| = |\mathbb{N}| \triangleq \aleph_0$$

Countable

(finite  $\vee$  countably infinite)

Uncountably Infinite

$$(\neg \text{finite}) \wedge (\neg (\text{countably infinite}))$$

$$(\neg \text{countable})$$

## Theorem ( $\aleph_0$ (1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$




# Theorem ( $\aleph_0$ (1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$

$$|\mathbb{Q}| = |\mathbb{N}| \text{ (UD 22.9)}$$

$$q \in \mathbb{Q}^+ : a/b \ (a, b \in \mathbb{N}^+)$$



$$|\mathbb{N}| = |\mathbb{Z}|$$

$$|\mathbb{N}| = |\mathbb{Z}|$$

$$|\mathbb{N}| = |\mathbb{N} \times \mathbb{N}|$$

$$|\mathbb{N}| = |\mathbb{Z}|$$

$$|\mathbb{N}| = |\mathbb{N} \times \mathbb{N}|$$

$$f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N} \quad f(m, n) = n + \frac{(m+n)(m+n+1)}{2}$$

Theorem ( $\mathbb{R}$  is uncountably infinite (1874) .)

$$|\mathbb{R}| \neq |\mathbb{N}| \quad (|\mathbb{R}| > |\mathbb{N}|)$$

Theorem ( $\mathbb{R}$  is uncountably infinite (1874) .)

$$|\mathbb{R}| \neq |\mathbb{N}| \quad (|\mathbb{R}| > |\mathbb{N}|)$$

Proof.

See UD Theorem 22.12.  $f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} (0, 1)$ .



Theorem ( $\mathbb{R}$  is uncountably infinite (1874) .)

$$|\mathbb{R}| \neq |\mathbb{N}| \quad (|\mathbb{R}| > |\mathbb{N}|)$$

Proof.

See UD Theorem 22.12.  $f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} (0, 1)$ .



Different “Sizes” of Infinity

Theorem ( $\mathbb{R}$  is uncountably infinite (1874) .)

$$|\mathbb{R}| \neq |\mathbb{N}| \quad (|\mathbb{R}| > |\mathbb{N}|)$$

Proof.

See UD Theorem 22.12.  $f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} (0, 1)$ .



Different “Sizes” of Infinity

Cantor’s Diagonal Argument (1890)



Theorem ( $\mathbb{R}$  is uncountably infinite (1874) .)

$$|\mathbb{R}| \neq |\mathbb{N}| \quad (|\mathbb{R}| > |\mathbb{N}|)$$

Proof.

See UD Theorem 22.12.  $f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} (0, 1)$ .



Different “Sizes” of Infinity

Cantor’s Diagonal Argument (1890)

Theorem (Cantor’s Theorem (1891))

$$|X| \neq |2^X| \quad (|X| < |2^X|)$$

## Infinite Sequences of 0's and 1's (UD 22.3)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

## Infinite Sequences of 0's and 1's (UD 22.3)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

Proof.

By Cantor's diagonal argument  $\implies$  uncountable. □

## Infinite Sequences of 0's and 1's (UD 22.3)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

Proof.

By Cantor's diagonal argument  $\implies$  uncountable. □

Nonproof.

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

$$f(x_0x_1\cdots) = \sum_{i=0}^{\infty} x_i 2^i$$
□

## Theorem ( $|\mathbb{R}|$ (1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

## Theorem ( $|\mathbb{R}|$ (1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

## Theorem ( $|\mathbb{R}|$ (1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots)$$

## Theorem ( $|\mathbb{R}|$ (1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$





## Theorem ( $|\mathbb{R}|$ (1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$



*“Je le vois, mais je ne le crois pas !”*

*“I see it, but I don't believe it !”*

— Cantor's letter to Dedekind (1877).

## Theorem ( $|\mathbb{R}|$ (1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$




*“Je le vois, mais je ne le crois pas !”*

*“I see it, but I don't believe it !”*

— Cantor's letter to Dedekind (1877).

*Q : Then, what is “dimension”?*



Definition ( $|A| \leq |B|$ )

$|A| \leq |B|$  if there exists an *one-to-one* function  $f$  from  $A$  into  $B$ .

Definition ( $|A| \leq |B|$ )

$|A| \leq |B|$  if there exists an *one-to-one* function  $f$  from  $A$  into  $B$ .

bijection  $f : A \rightarrow f(A) (\subseteq B)$

Definition ( $|A| \leq |B|$ )

$|A| \leq |B|$  if there exists an *one-to-one* function  $f$  from  $A$  into  $B$ .

bijection  $f : A \rightarrow f(A) (\subseteq B)$

*Q : What about onto function  $f : A \rightarrow B$ ?*

Definition ( $|A| \leq |B|$ )

$|A| \leq |B|$  if there exists an *one-to-one* function  $f$  from  $A$  into  $B$ .

bijection  $f : A \rightarrow f(A) (\subseteq B)$

*Q : What about onto function  $f : A \rightarrow B$ ?*

$$|B| \leq |A|$$

Definition ( $|A| \leq |B|$ )

$|A| \leq |B|$  if there exists an *one-to-one* function  $f$  from  $A$  into  $B$ .

bijection  $f : A \rightarrow f(A) (\subseteq B)$

*Q : What about onto function  $f : A \rightarrow B$ ?*

$|B| \leq |A|$  (Axiom of Choice)



Definition ( $|A| < |B|$ )

$$|A| < |B| \iff |A| \leq |B| \wedge |A| \neq |B|$$

Definition ( $|A| < |B|$ )

$$|A| < |B| \iff |A| \leq |B| \wedge |A| \neq |B|$$

$$|\mathbb{N}| < |\mathbb{R}|$$

$$|X| < |2^X|$$

## Definition (Countable Revisited)

$X$  is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

## Definition (Countable Revisited)

$X$  is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

## Theorem (Proof for Countable (UD Exercise 22.5))

$X$  is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

## Definition (Countable Revisited)

$X$  is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

## Theorem (Proof for Countable (UD Exercise 22.5))

$X$  is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

$X$  is countable iff

$$|X| \leq |\mathbb{N}|.$$

## Definition (Countable Revisited)

$X$  is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

## Theorem (Proof for Countable (UD Exercise 22.5))

$X$  is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

$X$  is countable iff

$$|X| \leq |\mathbb{N}|.$$

## Subsets of Countable Set (UD 22.6; UD Corollary 22.4)

Every subset  $B$  of a countable set  $A$  is countable.

## Set Union (UD 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.

## Set Union (UD 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.
- (b) a countably infinite collection of nonempty sets whose union is finite.



## Set Union (UD 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.
- (b) a countably infinite collection of nonempty sets whose union is finite.

$$\left( \{A_i : i \in \mathbb{R}\} \quad A_i = \{1\} \right) = \{\{1\}\}$$

## Set Union (UD 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.
- (b) a countably infinite collection of nonempty sets whose union is finite.

$$\left( \{A_i : i \in \mathbb{R}\} \quad A_i = \{1\} \right) = \{\{1\}\}$$

$$|A| = n \implies |2^A| = 2^n$$

Slope (UD 22.2 ( $e$ ))

( $e$ ) the set of all lines with rational slopes

Slope (UD 22.2 (e))

(e) the set of all lines with rational slopes

$$(\mathbb{Q}, \mathbb{R})$$

Slope (UD 22.2 (e))

(e) the set of all lines with rational slopes

$$(\mathbb{Q}, \mathbb{R})$$

$$|\mathbb{R}| \leq |\mathbb{Q} \times \mathbb{R}| \leq |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}|$$

*Q : Is " $\leq$ " a partial order?*

*Q : Is " $\leq$ " a partial order?*

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

*Q : Is “ $\leq$ ” a partial order?*

Theorem (Cantor-Schröder–Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$




*Q : Is “ $\leq$ ” a partial order?*

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$




*Q : Is “ $\leq$ ” a partial order?*

Theorem (Cantor-Schröder–Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$




*Q : Is “ $\leq$ ” a partial order?*

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$



*Q : Is " $\leq$ " a total order?*

*Q : Is " $\leq$ " a total order?*

Theorem (PCC)

*Principle of Cardinal Comparability (PCC)  $\iff$  Axiom of Choice*

# Finite Sets



# Finite Sets



“关于有穷，我原以为我是懂的”

## Definition (Finite)

$X$  is finite if

$$\exists n \in \mathbb{N} : |X| = n$$



## Definition (Finite)

$X$  is finite if

$$\exists n \in \mathbb{N} : |X| = n$$

## Theorem (Pigeonhole Principle (UD Theorem 21.2))

$$f : \{1, \dots, m\} \rightarrow \{1, \dots, n\} \quad (m, n \in \mathbb{N}^+, m > n)$$

$f$  is not one-to-one.

## $A \setminus \{a\}$ (UD Problem 21.15)

Let  $A$  be a nonempty finite set with  $|A| = n$  and let  $a \in A$ .

Prove that  $A \setminus \{a\}$  is finite and  $|A \setminus \{a\}| = n - 1$ .

## $A \setminus \{a\}$ (UD Problem 21.15)

Let  $A$  be a nonempty finite set with  $|A| = n$  and let  $a \in A$ .

Prove that  $A \setminus \{a\}$  is finite and  $|A \setminus \{a\}| = n - 1$ .

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

## $A \setminus \{a\}$ (UD Problem 21.15)

Let  $A$  be a nonempty finite set with  $|A| = n$  and let  $a \in A$ .

Prove that  $A \setminus \{a\}$  is finite and  $|A \setminus \{a\}| = n - 1$ .

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$$f|_{A \setminus \{a\}} : A \setminus \{a\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\} \setminus \{f(a)\}$$

## $A \setminus \{a\}$ (UD Problem 21.15)

Let  $A$  be a nonempty finite set with  $|A| = n$  and let  $a \in A$ .

Prove that  $A \setminus \{a\}$  is finite and  $|A \setminus \{a\}| = n - 1$ .

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$$f|_{A \setminus \{a\}} : A \setminus \{a\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\} \setminus \{f(a)\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n-1\}$$

$|A| \leq |B|$  (UD Problem 21.17)

$A$  and  $B$  are finite sets and  $f : A \rightarrow B$  is one-to-one.

Show that  $|A| \leq |B|$ .

$|A| \leq |B|$  (UD Problem 21.17)

$A$  and  $B$  are finite sets and  $f : A \rightarrow B$  is one-to-one.  
Show that  $|A| \leq |B|$ .



$|A| \leq |B|$  (UD Problem 21.17)

$A$  and  $B$  are finite sets and  $f : A \rightarrow B$  is one-to-one.  
Show that  $|A| \leq |B|$ .



By contradiction and the pigeonhole principle.



(UD Problem 21.16)

- (a)  $A$  is a finite set and  $B \subseteq A$ . We showed that  $B$  is finite (Corollary 20.11). Show that  $|B| \leq |A|$ .

(UD Problem 21.16)

- (a)  $A$  is a finite set and  $B \subseteq A$ . We showed that  $B$  is finite (Corollary 20.11). Show that  $|B| \leq |A|$ .

one-to-one  $f : B \rightarrow A$

(UD Problem 21.16)

- (a)  $A$  is a finite set and  $B \subseteq A$ . We showed that  $B$  is finite (Corollary 20.11). Show that  $|B| \leq |A|$ .

one-to-one  $f : B \rightarrow A$

- (b)  $A$  is a finite set and  $B \subseteq A$ . Show that if  $B \neq A$ , then  $|B| < |A|$ .

(UD Problem 21.16)

- (a)  $A$  is a finite set and  $B \subseteq A$ . We showed that  $B$  is finite (Corollary 20.11). Show that  $|B| \leq |A|$ .

one-to-one  $f : B \rightarrow A$

- (b)  $A$  is a finite set and  $B \subseteq A$ . Show that if  $B \neq A$ , then  $|B| < |A|$ .

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

### (UD Problem 21.16)

- (a)  $A$  is a finite set and  $B \subseteq A$ . We showed that  $B$  is finite (Corollary 20.11). Show that  $|B| \leq |A|$ .

one-to-one  $f : B \rightarrow A$

- (b)  $A$  is a finite set and  $B \subseteq A$ . Show that if  $B \neq A$ , then  $|B| < |A|$ .

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

- (c) If two finite sets  $A$  and  $B$  satisfy  $B \subseteq A$  and  $|A| \leq |B|$ , then  $A = B$ .

(UD Problem 21.16)

- (a)  $A$  is a finite set and  $B \subseteq A$ . We showed that  $B$  is finite (Corollary 20.11). Show that  $|B| \leq |A|$ .

one-to-one  $f : B \rightarrow A$

- (b)  $A$  is a finite set and  $B \subseteq A$ . Show that if  $B \neq A$ , then  $|B| < |A|$ .

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

- (c) If two finite sets  $A$  and  $B$  satisfy  $B \subseteq A$  and  $|A| \leq |B|$ , then  $A = B$ .

By contradiction and (b).

## Cardinality of $|\text{ran}(f)|$ (UD Problem 21.18)

Let  $A$  and  $B$  be sets with  $A$  finite.

$$f : A \rightarrow B$$

Prove that  $|\text{ran}(f)| \leq |A|$ .

## Cardinality of $|\text{ran}(f)|$ (UD Problem 21.18)

Let  $A$  and  $B$  be sets with  $A$  finite.

$$f : A \rightarrow B$$

Prove that  $|\text{ran}(f)| \leq |A|$ .

one-to-one  $g : \text{ran}(f) \rightarrow A$



## Cardinality of $|\text{ran}(f)|$ (UD Problem 21.18)

Let  $A$  and  $B$  be sets with  $A$  finite.

$$f : A \rightarrow B$$

Prove that  $|\text{ran}(f)| \leq |A|$ .

one-to-one  $g : \text{ran}(f) \rightarrow A$

(No Axiom of Choice Here)

$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$f$  is one-to-one  $\iff f$  is onto.

$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$\implies$

By contradiction.

$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$\implies$

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$



By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$$\iff$$

$$\implies$$

$$\forall y \in A \exists x \in A : y = f(x)$$

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$$\iff$$

$$\implies$$

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$$\forall y \in A \exists x \in A : y = f(x)$$

$$\forall y, \text{ choose } x : (g : g(y) = x)$$

$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$\iff$

$\implies$

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$$\forall y \in A \exists x \in A : y = f(x)$$

$$\forall y, \text{ choose } x : (g : g(y) = x)$$

$g$  is bijective.



$f : A \rightarrow A$  (UD Problem 21.19)

Let  $A$  be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$\Leftarrow$

$\Rightarrow$

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$$\forall y \in A \exists x \in A : y = f(x)$$

$$\forall y, \text{ choose } x : (g : g(y) = x)$$

$g$  is bijective.

$$f(g(y)) = f(x) = y \implies f = g^{-1}$$

# Dangerous Knowledge (BBC 2007)



Continuum Hypothesis (CH):

$$c = \aleph_1$$

Continuum Hypothesis (CH):

$$\boxed{c = \aleph_1}$$

$$c = 2^{\aleph_0} = \aleph_1$$

Continuum Hypothesis (CH):

$$\boxed{c = \aleph_1}$$

$$c = 2^{\aleph_0} = \aleph_1$$



Dangerous Knowledge (22:20)

Continuum Hypothesis (CH):

$$c = \aleph_1$$

$$c = 2^{\aleph_0} = \aleph_1$$


 Dangerous Knowledge (22:20)

Independence from ZFC:

Kurt Gödel (1940) CH cannot be disproved from ZF.

Paul Cohen (1964) CH cannot be proven from the ZFC axioms.

Thank  
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn