

## 2-9 Sorting and Selection

Hengfeng Wei

hfwei@nju.edu.cn

May 28, 2018




## How to Argue?




Am I Alone?



## QUICKSORT Invented by Tony Hoare in 1959/1960




## QUICKSORT Invented by Tony Hoare in 1959/1960



`null pointer`

## QUICKSORT Invented by Tony Hoare in 1959/1960



`null pointer`

*"I call it my billion-dollar mistake."*

## Best-Case Complexity of QUICKSORT (7.4 – 2)

**Show that** QUICKSORT's best-case running time is  $\Omega(n \log n)$ .

## Best-Case Complexity of QUICKSORT (7.4 – 2)

**Show that** QUICKSORT's best-case running time is  $\Omega(n \log n)$ .

$$T(n) = 2T\left(\frac{n}{2}\right) + \Theta(n)$$



## Best-Case Complexity of QUICKSORT (7.4 – 2)

**Show that** QUICKSORT's best-case running time is  $\Omega(n \log n)$ .

$$T(n) = 2T\left(\frac{n}{2}\right) + \Theta(n)$$

$$T(n) = \underbrace{n}_{\text{PARTITION}} \underbrace{\log n}_{\text{Height}} \quad (\text{Recursion Tree})$$

## Best-Case Complexity of QUICKSORT (7.4 – 2)

**Show that** QUICKSORT's best-case running time is  $\Omega(n \log n)$ .

$$T(n) = 2T\left(\frac{n}{2}\right) + \Theta(n)$$

$$T(n) = \underbrace{n}_{\text{PARTITION}} \underbrace{\log n}_{\text{Height}} \quad (\text{Recursion Tree})$$

$$T(n) = \min_{0 \leq q \leq n-1} \left( T(q) + T(n - q - 1) \right) + \Theta(n)$$

## Best-Case Complexity of QUICKSORT (7.4 – 2)

**Show that** QUICKSORT's best-case running time is  $\Omega(n \log n)$ .

$$T(n) = 2T\left(\frac{n}{2}\right) + \Theta(n)$$

$$T(n) = \underbrace{n}_{\text{PARTITION}} \underbrace{\log n}_{\text{Height}} \quad (\text{Recursion Tree})$$

$$T(n) = \min_{0 \leq q \leq n-1} \left( T(q) + T(n - q - 1) \right) + \Theta(n)$$

$$T(n) = \Omega(n \log n)$$

## Best-Case Complexity of QUICKSORT (7.4 – 2)

**Show that** QUICKSORT's best-case running time is  $\Omega(n \log n)$ .

$$T(n) = 2T\left(\frac{n}{2}\right) + \Theta(n)$$

$$T(n) = \underbrace{n}_{\text{PARTITION}} \underbrace{\log n}_{\text{Height}} \quad (\text{Recursion Tree})$$

$$T(n) = \min_{0 \leq q \leq n-1} \left( T(q) + T(n - q - 1) \right) + \Theta(n)$$

$$T(n) = \Omega(n \log n)$$

By substitution.

## Median-of-3 PARTITION (Problem 7 – 5)

**Argue that** in the  $\Omega(n \log n)$  running time of QUICKSORT, the *median-of-3* method affects only the constant factor.

## Median-of-3 PARTITION (Problem 7 – 5)

**Argue that** in the  $\Omega(n \log n)$  running time of QUICKSORT, the *median-of-3* method affects only the constant factor.



## Median-of-3 PARTITION (Problem 7 – 5)

**Argue that** in the  $\Omega(n \log n)$  running time of QUICKSORT, the *median-of-3* method affects only the constant factor.



$$T(n) = \min_{0 \leq q \leq n-1} (T(q) + T(n - q - 1)) + \Theta(n)$$

$$T(n) = \Omega(n \log n)$$






## The $\frac{n}{k}$ -sorted Problem (Problem 8.1 – 4)

Sorts an already  $\frac{n}{k}$ -sorted array

$n$  elements

$\frac{n}{k}$  blocks



$k$  elements

<


not sorted

## The $\frac{n}{k}$ -sorted Problem (Problem 8.1 – 4)

Sorts an already  $\frac{n}{k}$ -sorted array

$n$  elements

$\frac{n}{k}$  blocks



$k$  elements

not sorted


$$\Omega(n \log k)$$

## The $\frac{n}{k}$ -sorted Problem (Problem 8.1 – 4)

Sorts an already  $\frac{n}{k}$ -sorted array

$n$  elements

$\frac{n}{k}$  blocks



$k$  elements

not sorted


$$\Omega(n \log k) \quad O(n \log k)$$

## The $\frac{n}{k}$ -sorted Problem (Problem 8.1 – 4)

Sorts an already  $\frac{n}{k}$ -sorted array

$n$  elements

$\frac{n}{k}$  blocks




$k$  elements

not sorted


$$\Omega(n \log k) \quad O(n \log k)$$

$$(k!)^{\frac{n}{k}} \leq L \leq 2^H$$

$\frac{n}{k}$ -sorts an arbitrary array




$\frac{n}{k}$ -sorts an arbitrary array




$O(?)$

$\frac{n}{k}$ -sorts an arbitrary array



$O(?)$      $\Omega(?)$

$\frac{n}{k}$ -sorts an arbitrary array




$O(?)$      $\Omega(?)$

$$L \geq \binom{n}{\underbrace{k, \dots, k}_{\frac{n}{k}}} = \frac{n!}{(k!)^{\frac{n}{k}}}$$



$\frac{n}{k}$ -sorts an arbitrary array



$O(?)$      $\Omega(?)$

$$L \geq \binom{n}{\underbrace{k, \dots, k}_{\frac{n}{k}}} = \frac{n!}{(k!)^{\frac{n}{k}}} \implies \Omega(n \log(n/k))$$

## Sorting $[0, n^3 - 1]$ (Problem 8.3 - 4)

Sort  $n$  integers in  $[0, n^3 - 1]$  in  $O(n)$  time.

## Sorting in Place in Linear Time (Problem 8 – 2 (e))

Suppose that the  $n$  records have keys in the range  $[0, k]$ .

Modify COUNTING-SORT to sort them **in place** ( $O(k)$ ) in  $O(n + k)$  time.

## Sorting in Place in Linear Time (Problem 8 – 2 (e))

Suppose that the  $n$  records have keys in the range  $[0, k]$ .

Modify COUNTING-SORT to sort them **in place** ( $O(k)$ ) in  $O(n + k)$  time.

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| A: | 2 | 5 | 3 | 0 | 2 | 3 | 0 | 3 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
|    | 0 | 1 | 2 | 3 | 4 | 5 |
| C: | 2 | 0 | 2 | 3 | 0 | 1 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
| C: | 2 | 2 | 4 | 7 | 7 | 8 |
|----|---|---|---|---|---|---|

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| B: | 0 | 0 | 2 | 2 | 3 | 3 | 3 | 5 |

## Sorting in Place in Linear Time (Problem 8 – 2 (e))

Suppose that the  $n$  records have keys in the range  $[0, k]$ .

Modify COUNTING-SORT to sort them **in place** ( $O(k)$ ) in  $O(n + k)$  time.

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| A: | 2 | 5 | 3 | 0 | 2 | 3 | 0 | 3 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
|    | 0 | 1 | 2 | 3 | 4 | 5 |
| C: | 2 | 0 | 2 | 3 | 0 | 1 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
| C: | 2 | 2 | 4 | 7 | 7 | 8 |
|----|---|---|---|---|---|---|

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| B: | 0 | 0 | 2 | 2 | 3 | 3 | 3 | 5 |

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| A: | 2 | 0 | 3 | 0 | 2 | 3 | 3 | 5 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
|    | 0 | 1 | 2 | 3 | 4 | 5 |
| C: | 1 | 2 | 4 | 7 | 7 | 8 |

## Sorting in Place in Linear Time (Problem 8 – 2 (e))

Suppose that the  $n$  records have keys in the range  $[0, k]$ .

Modify COUNTING-SORT to sort them **in place** ( $O(k)$ ) in  $O(n + k)$  time.

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| A: | 2 | 5 | 3 | 0 | 2 | 3 | 0 | 3 |


|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
|    | 0 | 1 | 2 | 3 | 4 | 5 |
| C: | 2 | 0 | 2 | 3 | 0 | 1 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
| C: | 2 | 2 | 4 | 7 | 7 | 8 |
|----|---|---|---|---|---|---|

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| B: | 0 | 0 | 2 | 2 | 3 | 3 | 3 | 5 |

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| A: | 2 | 0 | 3 | 0 | 2 | 3 | 3 | 5 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
|    | 0 | 1 | 2 | 3 | 4 | 5 |
| C: | 1 | 2 | 4 | 7 | 7 | 8 |



## Sorting in Place in Linear Time (Problem 8 – 2 (e))

Suppose that the  $n$  records have keys in the range  $[0, k]$ .

Modify COUNTING-SORT to sort them **in place** ( $O(k)$ ) in  $O(n + k)$  time.

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| A: | 2 | 5 | 3 | 0 | 2 | 3 | 0 | 3 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
|    | 0 | 1 | 2 | 3 | 4 | 5 |
| C: | 2 | 0 | 2 | 3 | 0 | 1 |


|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
| C: | 2 | 2 | 4 | 7 | 7 | 8 |
|----|---|---|---|---|---|---|

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| B: | 0 | 0 | 2 | 2 | 3 | 3 | 3 | 5 |

|    |   |   |   |   |   |   |   |   |
|----|---|---|---|---|---|---|---|---|
|    | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| A: | 2 | 0 | 3 | 0 | 2 | 3 | 3 | 5 |

|    |   |   |   |   |   |   |
|----|---|---|---|---|---|---|
|    | 0 | 1 | 2 | 3 | 4 | 5 |
| C: | 1 | 2 | 4 | 7 | 7 | 8 |

While ( $i \geq 1$ ):



1 2 3 4 5 6 7 8  
 A: 


|   |   |   |   |   |   |   |   |
|---|---|---|---|---|---|---|---|
| 2 | 0 | 3 | 0 | 2 | 3 | 3 | 5 |
|---|---|---|---|---|---|---|---|

0 1 2 3 4 5  
 C: 

|   |   |   |   |   |   |
|---|---|---|---|---|---|
| 1 | 2 | 4 | 7 | 7 | 8 |
|---|---|---|---|---|---|

While ( $i \geq 1$ ):

Code here





## Finding the 2nd Smallest Element (Problem 9.1 – 1)

Show that the 2nd smallest of  $n$  elements can be found with  $n + \lceil \log n \rceil - 2$  comparisons in the worst case.

## Finding the 2nd Smallest Element (Problem 9.1 – 1)


Show that the 2nd smallest of  $n$  elements can be found with  $n + \lceil \log n \rceil - 2$  comparisons in the worst case.

$$(n - 1) + (n - 1 - 1) = 2n - 3$$


$$n + \lceil \log n \rceil - 2 = (n - 1) + (\lceil \log n \rceil - 1)$$



$$n + \lceil \log n \rceil - 2 = (n - 1) + (\lceil \log n \rceil - 1)$$




$$n + \lceil \log n \rceil - 2 = (n - 1) + (\lceil \log n \rceil - 1)$$



#**Potential** 2nd smallest elements  $\leq \lceil \log n \rceil$

$$n + \lceil \log n \rceil - 2 = (n - 1) + (\lceil \log n \rceil - 1)$$



#**Potential** 2nd smallest elements  $\leq \lceil \log n \rceil$


**Q** : Can we do even better?

$$\Omega = n + \lceil \log n \rceil - 2$$



$$\Omega = n + \lceil \log n \rceil - 2 = (n - 1) + (\lceil \log n \rceil - 1)$$

$$\Omega = n + \lceil \log n \rceil - 2 = (n - 1) + (\lceil \log n \rceil - 1)$$



## TAOCP Vol 3 (Page 209, Section 5.3.3)

## $k$ Numbers Closest to the Median (Problem 9.3 – 7)

$S : n$  distinct numbers       $k \leq n$

## $k$ Numbers Closest to the Median (Problem 9.3 – 7)

$S : n$  distinct numbers       $k \leq n$

$$\frac{n}{2} - \frac{k}{2} \sim \frac{n}{2} + \frac{k}{2}$$

## $k$ Numbers Closest to the Median (Problem 9.3 – 7)

$S$  :  $n$  distinct numbers       $k \leq n$

$$\frac{n}{2} - \frac{k}{2} \sim \frac{n}{2} + \frac{k}{2}$$

$$S = \{800, 6, 900, \textcolor{red}{50}, 7\}, \quad k = 2 \implies \{6, 7\}$$

## $k$ Numbers Closest to the Median (Problem 9.3 – 7)

$S$  :  $n$  distinct numbers       $k \leq n$

$$\frac{n}{2} - \frac{k}{2} \sim \frac{n}{2} + \frac{k}{2}$$

$$S = \{800, 6, 900, \textcolor{red}{50}, 7\}, \quad k = 2 \implies \{6, 7\}$$

$$S - 50 = \{750, -44, 850, \textcolor{red}{0}, -43\}$$

## $k$ Numbers Closest to the Median (Problem 9.3 – 7)

$S$  :  $n$  distinct numbers       $k \leq n$

$$\frac{n}{2} - \frac{k}{2} \sim \frac{n}{2} + \frac{k}{2}$$


$$S = \{800, 6, 900, \textcolor{red}{50}, 7\}, \quad k = 2 \implies \{6, 7\}$$

$$S - 50 = \{750, -44, 850, \textcolor{red}{0}, -43\}$$

median + subtraction +  $(k + 1)$ -th smallest + partition + add

Thank  
You!





Office 302

Mailbox: H016

hfwei@nju.edu.cn