

# 3-11 Matchings and Factors

## (Part II: Perfect Matchings)

Hengfeng Wei

hfwei@nju.edu.cn

December 24, 2018




# Chinese Postman Problem (CPP)



Shortest Simple Path in Undirected Graphs

# Chinese Postman Problem (CPP)



Shortest Simple Path in Undirected Graphs

# Chinese Postman Problem (CPP)

## (Postman Tour Problem, Route Inspection Problem)





管梅谷(1934-)

第 10 卷 第 3 期  
1960 年 12 月

数 学 学 报  
ACTA MATHEMATICA SINICA

Vol. 10, No. 3  
Dec., 1960

## 奇偶点图上作业法\*


管 梅 谷  
(山东师范学院)

### §1. 問題的提出

在邮局搞綫性规划时,发现了下述問題:“一个投递員每次上班,要走遍他負責送信的段<sup>1</sup>,然后回到邮局,問應該怎样走才能使所走的路程最短。”

《奇偶点图上作业法》, 1960

Translated into English in 1962



Jack Edmonds (1934-)

## MATCHING, EULER TOURS AND THE CHINESE POSTMAN

Jack EDMONDS

*University of Waterloo, Waterloo, Ontario, Canada*

and

Ellis L. JOHNSON

*IBM Watson Research Center, Yorktown Heights, New York, U.S.A.*

Received 20 May 1972

Revised manuscript received 3 April 1973

The solution of the Chinese postman problem using matching theory is given. The convex hull of integer solutions is described as a linear programming polyhedron. This polyhedron is used to show that a good algorithm gives an optimum solution. The algorithm is a specialization of the more general  $b$ -matching blossom algorithm. Algorithms for finding Euler tours and related problems are also discussed.

“Matching, Euler Tours and the Chinese Postman”, 1973 (1965)

## Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find the shortest tour such that each edge is traversed at least once.

## Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find the shortest tour such that each edge is traversed at least once.

*Q* : What is the relation between Postman Tour and Eulerian Tour?



## $P$ : A postman tour of $G$

## $P$ : A postman tour of $G$

$P$  contains every edge  $e$  at least once.

## $P$ : A postman tour of $G$

$P$  contains every edge  $e$  at least once.

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

## $P$ : A postman tour of $G$

$P$  contains every edge  $e$  at least once.

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

Construct  $G' = G + e \cdot x_e$

## $P$ : A postman tour of $G$

$P$  contains every edge  $e$  at least once.

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

Construct  $G' = G + e \cdot x_e$

$P$  is an Eulerian tour of  $G'$ .

## $P$ : A postman tour of $G$

$P$  contains every edge  $e$  at least once.

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

Construct  $G' = G + e \cdot x_e$

$P$  is an Eulerian tour of  $G'$ .

### Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find  $x_e \in \mathbb{N}$  for each edge  $e$  of  $G$

to minimize  $\sum_e w(e)x_e$ ,

such that  $G' = G + e \cdot x_e$  is an Eulerian graph.

## Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find  $x_e \in \mathbb{N}$  for each edge  $e$  of  $G$

to minimize  $\sum_e w(e)x_e$ ,

such that  $G' = G + e \cdot x_e$  is an Eulerian graph.

## Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find  $x_e \in \mathbb{N}$  for each edge  $e$  of  $G$

to minimize  $\sum_e w(e)x_e$ ,

such that  $G' = G + e \cdot x_e$  is an Eulerian graph.

*Q* : What are the possible values of each  $x_e$ ?

## Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find  $x_e \in \mathbb{N}$  for each edge  $e$  of  $G$

$$\text{to minimize } \sum_e w(e)x_e,$$

such that  $G' = G + e \cdot x_e$  is an Eulerian graph.

*Q* : What are the possible values of each  $x_e$ ?

## Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find  $x_e \in \{0, 1\}$  for each edge  $e$  of  $G$

$$\text{to minimize } \sum_e w(e)x_e,$$

such that  $G' = G + e \cdot x_e$  is an Eulerian graph.

---


---



1:  $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$




- 
- 
- 1:  $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
  - 2: Construct a complete weighted graph  $G_p$  with vertices  $V_o$ :
  - 3: **for**  $u, v \in V_o$  **do**
  - 4:      $w(u, v) \leftarrow$  the length of the shortest path between  $u$  and  $v$




- 
- 
- 1:  $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
  - 2: Construct a complete weighted graph  $G_p$  with vertices  $V_o$ :
  - 3: **for**  $u, v \in V_o$  **do**
  - 4:      $w(u, v) \leftarrow$  the length of the shortest path between  $u$  and  $v$
  - 5: Find a **minimum-weighted perfect matching**  $M$  of  $G_p$

- 
- 1:  $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
  - 2: Construct a complete weighted graph  $G_p$  with vertices  $V_o$ :
  - 3: **for**  $u, v \in V_o$  **do**
  - 4:      $w(u, v) \leftarrow$  the length of the shortest path between  $u$  and  $v$
  - 5: Find a **minimum-weighted perfect matching**  $M$  of  $G_p$
  - 6: **for**  $(u, v) \in M$  **do**
  - 7:      $p \leftarrow$  the shortest path between  $u$  and  $v$
  - 8:      $\forall e \in p : x_e \leftarrow 1$
-









- 
- 1:  $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
  - 2: Construct a complete weighted graph  $G_p$  with vertices  $V_o$ :
  - 3: **for**  $u, v \in V_o$  **do**
  - 4:      $w(u, v) \leftarrow$  the length of the shortest path between  $u$  and  $v$
  - 5: Find a minimum-weighted perfect matching  $M$  of  $G_p$
  - 6: **for**  $(u, v) \in M$  **do**
  - 7:      $p \leftarrow$  the shortest path between  $u$  and  $v$
  - 8:      $\forall e \in p : x_e \leftarrow 1$
-

- 
- 1:  $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
  - 2: Construct a complete weighted graph  $G_p$  with vertices  $V_o$ :
  - 3: **for**  $u, v \in V_o$  **do**
  - 4:      $w(u, v) \leftarrow$  the length of the shortest path between  $u$  and  $v$
  - 5: Find a minimum-weighted perfect matching  $M$  of  $G_p$
  - 6: **for**  $(u, v) \in M$  **do**
  - 7:      $p \leftarrow$  the shortest path between  $u$  and  $v$
  - 8:      $\forall e \in p : x_e \leftarrow 1$
- 

*Q* : What if some edge  $e \in E(G)$  is in two shortest paths corresponding to (two) matching edges of  $G_p$ ?

## Theorem (Edge-disjointness of Shortest Paths)

*No edge  $e \in E(G)$  is in two shortest paths corresponding to (two) matching edges of  $G_p$ .*

## Theorem (Edge-disjointness of Shortest Paths)

*No edge  $e \in E(G)$  is in two shortest paths corresponding to (two) matching edges of  $G_p$ .*

Proof.

## Theorem (Edge-disjointness of Shortest Paths)

*No edge  $e \in E(G)$  is in two shortest paths corresponding to (two) matching edges of  $G_p$ .*

Proof.


By Contradiction.

## Theorem (Edge-disjointness of Shortest Paths)

No edge  $e \in E(G)$  is in two shortest paths corresponding to (two) matching edges of  $G_p$ .

Proof.

By Contradiction.



Suppose that


$$\exists e \in E(G) : e \in u_1 \rightsquigarrow u_2 \wedge e \in v_1 \rightsquigarrow v_2$$

## Theorem (Edge-disjointness of Shortest Paths)

No edge  $e \in E(G)$  is in two shortest paths corresponding to (two) matching edges of  $G_p$ .

Proof.

By Contradiction.



Suppose that

$$\exists e \in E(G) : e \in u_1 \rightsquigarrow u_2 \wedge e \in v_1 \rightsquigarrow v_2$$

Contradiction:

$u_1 \rightsquigarrow v_1, u_2 \rightsquigarrow v_2 \implies$  smaller perfect matching



## Theorem (Property of CHINESE-POSTMAN)

*The edges with  $x_e = 1$  obtained by CHINESE-POSTMAN is a **minimum** collection of edge-disjoint paths connecting pairs of odd vertices.*

## Theorem (Property of CHINESE-POSTMAN)

*The edges with  $x_e = 1$  obtained by CHINESE-POSTMAN is a **minimum** collection of edge-disjoint paths connecting pairs of odd vertices.*



## Theorem (Property of CHINESE-POSTMAN)

*The edges with  $x_e = 1$  obtained by CHINESE-POSTMAN is a **minimum** collection of edge-disjoint paths connecting pairs of odd vertices.*

To prove that these  $x_e = 1$  obtained by CHINESE-POSTMAN satisfies:

## Definition (Chinese Postman Problem)

Given an undirected weighted graph  $G$  with  $w(e) > 0$ ,  
to find  $x_e \in \{0, 1\}$  for each edge  $e$  of  $G$

to minimize  $\sum_e w(e)x_e$ ,

such that  $G' = G + e \cdot x_e$  is an Eulerian graph.

Lemma (CHINESE-POSTMAN Gives a Postman Tour)

$G' = G + e \cdot x_e$  is an Eulerian graph.

Lemma (CHINESE-POSTMAN Gives a Postman Tour)

$G' = G + e \cdot x_e$  is an Eulerian graph.

Proof.

A collection of edge-disjoint paths connecting pairs of odd vertices.  $\square$

## Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$\sum_e w(e)x_e$  is minimized.

## Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$\sum_e w(e)x_e$  is minimized.

## Theorem (Property of CHINESE-POSTMAN)

*The edges with  $x_e = 1$  obtained by CHINESE-POSTMAN is a minimum collection of edge-disjoint simple paths connecting pairs of odd vertices.*

## Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$\sum_e w(e)x_e$  is minimized.

## Theorem (Property of CHINESE-POSTMAN)

*The edges with  $x_e = 1$  obtained by CHINESE-POSTMAN is a minimum collection of edge-disjoint simple paths connecting pairs of odd vertices.*

$P$ : An optimal postman tour of  $G$

## Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$\sum_e w(e)x_e$  is minimized.

## Theorem (Property of CHINESE-POSTMAN)

*The edges with  $x_e = 1$  obtained by CHINESE-POSTMAN is a minimum collection of edge-disjoint simple paths connecting pairs of odd vertices.*

$P$ : An optimal postman tour of  $G$

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

## Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$\sum_e w(e)x_e$  is minimized.

## Theorem (Property of CHINESE-POSTMAN)

*The edges with  $x_e = 1$  obtained by CHINESE-POSTMAN is a minimum collection of edge-disjoint simple paths connecting pairs of odd vertices.*

$P$ : An optimal postman tour of  $G$

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

We show that the edges with  $x_e = 1$  is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

## Lemma (Property of Optimal Postman Tours)

*P: An optimal postman tour of G*

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

The edges with  $x_e = 1$  is a collection of  
edge-disjoint simple paths connecting pairs of odd vertices.

## Lemma (Property of Optimal Postman Tours)

*P: An optimal postman tour of G*

*Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge e is in P.*

*The edges with  $x_e = 1$  is a collection of  
edge-disjoint simple paths connecting pairs of odd vertices.*

Proof.

By Construction.

## Lemma (Property of Optimal Postman Tours)

*P: An optimal postman tour of G*

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

The edges with  $x_e = 1$  is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

Proof.

By Construction.

An odd number of edges  $e$  with  $x_e = 1$  meet odd nodes.

An even number of edges  $e$  with  $x_e = 1$  meet even nodes.

## Lemma (Property of Optimal Postman Tours)

*P: An optimal postman tour of G*

Let  $1 + x_e$  ( $x_e \in \mathbb{N}$ ) be the number of times edge  $e$  is in  $P$ .

The edges with  $x_e = 1$  is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

Proof.


By Construction.

An odd number of edges  $e$  with  $x_e = 1$  meet odd nodes.

An even number of edges  $e$  with  $x_e = 1$  meet even nodes.







Office 302

Mailbox: H016

hfwei@nju.edu.cn