

A Simple Proof of Menger's Theorem

William McCuaig

DEPARTMENT OF MATHEMATICS
SIMON FRASER UNIVERSITY, BURNABY
BRITISH COLUMBIA, CANADA

ABSTRACT

A proof of Menger's theorem is presented.

We use the notation and terminology of Bondy and Murty [1].

Let D be a directed graph. If $\{u\}$, $\{v\}$, and S are disjoint subsets of $V(D)$ and u and v are nonadjacent, then S separates u and v if every (u, v) -path has a vertex in S .

Proofs of Menger's theorem are given in [2–14].

Menger's Theorem. If no set of fewer than n vertices separates nonadjacent vertices u and v in a directed graph D , then there are n internally disjoint (u, v) -paths.

Proof. The proof uses induction on n . The theorem is trivial for $n = 1$. Suppose u and v are separated by no set of less than $n + 1$ vertices ($n \geq 1$). By the induction hypothesis there are n internally disjoint (u, v) -paths P_1, \dots, P_n . Since the set of second vertices of P_1, \dots, P_n does not separate u and v , there is a (u, v) -path P whose initial arc is not on $P_i, i = 1, \dots, n$. Let x be the first vertex on P after u which is also on some $P_i, 1 \leq i \leq n$. Let P_{n+1} be the (u, x) -section of P . Assume P_1, \dots, P_n, P_{n+1} have been chosen so that the distance in $D - \{u\}$ from x to v is a minimum. If $x = v$ we are done, so assume not.

In $D - \{x\}$ there are n internally disjoint (u, v) -paths Q_1, \dots, Q_n , again by the induction hypothesis. Assume Q_1, \dots, Q_n have been chosen so that a minimum number of arcs in $B = A(D) - \bigcup_{i=1}^{n+1} A(P_i)$ are used. Let H be the directed graph consisting of the vertices and arcs of $Q_1,$

..., Q_n together with the vertex x . Choose some P_k , $1 \leq k \leq n + 1$, whose initial arc is not in $A(H)$. Let y be the first vertex on P_k after u which is in $V(H)$. If $y = v$ we are done, so assume not.

If $y = x$ then let R be the shortest (x, v) -path in $D - \{u\}$. Let z be the first vertex of R on some Q_j , $1 \leq j \leq n$. Then the distance in $D - \{u\}$ from z to v is less than the distance from x to v . This contradicts the choice of P_1, \dots, P_n, P_{n+1} .

If y is on some Q_i , $1 \leq i \leq n$, then the (u, y) -section of Q_i has an arc in B . Otherwise, two paths in $\{P_1, \dots, P_n, P_{n+1}\}$ intersect at a vertex other than u, v , or x . Now if we replace the (u, y) -section of Q_i by the (u, y) -section of P_k we get n internally disjoint (u, v) -paths in $D - \{x\}$ using less arcs in B than Q_1, \dots, Q_n . This is a contradiction. ■

A similar proof can be used for the undirected version of Menger's theorem.

References

- [1] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*. North-Holland, New York (1981).
- [2] G. B. Dantzig and D. R. Fulkerson, On the max-flow min-cut theorem of networks. Linear inequalities and related systems. *Annals of Mathematics Studies* 38. Princeton University, Princeton, NJ (1956), pp. 215–221.
- [3] G. A. Dirac, Short Proof of Menger's graph theorem. *Mathematika* 13 (1966) 42–44.
- [4] F. Escalante, Schnittverbände in Graphen. *Abh. Math. Sem. Univ. Hamburg* 38 (1972) 199–220.
- [5] T. Gallai, Maximum-minimum Sätze und verallgemeinerte Faktoren von Graphen. *Acta Math. Acad. Sci. Hungar.* 12 (1961) 131–173.
- [6] B. Grünbaum, On n -connected graphs. *Math. Nachr.* 39 (1969) 345–347.
- [7] E. V. Gutnikov, Relative joinability and separability of ultragraphs. *Prikl. Mat. i Programirovanie Vyp.* 11 (1974) 29–33, 127–128.
- [8] R. Halin, Über trennende Eckenmengen in Graphen und den Mengerschen Satz. *Math. Ann.* 157 (1964) 34–41.
- [9] L. Lovász, A remark on Menger's theorem. *Acta Math. Acad. Sci. Hungar.* 21 (1970) 365–368.
- [10] K. Menger, Zur allgemeinen Kurventheorie. *Fund. Math.* 10 (1927) 96–115.
- [11] C. St. J. A. Nash-Williams and W. T. Tutte, More proofs of Menger's theorem. *J. Graph Theory* 1 (1977) 13–17.
- [12] P. V. O'Neil, A new proof of Menger's theorem, *J. Graph Theory* 2 (1978) 257–259.

- [13] J. S. Pym, A proof of Menger's theorem. *Monatsh. Math.* **73** (1969) 81–83.
- [14] H. Whitney, Non-separable and planar graphs. *Trans. Amer. Math. Soc.* **34** (1932) 339–362.