

3-11 Matchings and Factors

(Part II: Perfect Matchings)

Hengfeng Wei

hfwei@nju.edu.cn

December 24, 2018



5.10 5.34 5.22 5.26

Chinese Postman Problem (CPP)

(Postman Tour Problem, Route Inspection Problem)





管梅谷(1934-)

第10卷 第3期
1960年12月

数 学 学 报
ACTA MATHEMATICA SINICA

Vol. 10, No. 3
Dec., 1960

奇 偶 点 图 上 作 业 法*

管 梅 谷

(山东师范学院)

§ 1. 问题的提出

在邮局搞线性规划时,发现了下述问题:“一个投递员每次上班,要走遍他负责送信的段^①,然后回到邮局,问应该怎样走才能使所走的路程最短.”

《奇偶点图上作业法》, 1960

Translated into English in 1962



Jack Edmonds (1934-)

MATCHING, EULER TOURS AND THE CHINESE POSTMAN

Jack EDMONDS

University of Waterloo, Waterloo, Ontario, Canada

and

Ellis L. JOHNSON

IBM Watson Research Center, Yorktown Heights, New York, U.S.A.

Received 20 May 1972

Revised manuscript received 3 April 1973

The solution of the Chinese postman problem using matching theory is given. The convex hull of integer solutions is described as a linear programming polyhedron. This polyhedron is used to show that a good algorithm gives an optimum solution. The algorithm is a specialization of the more general b -matching blossom algorithm. Algorithms for finding Euler tours and related problems are also discussed.

“Matching, Euler Tours and
the Chinese Postman”, 1973(1965)

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find the shortest **tour** such that **each edge is traversed at least once**.

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find the shortest **tour** such that **each edge is traversed at least once**.

Q : What is the relation between Postman Tour and Eulerian Tour?



P : A postman tour of G

P : A postman tour of G

P contains every edge e at least once.

P : A postman tour of G

P contains every edge e at least once.

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

P : A postman tour of G

P contains every edge e at least once.

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

Construct $G' = G + e \cdot x_e$

P : A postman tour of G

P contains every edge e at least once.

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

Construct $G' = G + e \cdot x_e$

P is an Eulerian tour of G' .

P : A postman tour of G

P contains every edge e at least once.

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

Construct $G' = G + e \cdot x_e$

P is an Eulerian tour of G' .

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find $x_e \in \mathbb{N}$ for each edge e of G

to minimize $\sum_e w(e)x_e$,

such that $G' = G + e \cdot x_e$ is an Eulerian graph.

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find $x_e \in \mathbb{N}$ for each edge e of G

$$\text{to minimize } \sum_e w(e)x_e,$$

such that $G' = G + e \cdot x_e$ is an Eulerian graph.

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find $x_e \in \mathbb{N}$ for each edge e of G

$$\text{to minimize } \sum_e w(e)x_e,$$

such that $G' = G + e \cdot x_e$ is an Eulerian graph.

Q : What are the possible values of each x_e ?

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find $x_e \in \mathbb{N}$ for each edge e of G

$$\text{to minimize } \sum_e w(e)x_e,$$

such that $G' = G + e \cdot x_e$ is an Eulerian graph.

Q : What are the possible values of each x_e ?

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find $x_e \in \{0, 1\}$ for each edge e of G

$$\text{to minimize } \sum_e w(e)x_e,$$

such that $G' = G + e \cdot x_e$ is an Eulerian graph.

1: $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$

-
- 1: $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
 - 2: Construct a complete weighted graph G_p with vertices V_o :
 - 3: **for** $u, v \in V_o$ **do**
 - 4: $w(u, v) \leftarrow$ the length of the shortest path between u and v

-
- 1: $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
 - 2: Construct a complete weighted graph G_p with vertices V_o :
 - 3: **for** $u, v \in V_o$ **do**
 - 4: $w(u, v) \leftarrow$ the length of the shortest path between u and v
 - 5: Find a **minimum-weighted perfect matching** M of G_p

-
- 1: $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
 - 2: Construct a complete weighted graph G_p with vertices V_o :
 - 3: **for** $u, v \in V_o$ **do**
 - 4: $w(u, v) \leftarrow$ the length of the shortest path between u and v
 - 5: Find a **minimum-weighted perfect matching** M of G_p
 - 6: **for** $(u, v) \in M$ **do**
 - 7: $p \leftarrow$ the shortest path between u and v
 - 8: $\forall e \in p : x_e \leftarrow 1$
-

-
- 1: $V_o \leftarrow \{v \in V(G) : \deg(v) \text{ is odd}\}$
 - 2: Construct a complete weighted graph G_p with vertices V_o :
 - 3: **for** $u, v \in V_o$ **do**
 - 4: $w(u, v) \leftarrow$ the length of the shortest path between u and v
 - 5: Find a **minimum-weighted perfect matching** M of G_p
 - 6: **for** $(u, v) \in M$ **do**
 - 7: $p \leftarrow$ the shortest path between u and v
 - 8: $\forall e \in p : x_e \leftarrow 1$
-

Q : What if some edge $e \in E(G)$ is in two shortest paths corresponding to (two) matching edges of G_p ?

Theorem (Edge-disjointness of Shortest Paths)

No edge $e \in E(G)$ is in two shortest paths corresponding to (two) matching edges of G_p .

Theorem (Edge-disjointness of Shortest Paths)

No edge $e \in E(G)$ is in two shortest paths corresponding to (two) matching edges of G_p .

Proof.

Theorem (Edge-disjointness of Shortest Paths)

No edge $e \in E(G)$ is in two shortest paths corresponding to (two) matching edges of G_p .

Proof.

By Contradiction.

Theorem (Edge-disjointness of Shortest Paths)


No edge $e \in E(G)$ is in two shortest paths corresponding to (two) matching edges of G_p .

Proof.

By Contradiction.

Suppose that

$$\exists e \in E(G) : e \in u_1 \rightsquigarrow u_2 \wedge e \in v_1 \rightsquigarrow v_2$$




Theorem (Edge-disjointness of Shortest Paths)

No edge $e \in E(G)$ is in two shortest paths corresponding to (two) matching edges of G_p .

Proof.

By Contradiction.



Suppose that

$$\exists e \in E(G) : e \in u_1 \rightsquigarrow u_2 \wedge e \in v_1 \rightsquigarrow v_2$$

Contradiction:

$$u_1 \rightsquigarrow v_1, u_2 \rightsquigarrow v_2 \implies \text{smaller perfect matching}$$




Theorem (Property of CHINESE-POSTMAN)

*The edges with $x_e = 1$ obtained by CHINESE-POSTMAN is a **minimum** collection of **edge-disjoint paths** connecting pairs of odd vertices.*

Theorem (Property of CHINESE-POSTMAN)

*The edges with $x_e = 1$ obtained by CHINESE-POSTMAN is a **minimum** collection of **edge-disjoint paths** connecting pairs of odd vertices.*



Theorem (Property of CHINESE-POSTMAN)

*The edges with $x_e = 1$ obtained by CHINESE-POSTMAN is a **minimum** collection of **edge-disjoint paths** connecting pairs of odd vertices.*

To prove that these $x_e = 1$ obtained by CHINESE-POSTMAN satisfies:

Definition (Chinese Postman Problem)

Given an undirected weighted graph G with $w(e) > 0$,
to find $x_e \in \{0, 1\}$ for each edge e of G

$$\text{to minimize } \sum_e w(e)x_e,$$

such that $G' = G + e \cdot x_e$ is an Eulerian graph.

Lemma (CHINESE-POSTMAN Gives a Postman Tour)

$G' = G + e \cdot x_e$ is an Eulerian graph.

Lemma (CHINESE-POSTMAN Gives a Postman Tour)

$G' = G + e \cdot x_e$ is an Eulerian graph.

Proof.

A collection of edge-disjoint paths connecting pairs of odd vertices. □

Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$$\sum_e w(e)x_e \text{ is minimized.}$$

Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$$\sum_e w(e)x_e \text{ is minimized.}$$

Theorem (Property of CHINESE-POSTMAN)

*The edges with $x_e = 1$ obtained by CHINESE-POSTMAN is a **minimum** collection of **edge-disjoint simple paths connecting pairs of odd vertices**.*

Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$$\sum_e w(e)x_e \text{ is minimized.}$$

Theorem (Property of CHINESE-POSTMAN)

*The edges with $x_e = 1$ obtained by CHINESE-POSTMAN is a **minimum** collection of **edge-disjoint simple paths connecting pairs of odd vertices**.*

P: An optimal postman tour of G

Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$$\sum_e w(e)x_e \text{ is minimized.}$$

Theorem (Property of CHINESE-POSTMAN)

*The edges with $x_e = 1$ obtained by CHINESE-POSTMAN is a **minimum** collection of **edge-disjoint simple paths connecting pairs of odd vertices**.*

P: An optimal postman tour of G

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

Lemma (CHINESE-POSTMAN Gives an Optimal Postman Tour)

$$\sum_e w(e)x_e \text{ is minimized.}$$

Theorem (Property of CHINESE-POSTMAN)

*The edges with $x_e = 1$ obtained by CHINESE-POSTMAN is a **minimum collection of edge-disjoint simple paths connecting pairs of odd vertices.***

P: An optimal postman tour of G

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

We show that the edges with $x_e = 1$ is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

Lemma (Property of Optimal Postman Tours)

P : An optimal postman tour of G

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

The edges with $x_e = 1$ is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

Lemma (Property of Optimal Postman Tours)

P : An optimal postman tour of G

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

The edges with $x_e = 1$ is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

Proof.

By Construction.

Lemma (Property of Optimal Postman Tours)

P : An optimal postman tour of G

Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

The edges with $x_e = 1$ is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

Proof.

By Construction.

An odd number of edges e with $x_e = 1$ meet odd nodes.

An even number of edges e with $x_e = 1$ meet even nodes.

Lemma (Property of Optimal Postman Tours)

P: An optimal postman tour of G


Let $1 + x_e$ ($x_e \in \mathbb{N}$) be the number of times edge e is in P .

The edges with $x_e = 1$ is a collection of edge-disjoint simple paths connecting pairs of odd vertices.

Proof.

By Construction.

An odd number of edges e with $x_e = 1$ meet odd nodes.
An even number of edges e with $x_e = 1$ meet even nodes.







Office 302

Mailbox: H016

hfwei@nju.edu.cn