

# 3-10 Traversability


## (Part II: Hamiltonian Graphs)

Hengfeng Wei

hfwei@nju.edu.cn

December 10, 2018





CZ 6.20

### Theorem (Necessary Condition; Theorem 6.5)

*If  $G$  is a Hamiltonian graph, then for each nonempty set  $S \subset V(G)$ ,*

$$k(G - S) \leq |S|.$$

### Theorem (Necessary Condition; Theorem 6.5)

*If  $G$  is a Hamiltonian graph, then for each nonempty set  $S \subset V(G)$ ,*

$$k(G - S) \leq |S|.$$

$\implies$  Hamiltonian graphs are 2-connected. (Why?)

### Theorem (Necessary Condition; Theorem 6.5)

*If  $G$  is a Hamiltonian graph, then for each nonempty set  $S \subset V(G)$ ,*

$$k(G - S) \leq |S|.$$

$\implies$  Hamiltonian graphs are 2-connected. (Why?)

Hamiltonian graphs has “good” connectedness.

### Theorem (Necessary Condition; Theorem 6.5)

*If  $G$  is a Hamiltonian graph, then for each nonempty set  $S \subset V(G)$ ,*

$$k(G - S) \leq |S|.$$

$\implies$  Hamiltonian graphs are 2-connected. (Why?)

Hamiltonian graphs has “good” connectedness.

Graphs with “bad” connectedness are **not** Hamiltonian.

## Theorem (Ore's Theorem, 1960; Theorem 6.6)

Let  $G$  be a graph of order  $n \geq 3$ . If

$$\deg(u) + \deg(v) \geq n$$

for each pair  $u, v$  of nonadjacent vertices of  $G$ , then  $G$  is Hamiltonian.

Theorem (Ore's Theorem, 1960; Theorem 6.6)

Let  $G$  be a graph of order  $n \geq 3$ . If

$$\deg(u) + \deg(v) \geq n$$

for each pair  $u, v$  of nonadjacent vertices of  $G$ , then  $G$  is Hamiltonian.

Proof.

By Contradiction and Extremality.



Theorem (Ore's Theorem, 1960; Theorem 6.6)

Let  $G$  be a graph of order  $n \geq 3$ . If

$$\deg(u) + \deg(v) \geq n$$

for each pair  $u, v$  of nonadjacent vertices of  $G$ , then  $G$  is Hamiltonian.

Proof.

By Contradiction and Extremality.

By Contradiction:

$\exists G$  satisfying Ore's Condition, but  $G$  is not Hamiltonian.

Theorem (Ore's Theorem, 1960; Theorem 6.6)

Let  $G$  be a graph of order  $n \geq 3$ . If

$$\deg(u) + \deg(v) \geq n$$

for each pair  $u, v$  of nonadjacent vertices of  $G$ , then  $G$  is Hamiltonian.

Proof.

By Contradiction and Extremality.

By Contradiction:

$\exists G$  satisfying Ore's Condition, but  $G$  is not Hamiltonian.

By Extremality:

Consider a critical  $G$ :  $G$  is not Hamiltonian but  $G + uv$  is Hamiltonian.

Theorem (Ore's Theorem, 1960; Theorem 6.6)

Let  $G$  be a graph of order  $n \geq 3$ . If

$$\deg(u) + \deg(v) \geq n$$

for each pair  $u, v$  of nonadjacent vertices of  $G$ , then  $G$  is Hamiltonian.

Proof.

By Contradiction and Extremality.

By Contradiction:

$\exists G$  satisfying Ore's Condition, but  $G$  is not Hamiltonian.

By Extremality:

Consider a critical  $G$ :  $G$  is not Hamiltonian but  $G + uv$  is Hamiltonian.

Contradiction: This critical  $G$  is actually Hamiltonian.



Theorem (Dirac's Theorem, 1952; Corollary 6.7)

*Let  $G$  be a graph of order  $n \geq 3$ . If*

$$\forall v \in V(G) : \deg(v) \geq n/2,$$

*then  $G$  is Hamiltonian.*

Theorem (Ore's Theorem, 1960; Theorem 6.8)

*Let  $u$  and  $v$  be nonadjacent vertices in a graph  $G$  of order  $n$  such that*

$$\deg(u) + \deg(v) \geq n.$$

*Then  $G + uv$  is Hamiltonian  $\iff G$  is Hamiltonian.*

### Theorem (Ore's Theorem, 1960; Theorem 6.8)

*Let  $u$  and  $v$  be nonadjacent vertices in a graph  $G$  of order  $n$  such that*

$$\deg(u) + \deg(v) \geq n.$$

*Then  $G + uv$  is Hamiltonian  $\iff G$  is Hamiltonian.*

### Definition (Closure $C(G)$ )

The closure  $C(G)$  of a graph  $G$  is the graph obtained from  $G$  by iteratively adding edges joining pairs of nonadjacent vertices  $u$  and  $v$  such that  $\deg(u) + \deg(v) \geq n$ , until no such pair remains.

Theorem (Bondy-Chavátal Theorem, 1976; Theorem 6.9)

$G$  is Hamiltonian  $\iff C(G)$  is Hamiltonian.

Theorem (Bondy-Chavátal Theorem, 1976; Theorem 6.9)

*$G$  is Hamiltonian  $\iff C(G)$  is Hamiltonian.*



Theorem (Bondy-Chavátal Theorem, 1976; Theorem 6.9)

*$G$  is Hamiltonian  $\iff C(G)$  is Hamiltonian.*

Corollary (Corollary 6.10)

*If  $G$  is a graph of order  $n \geq 3$  such that  $C(G) = K_n$ , then  $G$  is Hamiltonian.*

Theorem (Bondy-Chavátal Theorem, 1976; Theorem 6.9)

*$G$  is Hamiltonian  $\iff C(G)$  is Hamiltonian.*

Corollary (Corollary 6.10)

*If  $G$  is a graph of order  $n \geq 3$  such that  $C(G) = K_n$ , then  $G$  is Hamiltonian.*

Theorem (Lajos Pósa)

*Let  $G$  be a graph of order  $n \geq 3$ . If for each integer  $j$  with  $1 \leq j \leq \frac{n}{2}$ , the number of vertices of  $G$  with degree at most  $j$  is less than  $j$ , then  $G$  is Hamiltonian.*

## Theorem (Well-definedness of $C(G)$ )

$C(G)$  is well-defined.

## Theorem (Well-definedness of $C(G)$ )

$C(G)$  is well-defined.

$C(G)$  does not depend on the order in which we choose to add edges.

## Theorem (Well-definedness of $C(G)$ )

$C(G)$  is well-defined.

$C(G)$  does not depend on the order in which we choose to add edges.

$$\left( G + \langle e_1, \dots, e_r \rangle = G_1 \right) \wedge \left( G + \langle f_1, \dots, f_s \rangle = G_2 \right) \implies G_1 = G_2$$

## Theorem (Well-definedness of $C(G)$ )

$C(G)$  is well-defined.

$C(G)$  does not depend on the order in which we choose to add edges.

$$\left( G + \langle e_1, \dots, e_r \rangle = G_1 \right) \wedge \left( G + \langle f_1, \dots, f_s \rangle = G_2 \right) \implies G_1 = G_2$$

$$G_1 \subseteq G_2 \wedge G_2 \subseteq G_1$$

## Theorem (Well-definedness of $C(G)$ )

$C(G)$  is well-defined.

$C(G)$  does not depend on the order in which we choose to add edges.

$$\left( G + \langle e_1, \dots, e_r \rangle = G_1 \right) \wedge \left( G + \langle f_1, \dots, f_s \rangle = G_2 \right) \implies G_1 = G_2$$

$$G_1 \subseteq G_2 \wedge G_2 \subseteq G_1$$

$$\forall e_i \in G_1 : e_i \in G_2$$

## Theorem (Well-definedness of $C(G)$ )

$C(G)$  is well-defined.

$C(G)$  does not depend on the order in which we choose to add edges.

$$\left( G + \langle e_1, \dots, e_r \rangle = G_1 \right) \wedge \left( G + \langle f_1, \dots, f_s \rangle = G_2 \right) \implies G_1 = G_2$$

$$G_1 \subseteq G_2 \wedge G_2 \subseteq G_1$$

$$\forall e_i \in G_1 : e_i \in G_2$$

By induction on the order  $e_i$  is added to  $G_1$ .



## Hamiltonian Graphs and 2-Connectedness (Problem 6.20)

Let  $G$  be a graph of order  $n \geq 3$  having the property that for each  $v \in V(G)$ , there is a Hamiltonian path with initial vertex  $v$ .

Show that  $G$  is 2-connected but not necessarily Hamiltonian.

## Hamiltonian Graphs and 2-Connectedness (Problem 6.20)

Let  $G$  be a graph of order  $n \geq 3$  having the property that for each  $v \in V(G)$ , there is a Hamiltonian path with initial vertex  $v$ . Show that  $G$  is 2-connected but not necessarily Hamiltonian.

2-connected: Connected + No cut-vertex

## Hamiltonian Graphs and 2-Connectedness (Problem 6.20)

Let  $G$  be a graph of order  $n \geq 3$  having the property that for each  $v \in V(G)$ , there is a Hamiltonian path with initial vertex  $v$ . Show that  $G$  is 2-connected but not necessarily Hamiltonian.

2-connected: Connected + No cut-vertex


Suppose, by contradiction,  $v$  is a cut-vertex of  $G$ .

## Hamiltonian Graphs and 2-Connectedness (Problem 6.20)

Let  $G$  be a graph of order  $n \geq 3$  having the property that for each  $v \in V(G)$ , there is a Hamiltonian path with initial vertex  $v$ . Show that  $G$  is 2-connected but not necessarily Hamiltonian.

2-connected: Connected + No cut-vertex

Suppose, by contradiction,  $v$  is a cut-vertex of  $G$ .




## Hamiltonian Graphs and 2-Connectedness (Problem 6.20)

Let  $G$  be a graph of order  $n \geq 3$  having the property that for each  $v \in V(G)$ , there is a Hamiltonian path with initial vertex  $v$ . Show that  $G$  is 2-connected but not necessarily Hamiltonian.

2-connected: Connected + No cut-vertex

Suppose, by contradiction,  $v$  is a cut-vertex of  $G$ .




**Contradiction:** No Hamiltonian path with initial vertex  $v$ .

## Hamiltonian Graphs and 2-Connectedness (Problem 6.20)

Let  $G$  be a graph of order  $n \geq 3$  having the property that for each  $v \in V(G)$ , there is a Hamiltonian path with initial vertex  $v$ . Show that  $G$  is 2-connected but **not necessarily Hamiltonian**.

## Hamiltonian Graphs and 2-Connectedness (Problem 6.20)

Let  $G$  be a graph of order  $n \geq 3$  having the property that for each  $v \in V(G)$ , there is a Hamiltonian path with initial vertex  $v$ . Show that  $G$  is 2-connected but **not necessarily Hamiltonian**.









Office 302

Mailbox: H016

hfwei@nju.edu.cn