

# Law of total probability

In probability theory, the **law** (or **formula**) of **total probability** is a fundamental rule relating marginal probabilities to conditional probabilities. It expresses the total probability of an outcome which can be realized via several distinct events—hence the name.

## Contents

**Statement**

**Informal formulation**

**Example**

**Other names**

**See also**

**Notes**

**References**

## Statement

The law of total probability is<sup>[1]</sup> the proposition that if  $\{B_n : n = 1, 2, 3, \dots\}$  is a finite or countably infinite partition of a sample space (in other words, a set of pairwise disjoint events whose union is the entire sample space) and each event  $B_n$  is measurable, then for any event  $A$  of the same probability space:

$$\Pr(A) = \sum_n \Pr(A \cap B_n)$$

or, alternatively,<sup>[1]</sup>

$$\Pr(A) = \sum_n \Pr(A \mid B_n) \Pr(B_n),$$

where, for any  $n$  for which  $\Pr(B_n) = 0$  these terms are simply omitted from the summation, because  $\Pr(A \mid B_n)$  is finite.

The summation can be interpreted as a weighted average, and consequently the marginal probability,  $\Pr(A)$ , is sometimes called "average probability";<sup>[2]</sup> "overall probability" is sometimes used in less formal writings.<sup>[3]</sup>

The law of total probability can also be stated for conditional probabilities. Taking the  $B_n$  as above, and assuming  $C$  is an event independent with any of the  $B_n$ :

$$\Pr(A \mid C) = \sum_n \Pr(A \mid C \cap B_n) \Pr(B_n \mid C) = \sum_n \Pr(A \mid C \cap B_n) \Pr(B_n)$$

## Informal formulation

The above mathematical statement might be interpreted as follows: *given an outcome  $A$ , with known conditional probabilities given any of the  $B_n$  events, each with a known probability itself, what is the total probability that  $A$  will happen?* The answer to this question is given by  $\Pr(A)$ .

## Example

---

Suppose that two factories supply light bulbs to the market. Factory  $X$ 's bulbs work for over 5000 hours in 99% of cases, whereas factory  $Y$ 's bulbs work for over 5000 hours in 95% of cases. It is known that factory  $X$  supplies 60% of the total bulbs available and  $Y$  supplies 40% of the total bulbs available. What is the chance that a purchased bulb will work for longer than 5000 hours?

Applying the law of total probability, we have:

$$\begin{aligned}\Pr(A) &= \Pr(A \mid B_X) \cdot \Pr(B_X) + \Pr(A \mid B_Y) \cdot \Pr(B_Y) \\ &= \frac{99}{100} \cdot \frac{6}{10} + \frac{95}{100} \cdot \frac{4}{10} = \frac{594 + 380}{1000} = \frac{974}{1000}\end{aligned}$$

where

- $\Pr(B_X) = \frac{6}{10}$  is the probability that the purchased bulb was manufactured by factory  $X$ ;
- $\Pr(B_Y) = \frac{4}{10}$  is the probability that the purchased bulb was manufactured by factory  $Y$ ;
- $\Pr(A \mid B_X) = \frac{99}{100}$  is the probability that a bulb manufactured by  $X$  will work for over 5000 hours;
- $\Pr(A \mid B_Y) = \frac{95}{100}$  is the probability that a bulb manufactured by  $Y$  will work for over 5000 hours.

Thus each purchased light bulb has a 97.4% chance to work for more than 5000 hours.

## Other names

---

The term ***law of total probability*** is sometimes taken to mean the **law of alternatives**, which is a special case of the law of total probability applying to discrete random variables. One author even uses the terminology "continuous law of alternatives" in the continuous case.<sup>[4]</sup> This result is given by Grimmett and Welsh<sup>[5]</sup> as the **partition theorem**, a name that they also give to the related law of total expectation.

## See also

---

- Law of total expectation
- Law of total variance
- Law of total cumulance
- Marginal distribution

## Notes

---

1. Zwillinger, D., Kokoska, S. (2000) *CRC Standard Probability and Statistics Tables and Formulae*, CRC Press. ISBN 1-58488-059-7 page 31.
2. Paul E. Pfeiffer (1978). *Concepts of probability theory* ([https://books.google.com/books?id=\\_mayRBczVRwC&pg=PA47](https://books.google.com/books?id=_mayRBczVRwC&pg=PA47)). Courier Dover Publications. pp. 47–48. ISBN 978-0-486-63677-1.
3. Deborah Rumsey (2006). *Probability for dummies* (<https://books.google.com/books?id=Vj3NZ59ZcnoC&pg=PA58>). For Dummies. p. 58. ISBN 978-0-471-75141-0.
4. Kenneth Baclawski (2008). *Introduction to probability with R* (<https://books.google.com/books?id=KgIc9g5IPf4C&pg=PA179>). CRC Press. p. 179. ISBN 978-1-4200-6521-3.
5. *Probability: An Introduction*, by Geoffrey Grimmett and Dominic Welsh, Oxford Science Publications, 1986, Theorem 1B.

## References

---

- *Introduction to Probability and Statistics* by William Mendenhall, Robert J. Beaver, Barbara M. Beaver, Thomson Brooks/Cole, 2005, page 159.
  - *Theory of Statistics*, by Mark J. Schervish, Springer, 1995.
  - *Schaum's Outline of Probability, Second Edition*, by John J. Schiller, Seymour Lipschutz, McGraw–Hill Professional, 2010, page 89.
  - *A First Course in Stochastic Models*, by H. C. Tijms, John Wiley and Sons, 2003, pages 431–432.
  - *An Intermediate Course in Probability*, by Alan Gut, Springer, 1995, pages 5–6.
- 

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Law\\_of\\_total\\_probability&oldid=833049144](https://en.wikipedia.org/w/index.php?title=Law_of_total_probability&oldid=833049144)"

---

**This page was last edited on 2018-03-29, at 19:45:37.**

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.