3-5 Minimum Spanning Trees

Hengfeng Wei

hfwei@nju.edu.cn

October 22, 2018



 $\forall v \in V(G) : \deg(v) \ge 2 \implies G \text{ contains a cycle}$

$$\forall v \in V(G) : \deg(v) \ge 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$


$$m = n - k(G) \le n - 1$$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$m = n - k(G) \le n - 1$$

$$\sum_{v \in V(G)} \deg(v) \le 2(n-1)$$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$



maximal path $P_{u,v}$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \ge 2$$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \ge 2$$

$$\implies \nexists v \in V(G): \deg(v) = 1$$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \ge 2$$

$$\implies \nexists v \in V(G) : \deg(v) = 1$$

$$\xrightarrow{\text{Theorem 4.3}} G \text{ is not a tree}$$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \ge 2$$

$$\implies \nexists v \in V(G) : \deg(v) = 1$$

 $\xrightarrow{\text{Theorem 4.3}} G$ is not a tree

 $\xrightarrow{\text{Theorem } 4.2} \exists u, v \in V(G) : u, v \text{ are connected by } \geq 2 \text{ paths}$

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G): \deg(v) \geq 2$$

$$\implies \nexists v \in V(G) : \deg(v) = 1$$

 $\xrightarrow{\text{Theorem 4.3}} G$ is not a tree

 $\xrightarrow{\text{Theorem } 4.2} \exists u, v \in V(G) : u, v \text{ are connected by } \geq 2 \text{ paths}$

 $\implies G$ contains a cycle

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \ge 2$$

$$\implies \nexists v \in V(G) : \deg(v) = 1$$

 $\xrightarrow{\text{Theorem 4.3}} G$ is not a tree

 $\xrightarrow{\text{Theorem } 4.2} \exists u, v \in V(G) : u, v \text{ are connected by } \geq 2 \text{ paths}$

 $\implies G$ contains a cycle

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \ge 2$$

$$\implies \nexists v \in V(G) : \deg(v) = 1$$

 $\xrightarrow{\text{Theorem 4.3}} G$ is not a tree

Consider each component G' of G

 $\xrightarrow{\text{Theorem } 4.2} \exists u, v \in V(G') : u, v \text{ are connected by } \geq 2 \text{ paths}$

 $\implies G'$ contains a cycle

$$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$$

$$\forall v \in V(G) : \deg(v) \ge 2$$

$$\implies \nexists v \in V(G) : \deg(v) = 1$$

 $\xrightarrow{\text{Theorem 4.3}} G$ is not a tree

Consider each component G' of G

 $\xrightarrow{\text{Theorem } 4.2} \exists u, v \in V(G') : u, v \text{ are connected by } \geq 2 \text{ paths}$

- $\implies G'$ contains a cycle
- \implies G contains a cycle



G: a connected graph

 $e \in E(G)$ is a bridge $\iff e \in \forall ST \text{ of } G$

G: a connected graph

 $e \in E(G)$ is a bridge $\iff e \in \forall \mathrm{ST}$ of G



G: a connected graph

 $e \in E(G)$ is a bridge $\iff e \in \forall ST \text{ of } G$

G: a connected graph

 $e \in E(G)$ is a bridge $\iff e \in \forall ST \text{ of } G$


ST of
$$G - e$$

G: a connected graph

 $e \in E(G)$ is a bridge $\iff e \in \forall ST \text{ of } G$

By Contradiction.

ST of G - e



Cut Property

Cut Property (Version I)

X: A part of some MST T of G

 $(S, V \setminus S)$: A cut such that X does not cross $(S, V \setminus S)$

e: A lightest edge across $(S, V \setminus S)$

Cut Property (Version I)

X: A part of some MST T of G

 $(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$

e: A lightest edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of some MST T' of G.

Cut Property (Version I)


X: A part of some MST T of G


 $(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$

e: A lightest edge across $(S, V \setminus S)$


Then $X \cup \{e\}$ is a part of some MST T' of G.

Correctness of Prim's and Kruskal's algorithms.





$$T' = \underbrace{T + \{e\}}_{\text{if } e \notin T} - \{e'\}$$



$$T' = \underbrace{T + \{e\}}_{\text{if } e \notin T} - \{e'\}$$


$$\text{"a"} \to \text{"the"} \implies \text{"some"} \to \text{"all"}$$

Cut Property (Version II)

A cut $(S, V \setminus S)$

Let e = (u, v) be **a** lightest edge across $(S, V \setminus S)$

 \exists MST T of $G: e \in T$




Cut Property (Version II)

A cut
$$(S, V \setminus S)$$

Let e = (u, v) be **a** lightest edge across $(S, V \setminus S)$

\exists MST T of $G: e \in T$




$$T' = \underbrace{T + \{e\}}_{\text{if } e \notin T} - \{e'\}$$

Cut Property (Version II)

A cut
$$(S, V \setminus S)$$

Let e = (u, v) be **a** lightest edge across $(S, V \setminus S)$

\exists MST T of $G: e \in T$



$$T' = \underbrace{T + \{e\}}_{\text{if } e \notin T} - \{e'\}$$

"a" \rightarrow "the" \Longrightarrow " \exists " \rightarrow " \forall "



Application of Cut Property

$$e = (u, v) \in G$$
 is a lightest edge $\implies e \in \exists$ MST of G

Application of Cut Property

 $e = (u, v) \in G$ is the unique lightest edge $\implies e \in \forall MST$

Application of Cut Property

$$e = (u,v) \in G$$
 is a lightest edge $\implies e \in \exists$ MST of G

$$\left(S = \{u\}, V \setminus S\right)$$

Application of Cut Property

 $e = (u, v) \in G$ is the unique lightest edge $\implies e \in \forall$ MST

Wrong Divide&Conquer Algorithm for MST


$$(V_1, V_2) : ||V_1| - |V_2|| \le 1$$

 $T_1 + T_2 + \{e\} : e$ is a lightest edge across (V_1, V_2)

Wrong Divide&Conquer Algorithm for MST

$$(V_1, V_2) : ||V_1| - |V_2|| \le 1$$

 $T_1 + T_2 + \{e\}$: e is a lightest edge across (V_1, V_2)




Cycle Property

Cycle Property

- \blacktriangleright Let C be any cycle in G
- ▶ Let e = (u, v) be **a** maximum-weight edge in C


Then \exists MST T of $G: e \notin T$.



Cycle Property

- \blacktriangleright Let C be any cycle in G
- ▶ Let e = (u, v) be **a** maximum-weight edge in C

Then \exists MST T of $G: e \notin T$.




$$T' = \underbrace{T - \{e\}}_{\text{if } e \in T} + \{e'\}$$



Cycle Property

- \blacktriangleright Let C be any cycle in G
- ▶ Let e = (u, v) be a maximum-weight edge in C

Then \exists MST T of $G: e \notin T$.



$$T' = \underbrace{T - \{e\}}_{\text{if } e \in T} + \{e'\}$$

"a" \rightarrow "the" \Longrightarrow " \exists " \rightarrow " \forall "



Reverse-delete algorithm (wiki; clickable)

Reverse-delete algorithm (wiki; clickable)

$$O(m \log n ((\log \log n)^3))$$

Reverse-delete algorithm (wiki; clickable)

$$O(m \log n ((\log \log n)^3))$$

Proof.

Cycle Property

Reverse-delete algorithm (wiki; clickable)

$$O(m \log n ((\log \log n)^3))$$

Proof.

Cycle Property

$$T \subseteq F \implies \exists \ T' : T' \subseteq F - \{e\}$$



Reverse-delete algorithm (wiki; clickable)

$$O(m \log n ((\log \log n)^3))$$

Proof.

Cycle Property

$$T \subseteq F \implies \exists T' : T' \subseteq F - \{e\}$$

"On the Shortest Spanning Subtree of a Graph and the Traveling Salesman Problem"

— Kruskal, 1956.

$$G = (V, E), \quad |E| > |V| - 1$$

 \boldsymbol{e} : the unique maximum-weighted edge of G



 $e \notin \text{ any MST}$

$$G = (V, E), \quad |E| > |V| - 1$$

e : the unique maximum-weighted edge of G

$$\Longrightarrow$$

 $e \notin \text{any MST}$

Bridge



$$C \subseteq G$$
, $e \in C$

e: the unique maximum-weighted edge of G



 $e \notin \text{any MST}$

$$C \subseteq G$$
, $e \in C$

e: the unique maximum-weighted edge of G

$$\Longrightarrow$$

 $e \notin \text{any MST}$

Cycle Property

$$C \subseteq G, e \in C$$

e: the unique lightest edge of C

$$\implies$$


 $e \in \forall \ \mathrm{MST}$

$$C \subseteq G, e \in C$$

e: the unique lightest edge of C



 $e \in \forall MST$



Uniqueness of MST

Distinct weights \implies Unique MST.

Distinct weights \implies Unique MST.

Distinct weights \implies Unique MST.

By Contradiction.

 \exists MSTs $T_1 \neq T_2$

Distinct weights \implies Unique MST.

$$\exists$$
 MSTs $T_1 \neq T_2$

$$\Delta E = \{ e \mid e \in T_1 \setminus T_2 \lor e \in T_2 \setminus T_1 \}$$

Distinct weights \implies Unique MST.

$$\exists$$
 MSTs $T_1 \neq T_2$

$$\Delta E = \{ e \mid e \in T_1 \setminus T_2 \lor e \in T_2 \setminus T_1 \}$$

$$e = \min \Delta E$$

Distinct weights \implies Unique MST.


$$\exists MSTs T_1 \neq T_2$$


$$\Delta E = \{ e \mid e \in T_1 \setminus T_2 \lor e \in T_2 \setminus T_1 \}$$

$$e = \min \Delta E$$


$$e \in T_1 \setminus T_2 \ (w.l.o.g)$$








$$T_2 + \{e\} \implies C$$




$$T_2 + \{e\} \implies C$$

$$\exists (e' \in C) \notin T_1$$




$$T_2 + \{e\} \implies C$$

$$\exists (e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E$$



$$T_2 + \{e\} \implies C$$

$$\exists (e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E \implies w(e') > w(e)$$



$$T_2 + \{e\} \implies C$$

$$\exists (e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E \implies w(e') > w(e)$$

$$T' = T_2 + \{e\} - \{e'\} \implies w(T') < w(T_2)$$




Condition for Uniqueness of MST $\,$

Unique MST \implies Distinct weights.

Condition for Uniqueness of MST

Unique MST \implies Distinct weights.




Unique MST (Problem 4.30)

Unique MST \implies Maximum-weight edge in any cycle is unique.


Unique MST (Problem 4.30)

Unique MST \implies Maximum-weight edge in any cycle is unique.



Unique MST (Problem 4.30)

Unique MST \implies Maximum-weight edge in any cycle is unique.




Theorem (After-class Exercise)


Maximum-weight edge in any cycle is unique \implies Unique MST.

Unique MST \implies Minimum-weight edge across any cut is unique.

Unique MST \implies Minimum-weight edge across any cut is unique.



Unique MST \implies Minimum-weight edge across any cut is unique.



Theorem (After-class Exercise)

Minimum-weight edge across any cut is unique \implies Unique MST.


To decide whether a graph has a unique MST.

To decide whether a graph has a unique MST.

Ties in Prim's and Kruskal's algorithms


To decide whether a graph has a unique MST.

Ties in Prim's and Kruskal's algorithms



To decide whether a graph has a unique MST.


Ties in Prim's and Kruskal's algorithms



$$\underbrace{\frac{T}{\text{Any MST}}}_{\text{Cycle}} + \underbrace{\underbrace{\{e\}, \forall e \notin T}_{\text{Cycle}}}$$

To decide whether a graph has a unique MST.

Ties in Prim's and Kruskal's algorithms



$$\underbrace{\frac{T}{\text{Any MST}}}_{\text{Cycle}} + \underbrace{\{e\}, \forall e \notin T}_{\text{Cycle}}$$

By Kruskal Algorithm.







Office 302

Mailbox: H016

hfwei@nju.edu.cn