


# Symmetric group

In abstract algebra, the **symmetric group** defined over any set is the group whose elements are all the bijections from the set to itself, and whose group operation is the composition of functions. In particular, the finite symmetric group  $S_n$  defined over a finite set of  $n$  symbols consists of the permutation operations that can be performed on the  $n$  symbols.<sup>[1]</sup> Since there are  $n!$  ( $n$  factorial) possible permutation operations that can be performed on a tuple composed of  $n$  symbols, it follows that the number of elements (the order) of the symmetric group  $S_n$  is  $n!$ .

Although symmetric groups can be defined on infinite sets, this article focuses on the finite symmetric groups: their applications, their elements, their conjugacy classes, a finite presentation, their subgroups, their automorphism groups, and their representation theory. For the remainder of this article, "symmetric group" will mean a symmetric group on a finite set.


The symmetric group is important to diverse areas of mathematics such as Galois theory, invariant theory, the representation theory of Lie groups, and combinatorics. Cayley's theorem states that every group  $G$  is isomorphic to a subgroup of the symmetric group on  $G$ .



A Cayley graph of the symmetric group  $S_4$


Cayley table of the symmetric group  $S_3$   
(multiplication table of permutation matrices)

These are the positions of the six matrices:



Only the unity matrices are arranged symmetrically to the main diagonal - thus the symmetric group is not abelian.

## Contents

### Definition and first properties

### Applications

### Elements

- Multiplication
- Verification of group axioms
- Transpositions
- Cycles
- Special elements

### Conjugacy classes

### Low degree groups

- Maps between symmetric groups

### Properties

### Relation with alternating group

### Generators and relations

### Subgroup structure

- Normal subgroups
- Maximal subgroups
- Sylow subgroups

Transitive subgroups

**Automorphism group**

**Homology**

**Representation theory**

**See also**

**References**

**External links**

## Definition and first properties

---

The symmetric group on a finite set  $X$  is the group whose elements are all bijective functions from  $X$  to  $X$  and whose group operation is that of function composition.<sup>[1]</sup> For finite sets, "permutations" and "bijective functions" refer to the same operation, namely rearrangement. The symmetric group of **degree**  $n$  is the symmetric group on the set  $X = \{ 1, 2, \dots, n \}$ .

The symmetric group on a set  $X$  is denoted in various ways including  $S_X$ ,  $\mathfrak{S}_X$ ,  $\Sigma_X$ ,  $X!$  and  $\text{Sym}(X)$ .<sup>[1]</sup> If  $X$  is the set  $\{ 1, 2, \dots, n \}$ , then the symmetric group on  $X$  is also denoted  $S_n$ ,<sup>[1]</sup>  $\mathfrak{S}_n$ ,  $\Sigma_n$ , and  $\text{Sym}(n)$ .

Symmetric groups on infinite sets behave quite differently from symmetric groups on finite sets, and are discussed in (Scott 1987, Ch. 11), (Dixon & Mortimer 1996, Ch. 8), and (Cameron 1999).

The symmetric group on a set of  $n$  elements has order  $n!$  (the factorial of  $n$ ).<sup>[2]</sup> It is abelian if and only if  $n$  is less than or equal to 2.<sup>[3]</sup> For  $n = 0$  and  $n = 1$  (the empty set and the singleton set), the symmetric group is trivial (it has order  $0! = 1! = 1$ ). The group  $S_n$  is solvable if and only if  $n \leq 4$ . This is an essential part of the proof of the Abel–Ruffini theorem that shows that for every  $n > 4$  there are polynomials of degree  $n$  which are not solvable by radicals, that is, the solutions cannot be expressed by performing a finite number of operations of addition, subtraction, multiplication, division and root extraction on the polynomial's coefficients.

## Applications

---

The symmetric group on a set of size  $n$  is the Galois group of the general polynomial of degree  $n$  and plays an important role in Galois theory. In invariant theory, the symmetric group acts on the variables of a multi-variate function, and the functions left invariant are the so-called symmetric functions. In the representation theory of Lie groups, the representation theory of the symmetric group plays a fundamental role through the ideas of Schur functors. In the theory of Coxeter groups, the symmetric group is the Coxeter group of type  $A_n$  and occurs as the Weyl group of the general linear group. In combinatorics, the symmetric groups, their elements (permutations), and their representations provide a rich source of problems involving Young tableaux, plactic monoids, and the Bruhat order. Subgroups of symmetric groups are called permutation groups and are widely studied because of their importance in understanding group actions, homogeneous spaces, and automorphism groups of graphs, such as the Higman–Sims group and the Higman–Sims graph.

## Elements

---

The elements of the symmetric group on a set  $X$  are the permutations of  $X$ .

## Multiplication

The group operation in a symmetric group is function composition, denoted by the symbol  $\circ$  or simply by juxtaposition of the permutations. The composition  $f \circ g$  of permutations  $f$  and  $g$ , pronounced " $f$  of  $g$ ", maps any element  $x$  of  $X$  to  $f(g(x))$ . Concretely, let (see permutation for an explanation of notation):

$$f = (1\ 3)(4\ 5) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 2 & 1 & 5 & 4 \end{pmatrix}$$

$$g = (1\ 2\ 5)(3\ 4) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 5 & 4 & 3 & 1 \end{pmatrix}.$$

Applying  $f$  after  $g$  maps 1 first to 2 and then 2 to itself; 2 to 5 and then to 4; 3 to 4 and then to 5, and so on. So composing  $f$  and  $g$  gives

$$fg = f \circ g = (1\ 2\ 4)(3\ 5) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 4 & 5 & 1 & 3 \end{pmatrix}.$$

A cycle of length  $L = k \cdot m$ , taken to the  $k$ -th power, will decompose into  $k$  cycles of length  $m$ : For example,  $(k = 2, m = 3)$ ,

$$(1\ 2\ 3\ 4\ 5\ 6)^2 = (1\ 3\ 5)(2\ 4\ 6).$$

## Verification of group axioms

To check that the symmetric group on a set  $X$  is indeed a group, it is necessary to verify the group axioms of closure, associativity, identity, and inverses.<sup>[4]</sup>

1. The operation of function composition is closed in the set of permutations of the given set  $X$ .
2. Function composition is always associative.
3. The trivial bijection that assigns each element of  $X$  to itself serves as an identity for the group.
4. Every bijection has an inverse function that undoes its action, and thus each element of a symmetric group does have an inverse which is a permutation too.

## Transpositions

A **transposition** is a permutation which exchanges two elements and keeps all others fixed; for example  $(1\ 3)$  is a transposition. Every permutation can be written as a product of transpositions; for instance, the permutation  $g$  from above can be written as  $g = (1\ 2)(2\ 5)(3\ 4)$ . Since  $g$  can be written as a product of an odd number of transpositions, it is then called an odd permutation, whereas  $f$  is an even permutation.

The representation of a permutation as a product of transpositions is not unique; however, the number of transpositions needed to represent a given permutation is either always even or always odd. There are several short proofs of the invariance of this parity of a permutation.

The product of two even permutations is even, the product of two odd permutations is even, and all other products are odd. Thus we can define the **sign** of a permutation:

$$\text{sgn } f = \begin{cases} +1, & \text{if } f \text{ is even} \\ -1, & \text{if } f \text{ is odd.} \end{cases}$$

With this definition,

$$\text{sgn}: S_n \rightarrow \{+1, -1\}$$

is a group homomorphism ( $\{+1, -1\}$  is a group under multiplication, where  $+1$  is  $e$ , the neutral element). The kernel of this homomorphism, that is, the set of all even permutations, is called the **alternating group**  $A_n$ . It is a normal subgroup of  $S_n$ , and for  $n \geq 2$  it has  $n!/2$  elements. The group  $S_n$  is the semidirect product of  $A_n$  and any subgroup generated by a single transposition.

Furthermore, every permutation can be written as a product of adjacent transpositions, that is, transpositions of the form  $(a\ a+1)$ . For instance, the permutation  $g$  from above can also be written as  $g = (4\ 5)(3\ 4)(4\ 5)(1\ 2)(2\ 3)(3\ 4)(4\ 5)$ . The sorting algorithm bubble sort is an application of this fact. The representation of a permutation as a product of adjacent transpositions is also not unique.

## Cycles

A cycle of length  $k$  is a permutation  $f$  for which there exists an element  $x$  in  $\{1, \dots, n\}$  such that  $x, f(x), f^2(x), \dots, f^k(x) = x$  are the only elements moved by  $f$ ; it is required that  $k \geq 2$  since with  $k = 1$  the element  $x$  itself would not be moved either. The permutation  $h$  defined by

$$h = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 4 & 2 & 1 & 3 & 5 \end{pmatrix}$$

is a cycle of length three, since  $h(1) = 4$ ,  $h(4) = 3$  and  $h(3) = 1$ , leaving 2 and 5 untouched. We denote such a cycle by  $(1\ 4\ 3)$ , but it could equally well be written  $(4\ 3\ 1)$  or  $(3\ 1\ 4)$  by starting at a different point. The order of a cycle is equal to its length. Cycles of length two are transpositions. Two cycles are *disjoint* if they move disjoint subsets of elements. Disjoint cycles commute: for example, in  $S_6$  there is the equality  $(4\ 1\ 3)(2\ 5\ 6) = (2\ 5\ 6)(4\ 1\ 3)$ . Every element of  $S_n$  can be written as a product of disjoint cycles; this representation is unique up to the order of the factors, and the freedom present in representing each individual cycle by choosing its starting point.

Cycles admits the following conjugation property with any permutation  $\sigma$ , this property is often used to obtain its Generators and relations.

$$\sigma(a\ b\ c\ \dots)\sigma^{-1} = (\sigma(a)\ \sigma(b)\ \sigma(c)\ \dots)$$

## Special elements

Certain elements of the symmetric group of  $\{1, 2, \dots, n\}$  are of particular interest (these can be generalized to the symmetric group of any finite totally ordered set, but not to that of an unordered set).

The **order reversing permutation** is the one given by:

$$\begin{pmatrix} 1 & 2 & \dots & n \\ n & n-1 & \dots & 1 \end{pmatrix}.$$

This is the unique maximal element with respect to the Bruhat order and the longest element in the symmetric group with respect to generating set consisting of the adjacent transpositions  $(i\ i+1)$ ,  $1 \leq i \leq n-1$ .

This is an involution, and consists of  $\lfloor n/2 \rfloor$  (non-adjacent) transpositions

$$(1\ n)(2\ n-1)\dots, \text{ or } \sum_{k=1}^{n-1} k = \frac{n(n-1)}{2} \text{ adjacent transpositions:}$$

$$(n\ n-1)(n-1\ n-2)\dots(2\ 1)(n-1\ n-2)(n-2\ n-3)\dots,$$

so it thus has sign:

$$\text{sgn}(\rho_n) = (-1)^{\lfloor n/2 \rfloor} = (-1)^{n(n-1)/2} = \begin{cases} +1 & n \equiv 0, 1 \pmod{4} \\ -1 & n \equiv 2, 3 \pmod{4} \end{cases}$$

which is 4-periodic in  $n$ .

In  $S_{2n}$ , the perfect shuffle is the permutation that splits the set into 2 piles and interleaves them. Its sign is also  $(-1)^{\lfloor n/2 \rfloor}$ .

Note that the reverse on  $n$  elements and perfect shuffle on  $2n$  elements have the same sign; these are important to the classification of Clifford algebras, which are 8-periodic.

# Conjugacy classes

---

The conjugacy classes of  $S_n$  correspond to the cycle structures of permutations; that is, two elements of  $S_n$  are conjugate in  $S_n$  if and only if they consist of the same number of disjoint cycles of the same lengths. For instance, in  $S_5$ ,  $(1\ 2\ 3)(4\ 5)$  and  $(1\ 4\ 3)(2\ 5)$  are conjugate;  $(1\ 2\ 3)(4\ 5)$  and  $(1\ 2)(4\ 5)$  are not. A conjugating element of  $S_n$  can be constructed in "two line notation" by placing the "cycle notations" of the two conjugate permutations on top of one another. Continuing the previous example:

$$k = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 4 & 3 & 2 & 5 \end{pmatrix}$$

which can be written as the product of cycles, namely:  $(2\ 4)$ .

This permutation then relates  $(1\ 2\ 3)(4\ 5)$  and  $(1\ 4\ 3)(2\ 5)$  via conjugation, that is,

$$(2\ 4) \circ (1\ 2\ 3)(4\ 5) \circ (2\ 4) = (1\ 4\ 3)(2\ 5).$$

It is clear that such a permutation is not unique.

## Low degree groups

---

The low-degree symmetric groups have simpler and exceptional structure, and often must be treated separately.

### $S_0$ and $S_1$

The symmetric groups on the empty set and the singleton set are trivial, which corresponds to  $0! = 1! = 1$ . In this case the alternating group agrees with the symmetric group, rather than being an index 2 subgroup, and the sign map is trivial. In the case of  $S_0$ , its only member is the empty function.

### $S_2$

This group consists of exactly two elements: the identity and the permutation swapping the two points. It is a cyclic group and is thus abelian. In Galois theory, this corresponds to the fact that the quadratic formula gives a direct solution to the general quadratic polynomial after extracting only a single root. In invariant theory, the representation theory of the symmetric group on two points is quite simple and is seen as writing a function of two variables as a sum of its symmetric and anti-symmetric parts: Setting  $f_s(x, y) = f(x, y) + f(y, x)$ , and  $f_a(x, y) = f(x, y) - f(y, x)$ , one gets that  $2 \cdot f = f_s + f_a$ . This process is known as symmetrization.

### $S_3$

$S_3$  is the first nonabelian symmetric group. This group is isomorphic to the dihedral group of order 6, the group of reflection and rotation symmetries of an equilateral triangle, since these symmetries permute the three vertices of the triangle. Cycles of length two correspond to reflections, and cycles of length three are rotations. In Galois theory, the sign map from  $S_3$  to  $S_2$  corresponds to the resolving quadratic for a cubic polynomial, as discovered by Gerolamo Cardano, while the  $A_3$  kernel corresponds to the use of the discrete Fourier transform of order 3 in the solution, in the form of Lagrange resolvents.

### $S_4$

The group  $S_4$  is isomorphic to the group of proper rotations about opposite faces, opposite diagonals and opposite edges, 9, 8 and 6 permutations, of the cube.<sup>[5]</sup> Beyond the group  $A_4$ ,  $S_4$  has a Klein four-group  $V$  as a proper normal subgroup, namely the even transpositions  $\{(1), (1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3)\}$ , with quotient  $S_3$ . In Galois theory, this map corresponds to the resolving cubic to a quartic polynomial, which allows the quartic to be solved by radicals, as established by Lodovico Ferrari. The Klein group can be understood in

terms of the Lagrange resolvents of the quartic. The map from  $S_4$  to  $S_3$  also yields a 2-dimensional irreducible representation, which is an irreducible representation of a symmetric group of degree  $n$  of dimension below  $n - 1$ , which only occurs for  $n = 4$ .

## $S_5$

$S_5$  is the first non-solvable symmetric group. Along with the special linear group  $SL(2, 5)$  and the icosahedral group  $A_5 \times S_2$ ,  $S_5$  is one of the three non-solvable groups of order 120, up to isomorphism.  $S_5$  is the Galois group of the general quintic equation, and the fact that  $S_5$  is not a solvable group translates into the non-existence of a general formula to solve quintic polynomials by radicals. There is an exotic inclusion map  $S_5 \rightarrow S_6$  as a transitive subgroup; the obvious inclusion map  $S_n \rightarrow S_{n+1}$  fixes a point and thus is not transitive. This yields the outer automorphism of  $S_6$ , discussed below, and corresponds to the resolvent sextic of a quintic.

## $S_6$

Unlike all other symmetric groups,  $S_6$ , has an outer automorphism. Using the language of Galois theory, this can also be understood in terms of Lagrange resolvents. The resolvent of a quintic is of degree 6—this corresponds to an exotic inclusion map  $S_5 \rightarrow S_6$  as a transitive subgroup (the obvious inclusion map  $S_n \rightarrow S_{n+1}$  fixes a point and thus is not transitive) and, while this map does not make the general quintic solvable, it yields the exotic outer automorphism of  $S_6$ —see automorphisms of the symmetric and alternating groups for details.

Note that while  $A_6$  and  $A_7$  have an exceptional Schur multiplier (a triple cover) and that these extend to triple covers of  $S_6$  and  $S_7$ , these do not correspond to exceptional Schur multipliers of the symmetric group.

## Maps between symmetric groups

Other than the trivial map  $S_n \rightarrow 1 \cong S_0 \cong S_1$  and the sign map  $S_n \rightarrow S_2$ , the most notable homomorphisms between symmetric groups, in order of relative dimension, are:

- $S_4 \rightarrow S_3$  corresponding to the exceptional normal subgroup  $V < A_4 < S_4$ ;
- $S_6 \rightarrow S_6$  (or rather, a class of such maps up to inner automorphism) corresponding to the outer automorphism of  $S_6$ .
- $S_5 \rightarrow S_6$  as a transitive subgroup, yielding the outer automorphism of  $S_6$  as discussed above.

There are also a host of other homomorphisms  $S_m \rightarrow S_n$  where  $n > m$ .

## Properties

---

Symmetric groups are Coxeter groups and reflection groups. They can be realized as a group of reflections with respect to hyperplanes  $x_i = x_j$ ,  $1 \leq i < j \leq n$ . Braid groups  $B_n$  admit symmetric groups  $S_n$  as quotient groups.

Cayley's theorem states that every group  $G$  is isomorphic to a subgroup of the symmetric group on the elements of  $G$ , as a group acts on itself faithfully by (left or right) multiplication.

## Relation with alternating group

---

For  $n \geq 5$ , the alternating group  $A_n$  is simple, and the induced quotient is the sign map:  $A_n \rightarrow S_n \rightarrow S_2$  which is split by taking a transposition of two elements. Thus  $S_n$  is the semidirect product  $A_n \rtimes S_2$ , and has no other proper normal subgroups, as they would intersect  $A_n$  in either the identity (and thus themselves be the identity or a 2-element group, which is not normal), or in  $A_n$  (and thus themselves be  $A_n$  or  $S_n$ ).

$S_n$  acts on its subgroup  $A_n$  by conjugation, and for  $n \neq 6$ ,  $S_n$  is the full automorphism group of  $A_n$ :  $\text{Aut}(A_n) \cong S_n$ . Conjugation by even elements are inner automorphisms of  $A_n$  while the outer automorphism of  $A_n$  of order 2 corresponds to conjugation by an odd element. For  $n = 6$ , there is an exceptional outer automorphism of  $A_n$  so  $S_n$  is not the full automorphism group of  $A_n$ .

Conversely, for  $n \neq 6$ ,  $S_n$  has no outer automorphisms, and for  $n \neq 2$  it has no center, so for  $n \neq 2, 6$  it is a complete group, as discussed in automorphism group, below.

For  $n \geq 5$ ,  $S_n$  is an almost simple group, as it lies between the simple group  $A_n$  and its group of automorphisms.

$S_n$  can be embedded into  $A_{n+2}$  by appending the transposition  $(n+1, n+2)$  to all odd permutations, while embedding into  $A_{n+1}$  is impossible for  $n > 1$ .

## Generators and relations

---

The symmetric group on  $n$ -letters,  $S_n$ , may be described as follows. It has generators:  $\sigma_1, \dots, \sigma_{n-1}$  and relations:

- $\sigma_i^2 = 1$ ,
- $\sigma_i \sigma_j = \sigma_j \sigma_i$  if  $j \neq i \pm 1$ ,
- $(\sigma_i \sigma_{i+1})^3 = 1$ .

One thinks of  $\sigma_i$  as swapping the  $i$ th and  $(i+1)$ th position.

Other popular generating sets include the set of transpositions that swap 1 and  $i$  for  $2 \leq i \leq n$  and a set containing any  $n$ -cycle and a 2-cycle of adjacent elements in the  $n$ -cycle.

## Subgroup structure

---

A subgroup of a symmetric group is called a permutation group.

### Normal subgroups

The normal subgroups of the finite symmetric groups are well understood. If  $n \leq 2$ ,  $S_n$  has at most 2 elements, and so has no nontrivial proper subgroups. The alternating group of degree  $n$  is always a normal subgroup, a proper one for  $n \geq 2$  and nontrivial for  $n \geq 3$ ; for  $n \geq 3$  it is in fact the only non-identity proper normal subgroup of  $S_n$ , except when  $n = 4$  where there is one additional such normal subgroup, which is isomorphic to the Klein four group.

The symmetric group on an infinite set does not have an associated alternating group: not all elements can be written as a (finite) product of transpositions. However it does contain a normal subgroup  $S$  of permutations that fix all but finitely many elements, and such permutations can be classified as either even or odd. The even elements of  $S$  form the alternating subgroup  $A$  of  $S$ , and since  $A$  is even a characteristic subgroup of  $S$ , it is also a normal subgroup of the full symmetric group of the infinite set. The groups  $A$  and  $S$  are the only non-identity proper normal subgroups of the symmetric group on a countably infinite set. For more details see (Scott 1987, Ch. 11.3) or (Dixon & Mortimer 1996, Ch. 8.1).

### Maximal subgroups

The maximal subgroups of the finite symmetric groups fall into three classes: the intransitive, the imprimitive, and the primitive. The intransitive maximal subgroups are exactly those of the form  $\text{Sym}(k) \times \text{Sym}(n-k)$  for  $1 \leq k < n/2$ . The imprimitive maximal subgroups are exactly those of the form  $\text{Sym}(k) \text{ wr } \text{Sym}(n/k)$  where  $2 \leq k \leq n/2$  is a proper divisor of  $n$  and "wr" denotes the wreath product acting imprimitively. The primitive maximal subgroups are more difficult to identify, but with the assistance of the O'Nan–Scott theorem and the classification of finite simple groups, (Liebeck, Praeger & Saxl 1988) gave a fairly satisfactory description of the maximal subgroups of this type according to (Dixon & Mortimer 1996, p. 268).

# Sylow subgroups

The Sylow subgroups of the symmetric groups are important examples of p-groups. They are more easily described in special cases first:

The Sylow *p*-subgroups of the symmetric group of degree *p* are just the cyclic subgroups generated by *p*-cycles. There are  $(p-1)!/(p-1) = (p-2)!$  such subgroups simply by counting generators. The normalizer therefore has order  $p \cdot (p-1)$  and is known as a Frobenius group  $F_{p(p-1)}$  (especially for  $p=5$ ), and is the affine general linear group,  $AGL(1, p)$ .

The Sylow *p*-subgroups of the symmetric group of degree  $p^2$  are the wreath product of two cyclic groups of order *p*. For instance, when  $p=3$ , a Sylow 3-subgroup of  $Sym(9)$  is generated by  $a = (1\ 4\ 7)(2\ 5\ 8)(3\ 6\ 9)$  and the elements  $x = (1\ 2\ 3)$ ,  $y = (4\ 5\ 6)$ ,  $z = (7\ 8\ 9)$ , and every element of the Sylow 3-subgroup has the form  $a^i x^j y^k z^l$  for  $0 \leq i, j, k, l \leq 2$ .

The Sylow *p*-subgroups of the symmetric group of degree  $p^n$  are sometimes denoted  $W_p(n)$ , and using this notation one has that  $W_p(n+1)$  is the wreath product of  $W_p(n)$  and  $W_p(1)$ .

In general, the Sylow *p*-subgroups of the symmetric group of degree *n* are a direct product of  $a_i$  copies of  $W_p(i)$ , where  $0 \leq a_i \leq p-1$  and  $n = a_0 + p \cdot a_1 + \dots + p^k \cdot a_k$  (the base *p* expansion of *n*).

For instance,  $W_2(1) = C_2$  and  $W_2(2) = D_8$ , the dihedral group of order 8, and so a Sylow 2-subgroup of the symmetric group of degree 7 is generated by  $\{(1,3)(2,4), (1,2), (3,4), (5,6)\}$  and is isomorphic to  $D_8 \times C_2$ .

These calculations are attributed to (Kaloujnine 1948) and described in more detail in (Rotman 1995, p. 176). Note however that (Kerber 1971, p. 26) attributes the result to an 1844 work of Cauchy, and mentions that it is even covered in textbook form in (Netto 1882, §39–40).

# Transitive subgroups

A **transitive subgroup** of  $S_n$  is a subgroup whose action on  $\{1, 2, \dots, n\}$  is transitive. For example, the Galois group of a (finite) Galois extension is a transitive subgroup of  $S_n$ , for some *n*.

# Automorphism group

For  $n \neq 2, 6$ ,  $S_n$  is a complete group: its center and outer automorphism group are both trivial.

For  $n=2$ , the automorphism group is trivial, but  $S_2$  is not trivial: it is isomorphic to  $C_2$ , which is abelian, and hence the center is the whole group.

For  $n=6$ , it has an outer automorphism of order 2:  $Out(S_6) = C_2$ , and the automorphism group is a semidirect product

<i>n</i>	<b>Aut(<i>S<sub>n</sub></i>)</b>	<b>Out(<i>S<sub>n</sub></i>)</b>	<b>Z(<i>S<sub>n</sub></i>)</b>
<b><i>n</i> ≠ 2, 6</b>	<b><i>S<sub>n</sub></i></b>	1	1
<b><i>n</i> = 2</b>	1	1	<b><i>S<sub>2</sub></i></b>
<b><i>n</i> = 6</b>	<b><i>S<sub>6</sub></i> ⋊ <i>C<sub>2</sub></i></b>	<b><i>C<sub>2</sub></i></b>	1

**Aut(*S<sub>6</sub>*) = *S<sub>6</sub>* ⋊ *C<sub>2</sub>*.**

In fact, for any set *X* of cardinality other than 6, every automorphism of the symmetric group on *X* is inner, a result first due to (Schreier & Ulam 1937) according to (Dixon & Mortimer 1996, p. 259).

# Homology

The group homology of  $S_n$  is quite regular and stabilizes: the first homology (concretely, the abelianization) is:

$$H_1(S_n, \mathbf{Z}) = \begin{cases} 0 & n < 2 \\ \mathbf{Z}/2 & n \geq 2. \end{cases}$$



The first homology group is the abelianization, and corresponds to the sign map  $S_n \rightarrow S_2$  which is the abelianization for  $n \geq 2$ ; for  $n < 2$  the symmetric group is trivial. This homology is easily computed as follows:  $S_n$  is generated by involutions (2-cycles, which have order 2), so the only non-trivial maps  $S_n \rightarrow C_p$  are to  $S_2$  and all involutions are conjugate, hence map to the same element in the abelianization (since conjugation is trivial in abelian groups). Thus the only possible maps  $S_n \rightarrow S_2 \cong \{\pm 1\}$  send an involution to 1 (the trivial map) or to  $-1$  (the sign map). One must also show that the sign map is well-defined, but assuming that, this gives the first homology of  $S_n$ .

The second homology (concretely, the Schur multiplier) is:

$$H_2(S_n, \mathbf{Z}) = \begin{cases} 0 & n < 4 \\ \mathbf{Z}/2 & n \geq 4. \end{cases}$$

This was computed in (Schur 1911), and corresponds to the double cover of the symmetric group,  $2 \cdot S_n$ .

Note that the exceptional low-dimensional homology of the alternating group ( $H_1(\mathbf{A}_3) \cong H_1(\mathbf{A}_4) \cong \mathbf{C}_3$ , corresponding to non-trivial abelianization, and  $H_2(\mathbf{A}_6) \cong H_2(\mathbf{A}_7) \cong \mathbf{C}_6$ , due to the exceptional 3-fold cover) does not change the homology of the symmetric group; the alternating group phenomena do yield symmetric group phenomena – the map  $\mathbf{A}_4 \twoheadrightarrow \mathbf{C}_3$  extends to  $\mathbf{S}_4 \twoheadrightarrow \mathbf{S}_3$ , and the triple covers of  $\mathbf{A}_6$  and  $\mathbf{A}_7$  extend to triple covers of  $\mathbf{S}_6$  and  $\mathbf{S}_7$  – but these are not *homological* – the map  $\mathbf{S}_4 \twoheadrightarrow \mathbf{S}_3$  does not change the abelianization of  $\mathbf{S}_4$ , and the triple covers do not correspond to homology either.

The homology "stabilizes" in the sense of stable homotopy theory: there is an inclusion map  $S_n \rightarrow S_{n+1}$ , and for fixed  $k$ , the induced map on homology  $H_k(S_n) \rightarrow H_k(S_{n+1})$  is an isomorphism for sufficiently high  $n$ . This is analogous to the homology of families Lie groups stabilizing.

The homology of the infinite symmetric group is computed in (Nakaoka 1961), with the cohomology algebra forming a Hopf algebra.

## Representation theory

---

The representation theory of the symmetric group is a particular case of the representation theory of finite groups, for which a concrete and detailed theory can be obtained. This has a large area of potential applications, from symmetric function theory to problems of quantum mechanics for a number of identical particles.

The symmetric group  $S_n$  has order  $n!$ . Its conjugacy classes are labeled by partitions of  $n$ . Therefore, according to the representation theory of a finite group, the number of inequivalent irreducible representations, over the complex numbers, is equal to the number of partitions of  $n$ . Unlike the general situation for finite groups, there is in fact a natural way to parametrize irreducible representation by the same set that parametrizes conjugacy classes, namely by partitions of  $n$  or equivalently Young diagrams of size  $n$ .

Each such irreducible representation can be realized over the integers (every permutation acting by a matrix with integer coefficients); it can be explicitly constructed by computing the Young symmetrizers acting on a space generated by the Young tableaux of shape given by the Young diagram.

Over other fields the situation can become much more complicated. If the field  $K$  has characteristic equal to zero or greater than  $n$  then by Maschke's theorem the group algebra  $KS_n$  is semisimple. In these cases the irreducible representations defined over the integers give the complete set of irreducible representations (after reduction modulo the characteristic if necessary).

However, the irreducible representations of the symmetric group are not known in arbitrary characteristic. In this context it is more usual to use the language of modules rather than representations. The representation obtained from an irreducible representation defined over the integers by reducing modulo the characteristic will not in general be irreducible. The modules so constructed are called Specht modules, and every irreducible does arise inside some such module. There are now fewer irreducibles, and although they can be classified they are very poorly understood. For example, even their dimensions are not known in general.

The determination of the irreducible modules for the symmetric group over an arbitrary field is widely regarded as one of the most important open problems in representation theory.

## See also

---

- [History of group theory](#)
- [Symmetric inverse semigroup](#)
- [Signed symmetric group](#)
- [Generalized symmetric group](#)

## References

---

1. Jacobson (2009), p. 31.
  2. Jacobson (2009), p. 32. Theorem 1.1.
  3. "Symmetric Group is not Abelian/Proof 1" ([https://proofwiki.org/wiki/Symmetric\\_Group\\_is\\_not\\_Abelian/Proof\\_1](https://proofwiki.org/wiki/Symmetric_Group_is_not_Abelian/Proof_1)).
  4. modern algebra Author A. R. Vasishtha, A. K. Vasishtha Publisher Krishna Prakashan Media
  5. Die Untergruppenverbände der Gruppen der ordnung weniger als 100, Habilitationsschrift, J. Neubuser, Universität Kiel, Germany, 1967.
- Cameron, Peter J. (1999), *Permutation Groups*, London Mathematical Society Student Texts, **45**, [Cambridge University Press](#), ISBN 978-0-521-65378-7
  - Dixon, John D.; Mortimer, Brian (1996), *Permutation groups*, Graduate Texts in Mathematics, **163**, Berlin, New York: Springer-Verlag, ISBN 978-0-387-94599-6, MR 1409812 (<https://www.ams.org/mathscinet-getitem?mr=1409812>)
  - Jacobson, Nathan (2009), *Basic algebra*, **1** (2nd ed.), Dover, ISBN 978-0-486-47189-1.
  - Kaloujnine, Léo (1948), "La structure des p-groupes de Sylow des groupes symétriques finis" ([http://www.numdam.org/item?id=ASENS\\_1948\\_3\\_65\\_239\\_0](http://www.numdam.org/item?id=ASENS_1948_3_65_239_0)), *Annales Scientifiques de l'École Normale Supérieure*, Série 3, **65**: 239–276, ISSN 0012-9593 (<https://www.worldcat.org/issn/0012-9593>), MR 0028834 (<https://www.ams.org/mathscinet-getitem?mr=0028834>)
  - Kerber, Adalbert (1971), *Representations of permutation groups. I*, Lecture Notes in Mathematics, Vol. 240, **240**, Berlin, New York: Springer-Verlag, doi:10.1007/BFb0067943 (<https://doi.org/10.1007%2FBFb0067943>), ISBN 978-3-540-05693-5, MR 0325752 (<https://www.ams.org/mathscinet-getitem?mr=0325752>)
  - Liebeck, M.W.; Praeger, C.E.; Saxl, J. (1988), "On the O'Nan-Scott theorem for finite primitive permutation groups", *J. Austral. Math. Soc.*, **44** (3): 389–396, doi:10.1017/S144678870003216X (<https://doi.org/10.1017%2FS144678870003216X>)
  - Nakaoka, Minoru (March 1961), "Homology of the Infinite Symmetric Group", *The Annals of Mathematics*, **2**, Annals of Mathematics, **73** (2): 229–257, doi:10.2307/1970333 (<https://doi.org/10.2307%2F1970333>), JSTOR 1970333 (<https://www.jstor.org/stable/1970333>)
  - Netto, E. (1882), *Substitutionentheorie und ihre Anwendungen auf die Algebra* (in German), Leipzig: Teubner, JFM 14.0090.01 (<https://zbmath.org/?format=complete&q=an:14.0090.01>)
  - Scott, W.R. (1987), *Group Theory*, New York: Dover Publications, pp. 45–46, ISBN 978-0-486-65377-8
  - Schur, Issai (1911), "Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen", *Journal für die reine und angewandte Mathematik*, **139**: 155–250, doi:10.1515/crll.1911.139.155 (<https://doi.org/10.1515%2FCrll.1911.139.155>)
  - Schreier, J.; Ulam, Stanislaw (1936), "Über die Automorphismen der Permutationsgruppe der natürlichen Zahlenfolge" (<http://matwbn.icm.edu.pl/ksiazki/fm/fm28/fm28128.pdf>) (PDF), *Fundam. Math.* (in German), **28**: 258–260, Zbl 0016.20301 (<https://zbmath.org/?format=complete&q=an:0016.20301>)

## External links

---

- Hazewinkel, Michiel, ed. (2001) [1994], "Symmetric group" (<https://www.encyclopediaofmath.org/index.php?title=p/s091670>), *Encyclopedia of Mathematics*, Springer Science+Business Media B.V. / Kluwer Academic Publishers, ISBN 978-1-55608-010-4

- Weisstein, Eric W. "Symmetric group" (<http://mathworld.wolfram.com/SymmetricGroup.html>). *MathWorld*.
  - Weisstein, Eric W. "Symmetric group graph" (<http://mathworld.wolfram.com/SymmetricGroupGraph.html>). *MathWorld*.
  - Marcus du Sautoy: Symmetry, reality's riddle ([http://www.ted.com/talks/marcus\\_du\\_sautoy\\_symmetry\\_reality\\_s\\_riddle.html](http://www.ted.com/talks/marcus_du_sautoy_symmetry_reality_s_riddle.html)) (video of a talk)
  - OEIS Entries dealing with the Symmetric Group (<http://oeis.org/search?q=Symmetric+Group>)
- 

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Symmetric\\_group&oldid=826535064](https://en.wikipedia.org/w/index.php?title=Symmetric_group&oldid=826535064)"

---

**This page was last edited on 19 February 2018, at 18:07.**

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.