

## 2-4 Recurrences


魏恒峰

hfwei@nju.edu.cn

2018 年 04 月 18 日




## Triangulating Polygons



## Ear Lemma (Problem 4.1 – 16)




## # of triangles (Problem 4.1 – 17)




$$T(n) = n - 2$$


## # of triangles (Problem 4.1 – 17)





$$T(n) = n - 2$$



## # of triangles (Problem 4.1 – 17)



$$T(n) = n - 2$$



## Lemma (Ear Lemma)

A triangle has 3 ears, and a larger *triangulated* polygon has at least 2 non-adjacent ears.

## Lemma (Ear Lemma)

*A triangle has 3 ears, and a larger **triangulated** polygon has at least 2 non-adjacent ears.*

*Q : Can every polygon be triangulated?*

## Theorem (Existence of Triangulation)

*Any polygon can be triangulated.*

## Theorem (Existence of Triangulation)

*Any polygon can be triangulated.*

Proof.

*"To triangulate a polygon one keeps adding diagonals connecting pairs of vertices until no more diagonals can be added.*

## Theorem (Existence of Triangulation)

*Any polygon can be triangulated.*

Proof.

*"To triangulate a polygon one keeps adding diagonals connecting pairs of vertices until no more diagonals can be added.*

*These diagonals must lie entirely interior to the polygon and are not allowed to intersect.*

## Theorem (Existence of Triangulation)

*Any polygon can be triangulated.*

Proof.

*"To triangulate a polygon one keeps adding diagonals connecting pairs of vertices until no more diagonals can be added.*

*These diagonals must lie entirely interior to the polygon and are not allowed to intersect.*

*They break the interior of the polygon into a number of triangles, because any larger polygon can be split by adding a diagonal."*



## Theorem (Existence of Triangulation)

*Any polygon can be triangulated.*

Proof.

*"To triangulate a polygon one keeps adding diagonals connecting pairs of vertices until no more diagonals can be added.*

*These diagonals must lie entirely interior to the polygon and are not allowed to intersect.*

*They break the interior of the polygon into a number of triangles, because any larger polygon can be split by adding a diagonal."*



*"(This fact is perhaps not obvious,  
but we won't get sidetracked by proving it here.)"*

## Theorem (Existence of Triangulation)

*Any polygon can be triangulated.*

Proof.

*"To triangulate a polygon one keeps adding diagonals connecting pairs of vertices until no more diagonals can be added.*

*These diagonals must lie entirely interior to the polygon and are not allowed to intersect.*

*They break the interior of the polygon into a number of triangles, because any larger polygon can be split by adding a diagonal."*



*"(This fact is perhaps not obvious,  
but we won't get sidetracked by proving it here.)"*

## Theorem (Existence of Diagonal)

*Every polygon with  $n > 3$  has a diagonal.*

## Theorem (Existence of Diagonal)

*Every polygon with  $n > 3$  has a diagonal.*

## Definition (Convex Vertex)


A vertex  $v$  is **convex** if the *interior* angle at  $v$  is less than  $180^\circ$ .

## Theorem (Existence of Diagonal)

*Every polygon with  $n > 3$  has a diagonal.*


## Definition (Convex Vertex)

A vertex  $v$  is **convex** if the *interior* angle at  $v$  is less than  $180^\circ$ .




## Theorem (Coloring)

*Any triangulated polygon polygon is 3-colorable.*



## Theorem (Coloring)

Any triangulated polygon is 3-colorable.



# The Art Gallery Problem




## The Art Gallery Problem




Q : How many “BIG BROs” to hire?

## The Art Gallery Problem



*Q* : How many “BIG BROs” to hire?



## Theorem (The Art Gallery Theorem ( $O$ ))


For any art gallery with  $n$  walls,  $\lfloor \frac{n}{3} \rfloor$  "BIG BROs" suffice.

Theorem (The Art Gallery Theorem ( $\Omega$ ))

For any art gallery with  $n$  walls,  $\lfloor \frac{n}{3} \rfloor$  "BIG BROs" suffice.


## Theorem (The Art Gallery Theorem ( $\Omega$ ))

For any art gallery with  $n$  walls,  $\lfloor \frac{n}{3} \rfloor$  "BIG BROs" suffice.



$$n = 3m$$

# Fish Recurrence



## Fish Recurrence (Problem 4.2 – 8)

At the end of each year, a state fish hatchery puts 2000 fish into a lake.

The number of fish in the lake at the beginning of the year doubles by the end of the year due to reproduction.

Give a recurrence for the number of fish in the lake after  $n$  years, and solve the recurrence.

## Fish Recurrence (Problem 4.2 – 8)

At the end of each year, a state fish hatchery puts 2000 fish into a lake.

The number of fish in the lake at the beginning of the year doubles by the end of the year due to reproduction.

Give a recurrence for the number of fish in the lake after  $n$  years, and solve the recurrence.

$$T(n) = \# \text{ of fish at the end of the } n\text{-th year}$$

## Fish Recurrence (Problem 4.2 – 8)

At the end of each year, a state fish hatchery puts 2000 fish into a lake.

The number of fish in the lake at the beginning of the year doubles by the end of the year due to reproduction.

Give a recurrence for the number of fish in the lake after  $n$  years, and solve the recurrence.

$T(n) = \# \text{ of fish at the end of the } n\text{-th year}$

$$T(n) = 2T(n - 1) + 2000$$

## Fish Recurrence (Problem 4.2 – 8)

At the end of each year, a state fish hatchery puts 2000 fish into a lake.

The number of fish in the lake at the beginning of the year doubles by the end of the year due to reproduction.

Give a recurrence for the number of fish in the lake after  $n$  years, and solve the recurrence.

$T(n) = \# \text{ of fish at the end of the } n\text{-th year}$

$$T(n) = 2T(n - 1) + 2000$$

$$T(0) = 2000$$

# Solving Recurrence





Why again?



base cases

## First-order Linear Recurrence (CS 4.2 – 11)

$$T(n) = \begin{cases} 1 & n = 0 \\ 2T(n - 1) + n2^n & n > 0 \end{cases}$$

## First-order Linear Recurrence (CS 4.2 – 11)

$$T(n) = \begin{cases} 1 & n = 0 \\ 2T(n - 1) + n2^n & n > 0 \end{cases}$$

Theorem (First-order Linear Recurrences with Constant Coefficients (CS Theorem 4.5))

$$T(n) = \begin{cases} a & n = 0 \\ rT(n - 1) + g(n) & n > 0 \end{cases}$$

## First-order Linear Recurrence (CS 4.2 – 11)

$$T(n) = \begin{cases} 1 & n = 0 \\ 2T(n - 1) + n2^n & n > 0 \end{cases}$$

Theorem (First-order Linear Recurrences with Constant Coefficients (CS Theorem 4.5))

$$T(n) = \begin{cases} a & n = 0 \\ rT(n - 1) + g(n) & n > 0 \end{cases}$$

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i)$$

## Theorem (First-order Linear Recurrences)

$T(n) = \textcolor{red}{x_n} T(n-1) + y_n \quad \text{for } n > 0 \text{ with } T(0) = 0$

$$T(n) = y_0 + \sum_{1 \leq j < n} y_j x_{j+1} x_{j+2} \cdots x_n$$

## Theorem (First-order Linear Recurrences)

$$T(n) = \textcolor{red}{x_n} T(n-1) + y_n \quad \text{for } n > 0 \text{ with } T(0) = 0$$

$$T(n) = y_0 + \sum_{1 \leq j < n} y_j x_{j+1} x_{j+2} \cdots x_n$$

Proof.

$$\underbrace{\frac{T(n)}{x_n x_{n-1} \cdots x_1}}_{\text{summation factor}} = \frac{T(n-1)}{x_{n-1} \cdots x_1} + \frac{y_n}{x_n x_{n-1} \cdots x_1}$$

## Theorem (First-order Linear Recurrences)

$$T(n) = \textcolor{red}{x_n} T(n-1) + y_n \quad \text{for } n > 0 \text{ with } T(0) = 0$$

$$T(n) = y_0 + \sum_{1 \leq j < n} y_j x_{j+1} x_{j+2} \cdots x_n$$

Proof.

$$\underbrace{\frac{T(n)}{x_n x_{n-1} \cdots x_1}}_{\text{summation factor}} = \frac{T(n-1)}{x_{n-1} \cdots x_1} + \frac{y_n}{x_n x_{n-1} \cdots x_1}$$

$$S(n) \triangleq \frac{T(n)}{x_n x_{n-1} \cdots x_1}$$



$$T(n) = \left(1 + \frac{1}{n}\right)T(n-1) + 2 \quad \text{for } n > 1 \text{ with } T(1) = 0$$

$$T(n) = \left(1 + \frac{1}{n}\right)T(n-1) + 2 \quad \text{for } n > 1 \text{ with } T(1) = 0$$

$$x_n = 1 + \frac{1}{n} \implies$$

$$T(n) = \left(1 + \frac{1}{n}\right)T(n-1) + 2 \quad \text{for } n > 1 \text{ with } T(1) = 0$$

$$x_n = 1 + \frac{1}{n} \implies x_n x_{n-1} \cdots x_1 = n + 1$$

$$T(n) = \left(1 + \frac{1}{n}\right)T(n-1) + 2 \quad \text{for } n > 1 \text{ with } T(1) = 0$$

$$x_n = 1 + \frac{1}{n} \implies x_n x_{n-1} \cdots x_1 = n + 1$$

$$\frac{T(n)}{n+1} = \frac{T(n-1)}{n} + \frac{2}{n+1} \quad \text{for } n > 1$$

$$T(n) = \left(1 + \frac{1}{n}\right)T(n-1) + 2 \quad \text{for } n > 1 \text{ with } T(1) = 0$$

$$x_n = 1 + \frac{1}{n} \implies x_n x_{n-1} \cdots x_1 = n + 1$$

$$\frac{T(n)}{n+1} = \frac{T(n-1)}{n} + \frac{2}{n+1} \quad \text{for } n > 1$$

$$\frac{T(n)}{n+1} = \frac{T(1)}{2} + 2 \sum_{3 \leq k \leq n+1} \frac{1}{k}$$

$$T(n) = \left(1 + \frac{1}{n}\right)T(n-1) + 2 \quad \text{for } n > 1 \text{ with } T(1) = 0$$

$$x_n = 1 + \frac{1}{n} \implies x_n x_{n-1} \cdots x_1 = n + 1$$

$$\frac{T(n)}{n+1} = \frac{T(n-1)}{n} + \frac{2}{n+1} \quad \text{for } n > 1$$

$$\frac{T(n)}{n+1} = \frac{T(1)}{2} + 2 \sum_{3 \leq k \leq n+1} \frac{1}{k}$$

$$T(n) = 2(n+1)\left(H_{n+1} - \frac{3}{2}\right)$$

## After-class Exercise

$$T(n) = T(n-1) - \frac{2T(n-1)}{n} + 2 \left(1 - \frac{2T(n-1)}{n}\right), n > 0 \text{ with } T(0) = 0$$

## After-class Exercise

$$T(n) = T(n-1) - \frac{2T(n-1)}{n} + 2 \left(1 - \frac{2T(n-1)}{n}\right), n > 0 \text{ with } T(0) = 0$$



## Theorem (Linear Recurrences with Constant Coefficients)

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$a_0, a_1, \dots, a_{t-1}$$

## Theorem (Linear Recurrences with Constant Coefficients)

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$a_0, a_1, \dots, a_{t-1}$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

## Theorem (Linear Recurrences with Constant Coefficients)

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$a_0, a_1, \dots, a_{t-1}$$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

$$\beta_1(m_1), \beta_2(m_2), \dots, \beta_i(m_i), \dots, \beta_k(m_k)$$

$$m_1 + m_2 + \cdots + m_k = t$$

## Theorem (Linear Recurrences with Constant Coefficients)

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$a_0, a_1, \dots, a_{t-1}$$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

$$\beta_1(m_1), \beta_2(m_2), \dots, \beta_i(m_i), \dots, \beta_k(m_k)$$

$$m_1 + m_2 + \cdots + m_k = t$$

$$a_n = \sum_{0 \leq j < m_1} c_{1j} n^j \beta_1^n + \sum_{0 \leq j < m_2} c_{2j} n^j \beta_2^n + \cdots + \sum_{0 \leq j < m_k} c_{kj} n^j \beta_k^n$$

$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t}$  for  $n \geq t$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

Proof.

$$\beta \ (m = 2)$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

Proof.

$$\beta \ (m=2)$$

$$\beta^n = r_1 \beta^{n-1} + r_2 \beta^{n-2} + \cdots + r_t \beta^{n-t} \quad \text{for } n \geq t$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

Proof.

$$\beta \ (m=2)$$

$$\beta^n = r_1 \beta^{n-1} + r_2 \beta^{n-2} + \cdots + r_t \beta^{n-t} \quad \text{for } n \geq t$$

$$\beta^{n-t} q(\beta) = 0$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

Proof.

$$\beta \ (m=2)$$

$$\beta^n = r_1 \beta^{n-1} + r_2 \beta^{n-2} + \cdots + r_t \beta^{n-t} \quad \text{for } n \geq t$$

$$\beta^{n-t} q(\beta) = 0$$

$$n \beta^n = r_1(n-1) \beta^{n-1} + r_2(n-2) \beta^{n-2} + \cdots + r_t(n-t) \beta^{n-t} \quad \text{for } n \geq t$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$q(x) \equiv x^t - r_1 x^{t-1} - r_2 x^{t-2} - \cdots - r_t$$

Proof.

$$\beta \ (m=2)$$

$$\beta^n = r_1 \beta^{n-1} + r_2 \beta^{n-2} + \cdots + r_t \beta^{n-t} \quad \text{for } n \geq t$$

$$\beta^{n-t} q(\beta) = 0$$

$$n \beta^n = r_1(n-1) \beta^{n-1} + r_2(n-2) \beta^{n-2} + \cdots + r_t(n-t) \beta^{n-t} \quad \text{for } n \geq t$$

$$\beta^{n-t} ((n-t) q(\beta) + \beta q'(\beta)) = 0$$

$$a_n = 5a_{n-1} - 6a_{n-2} \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$a_n = 5a_{n-1} - 6a_{n-2} \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$\boxed{x^2 - 5x + 6 = (x - 2)(x - 3) = 0} \implies x = 2, 3$$

$$a_n = 5a_{n-1} - 6a_{n-2} \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$x^2 - 5x + 6 = (x - 2)(x - 3) = 0 \implies x = 2, 3$$

$$a_n = c_0 3^n + c_1 2^n$$

$$a_n = 5a_{n-1} - 6a_{n-2} \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$x^2 - 5x + 6 = (x - 2)(x - 3) = 0 \implies x = 2, 3$$

$$a_n = c_0 3^n + c_1 2^n$$

$$a_0 = 0 = c_0 + c_1$$

$$a_1 = 1 = 3c_0 + 2c_1$$

$$a_n = 5a_{n-1} - 6a_{n-2} \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$x^2 - 5x + 6 = (x - 2)(x - 3) = 0 \implies x = 2, 3$$

$$a_n = c_0 3^n + c_1 2^n$$

$$a_0 = 0 = c_0 + c_1$$

$$a_1 = 1 = 3c_0 + 2c_1$$

$$a_n = 3^n - 2^n$$

$$a_n = 5a_{n-1} - 8a_{n-2} + 4a_{n-3}, n \geq 3 \quad (a_0 = 0, a_1 = 1, a_2 = 4)$$

$$a_n = 5a_{n-1} - 8a_{n-2} + 4a_{n-3}, n \geq 3 \quad (a_0 = 0, a_1 = 1, a_2 = 4)$$

$$x^3 - 5x^2 + 8x - 4 = 0$$

$$a_n = 5a_{n-1} - 8a_{n-2} + 4a_{n-3}, n \geq 3 \quad (a_0 = 0, a_1 = 1, a_2 = 4)$$

$$x^3 - 5x^2 + 8x - 4 = 0$$

$$(x-1)(x-2)^2 = 0 \implies x_1 = 1, x_2 = 2, x'_2 = 2$$

$$a_n = 5a_{n-1} - 8a_{n-2} + 4a_{n-3}, n \geq 3 \quad (a_0 = 0, a_1 = 1, a_2 = 4)$$

$$x^3 - 5x^2 + 8x - 4 = 0$$

$$(x-1)(x-2)^2 = 0 \implies x_1 = 1, x_2 = 2, x'_2 = 2$$

$$a_n = c_1 \cdot 1^n + c_2 \cdot 2^n + c'_2 \cdot n2^n$$

$$a_n = 5a_{n-1} - 8a_{n-2} + 4a_{n-3}, n \geq 3 \quad (a_0 = 0, a_1 = 1, a_2 = 4)$$

$$x^3 - 5x^2 + 8x - 4 = 0$$

$$(x-1)(x-2)^2 = 0 \implies x_1 = 1, x_2 = 2, x'_2 = 2$$

$$a_n = c_1 \cdot 1^n + c_2 \cdot 2^n + c'_2 \cdot n2^n$$

$$a_0 = 0 = c_1 + c_2$$

$$a_1 = 1 = c_1 + 2c_2 + 2c'_2$$

$$a_2 = 4 = c_1 + 4c_2 + 8c'_2$$

$$a_n = 5a_{n-1} - 8a_{n-2} + 4a_{n-3}, n \geq 3 \quad (a_0 = 0, a_1 = 1, a_2 = 4)$$

$$x^3 - 5x^2 + 8x - 4 = 0$$

$$(x-1)(x-2)^2 = 0 \implies x_1 = 1, x_2 = 2, x'_2 = 2$$

$$a_n = c_1 \cdot 1^n + c_2 \cdot 2^n + c'_2 \cdot n2^n$$

$$a_0 = 0 = c_1 + c_2$$

$$a_1 = 1 = c_1 + 2c_2 + 2c'_2$$

$$a_2 = 4 = c_1 + 4c_2 + 8c'_2$$

$$a_n = n2^{n-1}$$

$$a_n = 2a_{n-1} - a_{n-2} + 2a_{n-3}, \quad n \geq 3 \quad (a_0 = 1, a_1 = 0, a_2 = -1)$$

$$a_n = 2a_{n-1} - a_{n-2} + 2a_{n-3}, \quad n \geq 3 \quad (a_0 = 1, a_1 = 0, a_2 = -1)$$

$$\boxed{x^3 - 2x^2 + x - 2} = (x^2 + 1)(x - 2) = 0 \implies x = 2, i, -i$$

$$a_n = 2a_{n-1} - a_{n-2} + 2a_{n-3}, \quad n \geq 3 \quad (a_0 = 1, a_1 = 0, a_2 = -1)$$

$$\boxed{x^3 - 2x^2 + x - 2} = (x^2 + 1)(x - 2) = 0 \implies x = 2, i, -i$$

$$\boxed{a_n = c_1 2^n + c_2 i^n + c_3 (-i)^n}$$

$$a_n = 2a_{n-1} - a_{n-2} + 2a_{n-3}, \quad n \geq 3 \quad (a_0 = 1, a_1 = 0, a_2 = -1)$$

$$\boxed{x^3 - 2x^2 + x - 2} = (x^2 + 1)(x - 2) = 0 \implies x = 2, i, -i$$

$$\boxed{a_n = c_1 2^n + c_2 i^n + c_3 (-i)^n}$$

$$a_0 = 1 = c_1 + c_2 + c_3$$

$$a_1 = 0 = 2c_1 + c_2 i - c_3 i$$

$$a_2 = -1 = 4c_1 - c_2 - c_3$$

$$a_n = 2a_{n-1} - a_{n-2} + 2a_{n-3}, \quad n \geq 3 \quad (a_0 = 1, a_1 = 0, a_2 = -1)$$

$$\boxed{x^3 - 2x^2 + x - 2} = (x^2 + 1)(x - 2) = 0 \implies x = 2, i, -i$$

$$\boxed{a_n = c_1 2^n + c_2 i^n + c_3 (-i)^n}$$

$$a_0 = 1 = c_1 + c_2 + c_3$$

$$a_1 = 0 = 2c_1 + c_2 i - c_3 i$$

$$a_2 = -1 = 4c_1 - c_2 - c_3$$

$$\boxed{a_n = \frac{1}{2} i^n (1 + (-1)^n)}$$

$$a_n = 2a_{n-1} - a_{n-2} + 2a_{n-3}, \quad n \geq 3 \quad (a_0 = 1, a_1 = 0, a_2 = -1)$$

$$\boxed{x^3 - 2x^2 + x - 2} = (x^2 + 1)(x - 2) = 0 \implies x = 2, i, -i$$

$$\boxed{a_n = c_1 2^n + c_2 i^n + c_3 (-i)^n}$$

$$a_0 = 1 = c_1 + c_2 + c_3$$

$$a_1 = 0 = 2c_1 + c_2 i - c_3 i$$

$$a_2 = -1 = 4c_1 - c_2 - c_3$$

$$\boxed{a_n = \frac{1}{2} i^n (1 + (-1)^n)}$$

$$1, \textcolor{red}{0}, \textcolor{blue}{-1}, \textcolor{red}{0}, \textcolor{blue}{1}, \textcolor{red}{0}, \textcolor{blue}{-1}, \textcolor{red}{0}, \textcolor{blue}{1}, \textcolor{red}{0}, \textcolor{blue}{-1}, \textcolor{red}{0} \dots$$

## After-class Exercise

Given initial conditions  $a_0, a_1$ , and  $a_2$  for which the growth rate of the solution to

$$a_n = 2a_{n-1} + a_{n-2} - 2a_{n-3}, \quad n > 2$$

is

- (i) constant,
- (ii) exponential,
- (iii) fluctuating in sign

## After-class Exercise

Given initial conditions  $a_0, a_1$ , and  $a_2$  for which the growth rate of the solution to

$$a_n = 2a_{n-1} + a_{n-2} - 2a_{n-3}, \quad n > 2$$

is

- (i) constant,
- (ii) exponential,
- (iii) fluctuating in sign



## First-order Linear Non-homogeneous Recurrences

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} + \textcolor{red}{r} \quad \text{for } n \geq t$$

# First-order Linear Non-homogeneous Recurrences

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} + \textcolor{red}{r} \quad \text{for } n \geq t$$

7. 费波那契数列的定义如下： $F_1 = 1, F_2 = 1, F_n = F_{n-1} + F_{n-2}$  ( $n \geq 3$ )。如果用下面的函数计算费波那契数列的第  $n$  项，则其时间复杂度为（ ）。

```
int F(int n)
{
    if (n <= 2)
        return 1;
    else
        return F(n - 1) + F(n - 2);
}
```

- A.  $O(1)$       B.  $O(n)$       C.  $O(n^2)$       D.  $O(F_n)$

$$F(n) = F(n - 1) + F(n - 2) + 2, \quad n \geq 3 \quad (F(1) = F(2) = 0)$$

# First-order Linear Non-homogeneous Recurrences

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} + \textcolor{red}{r} \quad \text{for } n \geq t$$

7. 费波那契数列的定义如下： $F_1 = 1, F_2 = 1, F_n = F_{n-1} + F_{n-2}$  ( $n \geq 3$ )。如果用下面的函数计算费波那契数列的第  $n$  项，则其时间复杂度为（ ）。

```
int F(int n)
{
    if (n <= 2)
        return 1;
    else
        return F(n - 1) + F(n - 2);
}
```

- A.  $O(1)$       B.  $O(n)$       C.  $O(n^2)$       D.  $O(F_n)$

$$F(n) = F(n - 1) + F(n - 2) + 2, \quad n \geq 3 \quad (F(1) = F(2) = 0)$$

$$T(n, k) = \begin{cases} 0, & k = 0 \vee n = k \\ T(n - 1, k) + T(n - 1, k - 1) + c, & \text{o.w.} \end{cases}$$

$$a_n = 5a_{n-1} - 6a_{n-2} + 2, \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$a_n = 5a_{n-1} - 6a_{n-2} + 2, \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$a_n = c_0 3^n + c_1 2^n + c_2$$

$$a_n = 5a_{n-1} - 6a_{n-2} + 2, \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$a_n = c_0 3^n + c_1 2^n + c_2$$

$$c_2 = 5c_2 - 6c_2 + 2 \implies c_2 = 1$$

$$a_n = 5a_{n-1} - 6a_{n-2} + 2, \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$a_n = c_0 3^n + c_1 2^n + c_2$$

$$c_2 = 5c_2 - 6c_2 + 2 \implies c_2 = 1$$

$$a_0 = 0 = c_0 + c_1 + 1$$

$$a_1 = 1 = 3c_0 + 2c_1 + 1$$

$$a_n = 5a_{n-1} - 6a_{n-2} + 2, \quad n \geq 2 \quad (a_0 = 0, a_1 = 1)$$

$$a_n = c_0 3^n + c_1 2^n + c_2$$

$$c_2 = 5c_2 - 6c_2 + 2 \implies c_2 = 1$$

$$a_0 = 0 = c_0 + c_1 + 1$$

$$a_1 = 1 = 3c_0 + 2c_1 + 1$$

$$a_n = 2 \cdot 3^n - 3 \cdot 2^n + 1$$

## More Issues about Linear Recurrences

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} + g(n) \quad \text{for } n \geq t$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} + g(n) \quad \text{for } n \geq t$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$t \geq 5$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$

$$t \geq 5$$

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_t a_{n-t} \quad \text{for } n \geq t$$


$$t \geq 5$$

## Generating Functions and Asymptotic Analysis


$$a_n = f_1(n)a_{n-1} + f_2(n)a_{n-2} + \cdots + f_t(n)a_{n-t} \quad \text{for } n \geq t$$

$$a_n = f_1(n)a_{n-1} + f_2(n)a_{n-2} + \cdots + f_t(n)a_{n-t} \quad \text{for } n \geq t$$

## Genearating Functions



# Thank You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn