

# 3-8 Cool? We are APSP Algorithms.

Hengfeng Wei

hfwei@nju.edu.cn

November 19, 2018





Please Help Me Out Here.

## Definition (Shortest Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{w(p) : u \rightsquigarrow^p v\} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

Path *vs.* Simple path

## Definition (Shortest Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{w(p) : u \rightsquigarrow^p v\} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

Path *vs.* Simple path

Shortest-path Problem *vs.* Longest-path Problem

Digraph *vs.* Undirected Graph

# Single Source      Digraph

Shortest-path Problem    *vs.*    Longest-path Problem

## Single Source      Digraph

Shortest-path Problem    *vs.*    Longest-path Problem

$$\text{SP in } G \iff \text{LP in } -G$$

## Single Source      Digraph

Shortest-path Problem    *vs.*    Longest-path Problem

$$\text{SP in } G \iff \text{LP in } -G$$


$O(VE)$  (Bellman-Ford)    *vs.*    NP-hard (I just told you.)

## Single Source      Digraph

Shortest-path Problem    *vs.*    Longest-path Problem

$$\text{SP in } G \iff \text{LP in } -G$$

$O(VE)$  (Bellman-Ford)    *vs.*    NP-hard (I just told you.)






Single Source

Digraph

Shortest-path Problem *vs.* Longest-path Problem

$$\text{SP in } G \iff \text{LP in } -G$$

$O(VE)$  (Bellman-Ford) *vs.* NP-hard (I just told you.)



## Definition (Shortest Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Path}}\} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

## Definition (Shortest Path)

$G = (V, E, w)$  : weighted digraph

$$\delta(u, v) = \begin{cases} \min \{w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Path}}\} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

$Q$  : How does Bellman-Ford Handle with Negative-weight Cycles?

## Definition (Shortest Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Path}}\} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

$Q$  : How does Bellman-Ford Handle with Negative-weight Cycles?

$A$  : Report it and Treat Shortest Path as “Undefined”.

## Definition (Shortest Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Path}}\} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

$Q$  : How does Bellman-Ford Handle with Negative-weight Cycles?

$A$  : Report it and Treat Shortest Path as “Undefined”.

$$O(VE)$$

## Definition (Shortest Simple Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{ w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Simple Path}} \} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

## Definition (Shortest Simple Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{ w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Simple Path}} \} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

**Q : How Should an Algorithm for Shortest Simple Path Problem Handle with Negative-weight Cycles?**

## Definition (Shortest Simple Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{ w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Simple Path}} \} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

$Q$  : How Should an Algorithm for Shortest Simple Path Problem Handle with Negative-weight Cycles?

$A$  : Still to Find Shortest Simple Path.



## Definition (Shortest Simple Path)

$G = (V, E, w) : \text{weighted digraph}$

$$\delta(u, v) = \begin{cases} \min \{ w(p) : \underbrace{u \rightsquigarrow^p v}_{\text{Simple Path}} \} & \text{if } u \rightsquigarrow v \\ \infty & \text{o.w.} \end{cases}$$

$Q$  : How Should an Algorithm for Shortest Simple Path Problem Handle with Negative-weight Cycles?

$A$  : Still to Find Shortest Simple Path.

NP-hard

# Shortest Path Problem

## Shortest Simple Path Problem

Shortest Path Problem

Longest Path Problem

Shortest Simple Path Problem

Longest Simple Path Problem

Shortest Path Problem



Longest Path Problem

Shortest Simple Path Problem



Longest Simple Path Problem

Single Source

Undirected Graph

Single Source      Undirected Graph

Negative-weight edges allowed    (Why?)

Single Source      Undirected Graph

Negative-weight edges allowed      (Why?)

Simple path      (Why?)

Single Source      Undirected Graph

Negative-weight edges allowed    (Why?)

Simple path                      (Why?)

No negative-weight cycles    (o.w., NP-hard)



Single Source      Undirected Graph

Negative-weight edges allowed    (Why?)

Simple path                      (Why?)

No negative-weight cycles    (o.w., NP-hard)

Single-source  $s \rightsquigarrow$  Single-target  $t$

## Shortest Path Algorithms

Luis Goddyn, Math 408

Given an edge weighted graph  $(G, d)$ ,  $d : E(G) \rightarrow \mathbb{Q}$  and two vertices  $s, t \in V(G)$ , the *Shortest Path Problem* is to find an  $s, t$ -path  $P$  whose total weight is as small as possible. Here,  $G$  may be either directed or undirected. A path in a graph is a sequence  $v_0 e_1 v_1, \dots, v_k$  of vertices and edges such that no vertex or edge appears twice, and  $e_i$  joins  $v_{i-1}$  to  $v_i$ . If  $G$  is directed, then  $e_i$  should be oriented from  $v_{i-1}$  to  $v_i$ .

## Minimum-weight Perfect Matching

## Shortest Path Algorithms

Luis Goddyn, Math 408

Given an edge weighted graph  $(G, d)$ ,  $d : E(G) \rightarrow \mathbb{Q}$  and two vertices  $s, t \in V(G)$ , the *Shortest Path Problem* is to find an  $s, t$ -path  $P$  whose total weight is as small as possible. Here,  $G$  may be either directed or undirected. A path in a graph is a sequence  $v_0 e_1 v_1, \dots, v_k$  of vertices and edges such that no vertex or edge appears twice, and  $e_i$  joins  $v_{i-1}$  to  $v_i$ . If  $G$  is directed, then  $e_i$  should be oriented from  $v_{i-1}$  to  $v_i$ .

## Minimum-weight Perfect Matching

We leave it to the reader to  $\dots$

## Shortest Path Algorithms

Luis Goddyn, Math 408

Given an edge weighted graph  $(G, d)$ ,  $d : E(G) \rightarrow \mathbb{Q}$  and two vertices  $s, t \in V(G)$ , the *Shortest Path Problem* is to find an  $s, t$ -path  $P$  whose total weight is as small as possible. Here,  $G$  may be either directed or undirected. A path in a graph is a sequence  $v_0 e_1 v_1, \dots, v_k$  of vertices and edges such that no vertex or edge appears twice, and  $e_i$  joins  $v_{i-1}$  to  $v_i$ . If  $G$  is directed, then  $e_i$  should be oriented from  $v_{i-1}$  to  $v_i$ .

## Minimum-weight Perfect Matching

We leave it to the reader to  $\dots$

It is easy to check that  $\dots$

## Shortest Path Algorithms

Luis Goddyn, Math 408

Given an edge weighted graph  $(G, d)$ ,  $d : E(G) \rightarrow \mathbb{Q}$  and two vertices  $s, t \in V(G)$ , the *Shortest Path Problem* is to find an  $s, t$ -path  $P$  whose total weight is as small as possible. Here,  $G$  may be either directed or undirected. A path in a graph is a sequence  $v_0 e_1 v_1 \dots v_k$  of vertices and edges such that no vertex or edge appears twice, and  $e_i$  joins  $v_{i-1}$  to  $v_i$ . If  $G$  is directed, then  $e_i$  should be oriented from  $v_{i-1}$  to  $v_i$ .

## Minimum-weight Perfect Matching

We leave it to the reader to  $\dots$

It is easy to check that  $\dots$

Why? This will be a homework question.

## Shortest Path Algorithms

Luis Goddyn, Math 408

Given an edge weighted graph  $(G, d)$ ,  $d : E(G) \rightarrow \mathbb{Q}$  and two vertices  $s, t \in V(G)$ , the *Shortest Path Problem* is to find an  $s, t$ -path  $P$  whose total weight is as small as possible. Here,  $G$  may be either directed or undirected. A path in a graph is a sequence  $v_0 e_1 v_1 \dots v_k$  of vertices and edges such that no vertex or edge appears twice, and  $e_i$  joins  $v_{i-1}$  to  $v_i$ . If  $G$  is directed, then  $e_i$  should be oriented from  $v_{i-1}$  to  $v_i$ .

## Minimum-weight Perfect Matching

We leave it to the reader to  $\dots$

It is easy to check that  $\dots$

Why? This will be a homework question.

And Errors.

INTERESTED?  
*let's talk.*



## Robert W. Floyd (1936–2001)

*For having a clear influence on **methodologies** for the creation of efficient and reliable software, and for helping to **found** the following important subfields of computer science:*





## Robert W. Floyd (1936–2001)

*For having a clear influence on **methodologies** for the creation of efficient and reliable software, and for helping to **found** the following important subfields of computer science:*

*the theory of parsing, the semantics of programming languages, automatic program verification, automatic program synthesis, and analysis of algorithms*

— *Turing Award*, 1978

$$d_{ij}^{(k)} :$$


the weight of a shortest path from  $i$  to  $j$   
for which all *intermediate* vertices are in  $\{1, 2, \dots, k\}$

$$d_{ij}^{(k)} :$$

the weight of a shortest path from  $i$  to  $j$   
for which all *intermediate* vertices are in  $\{1, 2, \dots, k\}$


$$D^{(n)} \triangleq \left( d_{ij}^{(n)} \right)$$

$k \in \text{SP}_{ij}^{(k)}?$




$$d_{ij}^{(k)} = \begin{cases} w_{ij} & k = 0 \\ \min \left\{ d_{ij}^{(k-1)}, \underbrace{d_{ik}^{(k-1)} + d_{kj}^{(k-1)}} \right\} & k \geq 1 \end{cases}$$

$k \in \text{SP}_{ij}^{(k)}?$



$$d_{ij}^{(k)} = \begin{cases} w_{ij} & k = 0 \\ \min \left\{ d_{ij}^{(k-1)}, \underbrace{d_{ik}^{(k-1)} + d_{kj}^{(k-1)}}_{\text{why?}} \right\} & k \geq 1 \end{cases}$$

$k \in \text{SP}_{ij}^{(k)}?$



$$d_{ij}^{(k)} = \begin{cases} w_{ij} & k = 0 \\ \min \left\{ d_{ij}^{(k-1)}, \underbrace{d_{ik}^{(k-1)} + d_{kj}^{(k-1)}}_{\text{why?}} \right\} & k \geq 1 \end{cases}$$

$\dots$ , but we assume that there are **no** negative-weight cycles.

— Section 25.2 of CLRS

---

```
1: procedure FLOYD-WARSHALL( $W$ )
2:    $D^{(0)} = W$ 
3:   for  $k \leftarrow 1$  to  $n$  do
4:      $D^{(k)} \triangleq \left( d_{ij}^{(k)} \right) \leftarrow$  a new  $n \times n$  matrix
5:     for  $i \leftarrow 1$  to  $n$  do
6:       for  $j \leftarrow 1$  to  $n$  do
7:          $d_{ij}^{(k)} = \min \left\{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \right\}$ 
8:   return  $D^{(n)}$ 
```

---

---

```
1: procedure FLOYD-WARSHALL( $W$ )
2:    $D^{(0)} = W$ 
3:   for  $k \leftarrow 1$  to  $n$  do
4:      $D^{(k)} \triangleq \left(d_{ij}^{(k)}\right) \leftarrow$  a new  $n \times n$  matrix
5:     for  $i \leftarrow 1$  to  $n$  do
6:       for  $j \leftarrow 1$  to  $n$  do
7:          $d_{ij}^{(k)} = \min \left\{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \right\}$ 
8:   return  $D^{(n)}$ 
```

---

Space :  $\Theta(n^3)$



---

```

1: procedure FLOYD-WARSHALL( $W$ )
2:    $D^{(0)} = W$ 
3:   for  $k \leftarrow 1$  to  $n$  do
4:      $D^{(k)} \triangleq \left( d_{ij}^{(k)} \right) \leftarrow$  a new  $n \times n$  matrix
5:     for  $i \leftarrow 1$  to  $n$  do
6:       for  $j \leftarrow 1$  to  $n$  do
7:          $d_{ij}^{(k)} = \min \left\{ d_{ij}^{(k-1)}, d_{ik}^{(k-1)} + d_{kj}^{(k-1)} \right\}$ 
8:   return  $D^{(n)}$ 

```

---

Space :  $\Theta(n^3) \implies \Theta(n^2)$

## FLOYD-WARSHALL Made Simple (Problem 25.2-4)

---

```
1: procedure FLOYD-WARSHALL-SIMPLIFIED( $W$ )
2:    $D = W$ 
3:   for  $k \leftarrow 1$  to  $n$  do
4:     for  $i \leftarrow 1$  to  $n$  do
5:       for  $j \leftarrow 1$  to  $n$  do
6:          $d_{ij} = \min \{d_{ij}, d_{ik} + d_{kj}\}$ 
7:   return  $D$ 
```

---

## FLOYD-WARSHALL Made Simple (Problem 25.2-4)

---

```
1: procedure FLOYD-WARSHALL-SIMPLIFIED( $W$ )
2:    $D = W$ 
3:   for  $k \leftarrow 1$  to  $n$  do
4:     for  $i \leftarrow 1$  to  $n$  do
5:       for  $j \leftarrow 1$  to  $n$  do
6:          $d_{ij} = \min \{d_{ij}, d_{ik} + d_{kj}\}$ 
7:   return  $D$ 
```

---

“Decrease” does no harm to the correctness.

## Negative-weight Cycle Detection (Problem 25.2-6)

To detect negative-weight cycle (NC) using FLOYD-WARSHALL.

## Negative-weight Cycle Detection (Problem 25.2-6)

To detect negative-weight cycle (NC) using FLOYD-WARSHALL.

$$\exists i : d_{ii}^{(n)} < 0$$

## Negative-weight Cycle Detection (Problem 25.2-6)

To detect negative-weight cycle (NC) using FLOYD-WARSHALL.

$$\exists i : d_{ii}^{(n)} < 0$$

Proof.

## Negative-weight Cycle Detection (Problem 25.2-6)

To detect negative-weight cycle (NC) using FLOYD-WARSHALL.

$$\exists i : d_{ii}^{(n)} < 0$$

Proof.

**The  
proof is  
trivial.**


## Negative-weight Cycle Detection (Problem 25.2-6)

To detect negative-weight cycle (NC) using FLOYD-WARSHALL.

$$\exists i : d_{ii}^{(n)} < 0$$

Proof.

**The  
proof is  
trivial.**






## Negative-weight Cycle Detection (Problem 25.2-6)

To detect negative-weight cycle (NC) using FLOYD-WARSHALL.

$$\exists i : d_{ii}^{(n)} < 0$$

Proof.

**The  
proof is  
trivial.**



$$\exists i : d_{ii}^{(n)} < 0 \iff \exists \text{ NC} \subseteq G$$



$$\exists i : d_{ii}^{(n)} < 0 \iff \exists \text{NC} \subseteq G$$

Proof.

$$\exists i : d_{ii}^{(n)} < 0 \iff \exists \text{ NC} \subseteq G$$


Proof.

“  $\implies$  ”

$$\exists i : d_{ii}^{(n)} < 0 \iff \exists \text{ NC} \subseteq G$$

Proof.

“  $\implies$  ”




A Negative Cycle

$$\exists i : d_{ii}^{(n)} < 0 \iff \exists \text{ NC} \subseteq G$$

Proof.

“  $\Leftarrow$  ”

“  $\Rightarrow$  ”




A Negative Cycle

$$\exists i : d_{ii}^{(n)} < 0 \iff \exists \text{ NC} \subseteq G$$


Proof.

“  $\implies$  ”



A Negative Cycle

“  $\impliedby$  ”




A Simple Negative Cycle

$k : \max \#$

$$\exists i : d_{ii}^{(n)} < 0 \iff \exists \text{ NC} \subseteq G$$


Proof.

“  $\implies$  ”



A Negative Cycle

“  $\impliedby$  ”



A Simple Negative Cycle

$k : \max \#$

$$d_{ii}^{(k)} < 0$$



$l_{ij}^{(m)}$  : the length of a shortest path from  $i$  to  $j$  consisting of  $\leq m$  edges



$l_{ij}^{(m)}$  : the length of a shortest path from  $i$  to  $j$  consisting of  $\leq m$  edges

$$l_{ij}^{(m)} = \min_{1 \leq k \leq n} \{l_{ik}^{(m-1)} + w_{kj}\}, \quad m \geq 2$$

$l_{ij}^{(m)}$  : the length of a shortest path from  $i$  to  $j$  consisting of  $\leq m$  edges

$$l_{ij}^{(m)} = \min_{1 \leq k \leq n} \{l_{ik}^{(m-1)} + w_{kj}\}, \quad m \geq 2$$

$$l_{ij}^{(1)} = w_{ij}$$

$l_{ij}^{(m)}$  : the length of a shortest path from  $i$  to  $j$  consisting of  $\leq m$  edges

$$l_{ij}^{(m)} = \min_{1 \leq k \leq n} \{l_{ik}^{(m-1)} + w_{kj}\}, \quad m \geq 2$$

$$l_{ij}^{(1)} = w_{ij}$$

$$L^{(n-1)} \triangleq (l_{ij}^{(n-1)})$$

$l_{ij}^{(m)}$  : the length of a shortest path from  $i$  to  $j$  consisting of  $\leq m$  edges

$$l_{ij}^{(m)} = \min_{1 \leq k \leq n} \{l_{ik}^{(m-1)} + w_{kj}\}, \quad m \geq 2$$

$$l_{ij}^{(1)} = w_{ij}$$

$$L^{(n-1)} \triangleq \left( l_{ij}^{(n-1)} \right)$$

$$L^{(n-1)} = W^{n-1} \triangleq \underbrace{((W \cdot W) \cdot \dots)}_{n-1} \cdot W$$

## Associative EXTEND-SHORTEST-PATHS

Show that matrix multiplication defined by EXTEND-SHORTEST-PATHS is associative.

## Associative EXTEND-SHORTEST-PATHS

Show that matrix multiplication defined by EXTEND-SHORTEST-PATHS is associative.

$$(W^a W^b) W^c = W^a (W^b W^c)$$

## Associative EXTEND-SHORTEST-PATHS

Show that matrix multiplication defined by EXTEND-SHORTEST-PATHS is associative.

$$(W^a W^b) W^c = W^a (W^b W^c)$$

$$(i, j)$$

$$\begin{aligned} & \min_{1 \leq k \leq n} \left( (W^a W^b)_{ik} + W^c_{kj} \right) & \min_{1 \leq k \leq n} \left( W^a_{ik} + (W^b W^c)_{kj} \right) \\ = & \min_{1 \leq k \leq n} \left( \min_{1 \leq q \leq n} (W^a_{iq} + W^b_{qk}) + W^c_{kj} \right) & = \min_{1 \leq k \leq n} \left( W^a_{ik} + \min_{1 \leq q \leq n} (W^b_{kq} + W^c_{qj}) \right) \end{aligned}$$

## Associative EXTEND-SHORTEST-PATHS

Show that matrix multiplication defined by EXTEND-SHORTEST-PATHS is associative.

$$(W^a W^b) W^c = W^a (W^b W^c)$$

$(i, j)$

$$\begin{aligned} & \min_{1 \leq k \leq n} \left( (W^a W^b)_{ik} + W^c_{kj} \right) & \min_{1 \leq k \leq n} \left( W^a_{ik} + (W^b W^c)_{kj} \right) \\ = & \min_{1 \leq k \leq n} \left( \min_{1 \leq q \leq n} (W^a_{iq} + W^b_{qk}) + W^c_{kj} \right) & = \min_{1 \leq k \leq n} \left( W^a_{ik} + \min_{1 \leq q \leq n} (W^b_{kq} + W^c_{qj}) \right) \end{aligned}$$

$Q$  : Why do we care about this?



## Associative EXTEND-SHORTEST-PATHS

Show that matrix multiplication defined by EXTEND-SHORTEST-PATHS is associative.

$$(W^a W^b) W^c = W^a (W^b W^c)$$

$$(i, j)$$

$$\begin{aligned} & \min_{1 \leq k \leq n} \left( (W^a W^b)_{ik} + W^c_{kj} \right) & \min_{1 \leq k \leq n} \left( W^a_{ik} + (W^b W^c)_{kj} \right) \\ = & \min_{1 \leq k \leq n} \left( \min_{1 \leq q \leq n} (W^a_{iq} + W^b_{qk}) + W^c_{kj} \right) & = \min_{1 \leq k \leq n} \left( W^a_{ik} + \min_{1 \leq q \leq n} (W^b_{kq} + W^c_{qj}) \right) \end{aligned}$$

*Q* : Why do we care about this?

*A* : Repeated squaring.

## Associative EXTEND-SHORTEST-PATHS

Show that matrix multiplication defined by EXTEND-SHORTEST-PATHS is associative.

$$(W^a W^b) W^c = W^a (W^b W^c) \quad \text{vs.} \quad (WW)W = W(WW)$$

$(i, j)$

$$\begin{aligned} & \min_{1 \leq k \leq n} \left( (W^a W^b)_{ik} + W^c_{kj} \right) & \min_{1 \leq k \leq n} \left( W^a_{ik} + (W^b W^c)_{kj} \right) \\ = & \min_{1 \leq k \leq n} \left( \min_{1 \leq q \leq n} (W^a_{iq} + W^b_{qk}) + W^c_{kj} \right) & = \min_{1 \leq k \leq n} \left( W^a_{ik} + \min_{1 \leq q \leq n} (W^b_{kq} + W^c_{qj}) \right) \end{aligned}$$

$Q$  : Why do we care about this?

$A$  : Repeated squaring.

SSSP from  $s$

## SSSP from $s$

$l_v^{(m)}$  : the length of a shortest path  $s \rightsquigarrow v$  consisting of  $\leq m$  edges

## SSSP from $s$

$l_v^{(m)}$  : the length of a shortest path  $s \rightsquigarrow v$  consisting of  $\leq m$  edges

$$l_v^{(m)} = \min_{1 \leq u \leq n} \left\{ l_u^{(m-1)} + w_{uv} \right\}, \quad m \geq 2$$

SSSP from  $s$

$l_v^{(m)}$  : the length of a shortest path  $s \rightsquigarrow v$  consisting of  $\leq m$  edges

$$l_v^{(m)} = \min_{1 \leq u \leq n} \left\{ l_u^{(m-1)} + w_{uv} \right\}, \quad m \geq 2$$

$$l_v^{(1)} = w_{sv}$$

SSSP from  $s$

$l_v^{(m)}$  : the length of a shortest path  $s \rightsquigarrow v$  consisting of  $\leq m$  edges

$$l_v^{(m)} = \min_{1 \leq u \leq n} \left\{ l_u^{(m-1)} + w_{uv} \right\}, \quad m \geq 2$$

$$l_v^{(1)} = w_{sv}$$

$$L^{(n-1)} \triangleq \left( l_v^{(n-1)} \right)$$

## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

To express SSSP as a **product** of matrices and a vector.



## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

To express SSSP as a **product** of matrices and a vector.

$$L^{(1)} = \langle w_{sv} \rangle_{v \in V}$$

$$L^{(2)} = L^{(1)} \cdot W$$

$$\dots = \dots$$

$$L^{(n-1)} = \left( \left( \left( L^{(1)} \cdot W \right) \cdot W \right) \cdot \dots \cdot W \right)$$

## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

What is the **relationship** between it and the Bellman-Ford algorithm?

## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

What is the **relationship** between it and the Bellman-Ford algorithm?

$$l_v^{(m)} = \min_{1 \leq u \leq n} \left\{ l_u^{(i-1)} + w_{uv} \right\}, \quad m \geq 2$$

## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

What is the **relationship** between it and the Bellman-Ford algorithm?

$$l_v^{(m)} = \min_{1 \leq u \leq n} \left\{ l_u^{(i-1)} + w_{uv} \right\}, \quad m \geq 2$$

$$d_v^{(i)} = \min_{u \rightarrow v} \left\{ d_u^{(i-1)} + w_{uv} \right\}, \quad i \geq 1$$

## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

What is the **relationship** between it and the Bellman-Ford algorithm?

$$l_v^{(m)} = \min_{1 \leq u \leq n} \{ l_u^{(i-1)} + w_{uv} \}, \quad m \geq 2$$

$$d_v^{(i)} = \min_{u \rightarrow v} \{ d_u^{(i-1)} + w_{uv} \}, \quad i \geq 1$$

$$d_v^{(0)} = \begin{cases} 0 & v = s \\ \infty & v \neq s \end{cases}$$

---

```
1: procedure BELLMAN-FORD-DP( $G, w, s$ )
2:    $d[0, s] \leftarrow 0$ 
3:   for  $(v \neq s) \in V$  do
4:      $d[0, v] \leftarrow \infty$ 
5:   for  $i \leftarrow 1$  to  $|V| - 1$  do
6:     for  $v \in V$  do
7:        $d[i, v] = \infty$ 
8:       for  $(u, v) \in E$  do
9:         if  $d[i, v] > d[i - 1, u] + w(u, v)$  then ▷
10:           $d[i, v] = d[i - 1, u] + w(u, v)$ 
```

---

---

```
1: procedure BELLMAN-FORD-DP( $G, w, s$ )
2:    $d[0, s] \leftarrow 0$ 
3:   for  $(v \neq s) \in V$  do
4:      $d[0, v] \leftarrow \infty$ 
5:   for  $i \leftarrow 1$  to  $|V| - 1$  do
6:     for  $v \in V$  do
7:        $d[i, v] = \infty$ 
8:       for  $(u, v) \in E$  do
9:         if  $d[i, v] > d[i - 1, u] + w(u, v)$  then           ▷ Simplify?
10:           $d[i, v] = d[i - 1, u] + w(u, v)$ 
```

---

---

```
1: procedure BELLMAN-FORD-DP-SIMPLIFIED( $G, w, s$ )
2:    $d[s] \leftarrow 0$ 
3:   for  $(v \neq s) \in V$  do
4:      $d[v] \leftarrow \infty$ 
5:   for  $i \leftarrow 1$  to  $|V| - 1$  do
6:     for  $v \in V$  do
7:       for  $(u, v) \in E$  do
8:         if  $d[v] > d[u] + w(u, v)$  then ▷
9:            $d[v] = d[u] + w(u, v)$ 
```

---



---

```
1: procedure BELLMAN-FORD-DP-SIMPLIFIED( $G, w, s$ )
2:    $d[s] \leftarrow 0$ 
3:   for  $(v \neq s) \in V$  do
4:      $d[v] \leftarrow \infty$ 
5:   for  $i \leftarrow 1$  to  $|V| - 1$  do
6:     for  $v \in V$  do
7:       for  $(u, v) \in E$  do
8:         if  $d[v] > d[u] + w(u, v)$  then                                 $\triangleright$  Relax!
9:            $d[v] = d[u] + w(u, v)$ 
```

---

---

```
1: procedure BELLMAN-FORD-DP-SIMPLIFIED( $G, w, s$ )
2:    $d[s] \leftarrow 0$ 
3:   for  $(v \neq s) \in V$  do
4:      $d[v] \leftarrow \infty$ 
5:   for  $i \leftarrow 1$  to  $|V| - 1$  do
6:     for  $v \in V$  do
7:       for  $(u, v) \in E$  do
8:         if  $d[v] > d[u] + w(u, v)$  then                                 $\triangleright$  Relax!
9:            $d[v] = d[u] + w(u, v)$ 
```

---

---

```
1: procedure BELLMAN-FORD-WITHOUT-NE( $G, w, s$ )
2:   INIT-SINGLE-SOURCE( $G, s$ )
3:   for  $i \leftarrow 1$  to  $|V| - 1$  do
4:     for  $(u, v) \in E$  do
5:       RELAX( $u, v, w$ )
```

---

Bellman-Ford:  $L \cdot W$

SLOW-ALL-PAIRS-SHORTEST-PATHS:  $W \cdot W$

Bellman-Ford:  $L \cdot W$

SLOW-ALL-PAIRS-SHORTEST-PATHS:  $W \cdot W$

SLOW-ALL-PAIRS-SHORTEST-PATHS is  $n$  instances of Bellman-Ford,  
one for each source.

## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

To express SSSP as a **product** of matrices and a vector.

$$L^{(n-1)} = \left( \left( \left( L^{(1)} \cdot W \right) \cdot W \right) \cdot \dots \cdot W \right)$$

## SSSP as a Product of Matrices and a Vector (Problem 25.1-5)

To express SSSP as a **product** of matrices and a vector.

$$L^{(n-1)} = \left( \left( \left( L^{(1)} \cdot W \right) \cdot W \right) \cdot \dots \cdot W \right)$$

*Q* : Associative? Repeated Squaring?

## Negative-weight Cycle Detection (Problem 25.1-9)

To detect negative-weight cycle (NC) using  
FASTER-ALL-PAIRS-SHORTEST-PATHS.

## Minimum-length Negative-weight Cycle (Problem 25.1-10)

To find the length of a minimum-length negative-weight cycle (NC).







Office 302

Mailbox: H016

hfwei@nju.edu.cn