

# 3-10 Traversability


## (Part I: Eulerian Graphs)

Hengfeng Wei


hfwei@nju.edu.cn

December 03, 2018





Leonhard Euler (1707 – 1783)





Leonhard Euler (1707 – 1783)

Graph Theory

Topology







## Theorem (Leonhard Euler 1735)

*A connected graph  $G$  is **Eulerian** if and only if the degree of each vertex of  $G$  is even.*

## Theorem (Leonhard Euler 1735)

*A connected graph  $G$  is **Eulerian** if and only if the degree of each vertex of  $G$  is even.*

“ $\Leftarrow$ ” (Carl Hierholzer 1873);



## Theorem (Leonhard Euler 1735)

*A connected graph  $G$  is **Eulerian** if and only if the degree of each vertex of  $G$  is even.*

“ $\Leftarrow$ ” (Carl Hierholzer 1873); (Proofs on PPT from Tao & on CZ).

## Theorem (Leonhard Euler 1735)

A connected graph  $G$  is *Eulerian* if and only if the degree of each vertex of  $G$  is even.

“ $\Leftarrow$ ” (Carl Hierholzer 1873); (Proofs on PPT from Tao & on CZ).

By strong mathematical induction on the number  $m$  of edges of  $G$ .

## Theorem (Leonhard Euler 1735)

A connected graph  $G$  is *Eulerian* if and only if the degree of each vertex of  $G$  is even.

“ $\Leftarrow$ ” (Carl Hierholzer 1873); (Proofs on PPT from Tao & on CZ).

By strong mathematical induction on the number  $m$  of edges of  $G$ .

Inductive Step :  $m = k + 1$

## Theorem (Leonhard Euler 1735)

A connected graph  $G$  is *Eulerian* if and only if the degree of each vertex of  $G$  is even.

“ $\Leftarrow$ ” (Carl Hierholzer 1873); (Proofs on PPT from Tao & on CZ).

By strong mathematical induction on the number  $m$  of edges of  $G$ .

Inductive Step :  $m = k + 1$

Let  $C = u \sim v$  be any trail in  $G$  of the maximum length.

## Theorem (Leonhard Euler 1735)

A connected graph  $G$  is *Eulerian* if and only if the degree of each vertex of  $G$  is even.

“ $\Leftarrow$ ” (Carl Hierholzer 1873); (Proofs on PPT from Tao & on CZ).

By strong mathematical induction on the number  $m$  of edges of  $G$ .

Inductive Step :  $m = k + 1$

Let  $C = u \sim v$  be any trail in  $G$  of the maximum length.

$$v = u$$

## Theorem (Leonhard Euler 1735)

A connected graph  $G$  is *Eulerian* if and only if the degree of each vertex of  $G$  is even.

“ $\Leftarrow$ ” (Carl Hierholzer 1873); (Proofs on PPT from Tao & on CZ).

By strong mathematical induction on the number  $m$  of edges of  $G$ .

Inductive Step :  $m = k + 1$

Let  $C = u \sim v$  be any trail in  $G$  of the maximum length.

$$v = u \implies C = u \sim u$$

## Theorem (Leonhard Euler 1735)

A connected graph  $G$  is *Eulerian* if and only if the degree of each vertex of  $G$  is even.

“ $\Leftarrow$ ” (Carl Hierholzer 1873); (Proofs on PPT from Tao & on CZ).

By strong mathematical induction on the number  $m$  of edges of  $G$ .

Inductive Step :  $m = k + 1$


Let  $C = u \sim v$  be any trail in  $G$  of the maximum length.

$$v = u \implies C = u \sim u$$

$$H = G - E(C) = \bigcup H_i$$




$$H = G - E(C) = \bigcup H_i$$






$$H = G - E(C) = \bigcup H_i$$

- (I)  $\forall v \in H : \deg(v)$  is even
- (II)  $\forall i : |E(H_i)| < m$



$$H = G - E(C) = \bigcup H_i$$

- (I)  $\forall v \in H : \deg(v)$  is even
- (II)  $\forall i : |E(H_i)| < m$




By I.H., each  $H_i$  has an Eulerian circuit  $C_i$ .

$$H = G - E(C) = \bigcup H_i$$

(I)  $\forall v \in H : \deg(v)$  is even

(II)  $\forall i : |E(H_i)| < m$




By I.H., each  $H_i$  has an Eulerian circuit  $C_i$ .

$$\forall i : V(H_i) \cap V(C) \neq \emptyset$$

$$H = G - E(C) = \bigcup H_i$$

- (I)  $\forall v \in H : \deg(v)$  is even
- (II)  $\forall i : |E(H_i)| < m$



By I.H., each  $H_i$  has an Eulerian circuit  $C_i$ .

$$\forall i : V(H_i) \cap V(C) \neq \emptyset$$

Glue together each  $C_i$  with  $C$  to get an Eulerian circuit of  $G$ .


---

```

1: procedure HIERHOLZER( $G$ )
2:    $u \in V(G)$ 
3:    $C \leftarrow$  any circuit  $u \sim u$  in  $G$ 
4:   while  $\exists v \in C : \deg(v) > 0$  do
5:      $H \leftarrow G - E(C)$ 
6:      $v \leftarrow$  any vertex in  $V(C)$  such that  $\deg(v) > 0$ 
7:      $C' \leftarrow$  any circuit  $v \sim v$  in  $H$ 
8:      $C \leftarrow C \otimes C'$  ▷ Glue  $C' = v \sim v$  with  $C$  via  $v$ 
9:   return  $C$ 

```

---




---

```

1: procedure HIERHOLZER( $G$ )
2:    $u \in V(G)$ 
3:    $C \leftarrow$  any circuit  $u \sim u$  in  $G$ 
4:   while  $\exists v \in C : \deg(v) > 0$  do
5:      $H \leftarrow G - E(C)$ 
6:      $v \leftarrow$  any vertex in  $V(C)$  such that  $\deg(v) > 0$ 
7:      $C' \leftarrow$  any circuit  $v \sim v$  in  $H$ 
8:      $C \leftarrow C \otimes C'$  ▷ Glue  $C' = v \sim v$  with  $C$  via  $v$ 
9:   return  $C$ 

```

---



$Q$  : Time Complexity?


---

```

1: procedure HIERHOLZER( $G$ )
2:    $u \in V(G)$ 
3:    $C \leftarrow$  any circuit  $u \sim u$  in  $G$ 
4:   while  $\exists v \in C : \deg(v) > 0$  do
5:      $H \leftarrow G - E(C)$ 
6:      $v \leftarrow$  any vertex in  $V(C)$  such that  $\deg(v) > 0$ 
7:      $C' \leftarrow$  any circuit  $v \sim v$  in  $H$ 
8:      $C \leftarrow C \otimes C'$  ▷ Glue  $C' = v \sim v$  with  $C$  via  $v$ 
9:   return  $C$ 

```

---



$Q$  : Time Complexity?

$Q$  : Data Structures?


---

```

1: procedure HIERHOLZER( $G$ )
2:    $u \in V(G)$ 
3:    $C \leftarrow$  any circuit  $u \sim u$  in  $G$ 
4:   while  $\exists v \in C : \deg(v) > 0$  do
5:      $H \leftarrow G - E(C)$ 
6:      $v \leftarrow$  any vertex in  $V(C)$  such that  $\deg(v) > 0$ 
7:      $C' \leftarrow$  any circuit  $v \sim v$  in  $H$ 
8:      $C \leftarrow C \otimes C'$  ▷ Glue  $C' = v \sim v$  with  $C$  via  $v$ 
9:   return  $C$ 

```

---



$Q$  : Time Complexity?

$Q$  : Data Structures?

$O(m)$  : Using doubly linked list



## Fleury's Algorithm (1883)

(I)  $v_0 \in V(G)$ ;  $P_0 = v_0$

(II) Suppose  $C_i = v_0 e_1 v_1 e_2 \cdots e_i v_i$ .

Choose  $e_{i+1}$  from  $E(G) - \{e_1, e_2, \cdots, e_i\}$  as follows:

- (a)  $e_{i+1}$  is incident with  $v_i$
- (b) Unless there is no alternative,  $e_{i+1}$  is not a bridge of  $G - \{e_1, e_2, \cdots, e_i\}$

(III) Stop when step (II) can no longer be implemented

## Fleury's Algorithm (1883)

(I)  $v_0 \in V(G)$ ;  $P_0 = v_0$

(II) Suppose  $C_i = v_0 e_1 v_1 e_2 \cdots e_i v_i$ .

Choose  $e_{i+1}$  from  $E(G) - \{e_1, e_2, \cdots, e_i\}$  as follows:

- (a)  $e_{i+1}$  is incident with  $v_i$
- (b) Unless there is no alternative,  $e_{i+1}$  is not a bridge of  $G - \{e_1, e_2, \cdots, e_i\}$

(III) Stop when step (II) can no longer be implemented



## Fleury's Algorithm (1883)

(I)  $v_0 \in V(G)$ ;  $P_0 = v_0$

(II) Suppose  $C_i = v_0 e_1 v_1 e_2 \cdots e_i v_i$ .

Choose  $e_{i+1}$  from  $E(G) - \{e_1, e_2, \cdots, e_i\}$  as follows:

- (a)  $e_{i+1}$  is incident with  $v_i$
- (b) Unless there is no alternative,  $e_{i+1}$  is not a bridge of  $G - \{e_1, e_2, \cdots, e_i\}$

(III) Stop when step (II) can no longer be implemented



## Fleury's Algorithm (1883)

(I)  $v_0 \in V(G)$ ;  $P_0 = v_0$

(II) Suppose  $C_i = v_0 e_1 v_1 e_2 \cdots e_i v_i$ .

Choose  $e_{i+1}$  from  $E(G) - \{e_1, e_2, \cdots, e_i\}$  as follows:

- (a)  $e_{i+1}$  is incident with  $v_i$
- (b) Unless there is no alternative,  $e_{i+1}$  is not a bridge of  $G - \{e_1, e_2, \cdots, e_i\}$

(III) Stop when step (II) can no longer be implemented



## Fleury's Algorithm (1883)

(I)  $v_0 \in V(G)$ ;  $P_0 = v_0$

(II) Suppose  $C_i = v_0 e_1 v_1 e_2 \cdots e_i v_i$ .

Choose  $e_{i+1}$  from  $E(G) - \{e_1, e_2, \cdots, e_i\}$  as follows:

(a)  $e_{i+1}$  is incident with  $v_i$

(b) Unless there is no alternative,  $e_{i+1}$  is not a bridge of  $G - \{e_1, e_2, \cdots, e_i\}$

(III) Stop when step (II) can no longer be implemented



# PROOF

## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

**By Contradiction.**


Suppose that  $v_i$  is incident with  $\geq 2$  bridges in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .

## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

By Contradiction.

Suppose that  $v_i$  is incident with  $\geq 2$  bridges in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .






## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

By Contradiction.

Suppose that  $v_i$  is incident with  $\geq 2$  bridges in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .




$$\exists x' \in C_x : \deg(x) \text{ is odd}$$

## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

By Contradiction.

Suppose that  $v_i$  is incident with  $\geq 2$  bridges in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .



$\exists x' \in C_x : \deg(x) \text{ is odd}$


$\exists y' \in C_y : \deg(y) \text{ is odd}$

## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

By Contradiction.

Suppose that  $v_i$  is incident with  $\geq 2$  bridges in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .



$$\exists x' \in C_x : \deg(x) \text{ is odd}$$

$$\exists y' \in C_y : \deg(y) \text{ is odd}$$


We have found 2 odd vertices.

## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

By Contradiction.

Suppose that  $v_i$  is incident with  $\geq 2$  bridges in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .



$\exists x' \in C_x : \deg(x) \text{ is odd}$

$\exists y' \in C_y : \deg(y) \text{ is odd}$

We have found 2 odd vertices.


*Q : What is the contradiction?*

## Theorem (Bridges in Fleury's Algorithm)

*At any stage,  $v_i$  is incident with  $\leq 1$  bridge in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .*

By Contradiction.

Suppose that  $v_i$  is incident with  $\geq 2$  bridges in  $E(G) - \{e_1, e_2, \dots, e_i\}$ .



$\exists x' \in C_x : \deg(x) \text{ is odd}$

$\exists y' \in C_y : \deg(y) \text{ is odd}$

We have found 2 odd vertices.

*Q : What is the contradiction?*

Is  $\deg(v_i)$  odd or even?

---

1: **procedure** FLEURY( $G$ )

2:      $v_0 \in V(G)$

3:      $C \leftarrow v_0$

4:      $i \leftarrow 0$ ,  $V_0 \leftarrow V(G)$ ,  $E_0 \leftarrow E(G)$

▷ Choose any starting vertex

▷ Keep track of the circuit

▷ Stop otherwise

---

```

1: procedure FLEURY( $G$ )
2:    $v_0 \in V(G)$                                 ▷ Choose any starting vertex
3:    $C \leftarrow v_0$                                ▷ Keep track of the circuit
4:    $i \leftarrow 0$ ,  $V_0 \leftarrow V(G)$ ,  $E_0 \leftarrow E(G)$ 

5:   while  $\deg(v_i) > 0$  in  $E_i$  do                ▷ Stop otherwise
6:     if  $\deg(v_i) = 1$  in  $E_i$  then                ▷ No alternative: go the bridge
7:        $e_{i+1} \triangleq v_i v_{i+1}$ 
8:     else                                          ▷ Delete the isolated vertex  $v_i$ 
9:        $\triangleright$  Have alternatives: don't go the bridge
10:      Choose  $e_{i+1} \triangleq v_i v_{i+1}$  that is not a bridge of  $(V_i, E_i)$ 
11:       $\triangleright$  No isolated vertex produced

15:  return  $C$ 

```

---



---

```

1: procedure FLEURY( $G$ )
2:    $v_0 \in V(G)$                                 ▷ Choose any starting vertex
3:    $C \leftarrow v_0$                                 ▷ Keep track of the circuit
4:    $i \leftarrow 0$ ,  $V_0 \leftarrow V(G)$ ,  $E_0 \leftarrow E(G)$ 

5:   while  $\deg(v_i) > 0$  in  $E_i$  do                                ▷ Stop otherwise
6:     if  $\deg(v_i) = 1$  in  $E_i$  then                                ▷ No alternative: go the bridge
7:        $e_{i+1} \triangleq v_i v_{i+1}$ 
8:                                     ▷ Delete the isolated vertex  $v_i$ 
9:     else                                ▷ Have alternatives: don't go the bridge
10:      Choose  $e_{i+1} \triangleq v_i v_{i+1}$  that is not a bridge of  $(V_i, E_i)$ 
11:                                     ▷ No isolated vertex produced

12:       $C \leftarrow C e_{i+1} v_{i+1}$ 
13:       $E_{i+1} \leftarrow E_i - \{e_{i+1}\}$ 
14:       $i \leftarrow i + 1$ 

15:   return  $C$ 

```

---

---

```

1: procedure FLEURY( $G$ )
2:    $v_0 \in V(G)$                                 ▷ Choose any starting vertex
3:    $C \leftarrow v_0$                                 ▷ Keep track of the circuit
4:    $i \leftarrow 0, V_0 \leftarrow V(G), E_0 \leftarrow E(G)$ 

5:   while  $\deg(v_i) > 0$  in  $E_i$  do                                ▷ Stop otherwise
6:     if  $\deg(v_i) = 1$  in  $E_i$  then                                ▷ No alternative: go the bridge
7:        $e_{i+1} \triangleq v_i v_{i+1}$ 
8:        $V_{i+1} \leftarrow V_i - \{v_i\}$                                 ▷ Delete the isolated vertex  $v_i$ 
9:     else                                ▷ Have alternatives: don't go the bridge
10:      Choose  $e_{i+1} \triangleq v_i v_{i+1}$  that is not a bridge of  $(V_i, E_i)$ 
11:       $V_{i+1} \leftarrow V_i$                                 ▷ No isolated vertex produced

12:       $C \leftarrow C e_{i+1} v_{i+1}$ 
13:       $E_{i+1} \leftarrow E_i - \{e_{i+1}\}$ 
14:       $i \leftarrow i + 1$ 

15:   return  $C$ 

```

---

complexity





Office 302

Mailbox: H016

hfwei@nju.edu.cn