

3-5 Minimum Spanning Trees

Hengfeng Wei

hfwei@nju.edu.cn

October 22, 2018



[Problem: 4.8]

$\forall v \in V(G) : \deg(v) \geq 2 \implies G \text{ contains a cycle}$

Cut Property

$$G = (V, E, w)$$

Cut Property (I)

X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$

e : *A* lightest edge across $(S, V \setminus S)$

Cut Property (I)

X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$

e : *A* lightest edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of some MST T' of G .

Cut Property (I)

X : A part of some MST T of G

$(S, V \setminus S)$: A *cut* such that X does *not* cross $(S, V \setminus S)$


e : *A* lightest edge across $(S, V \setminus S)$

Then $X \cup \{e\}$ is a part of some MST T' of G .


Correctness of Prim's and Kruskal's algorithms.

By Exchange Argument.

By Exchange Argument.




By Exchange Argument.



$$T' = \underbrace{T + \{e\}}_{\text{if } e \notin T} - \{e'\}$$

By Exchange Argument.



$$T' = T + \underbrace{\{e\}}_{\text{if } e \notin T} - \{e'\}$$


“a” \rightarrow “the” \Rightarrow “some” \rightarrow “all”

Cut Property (II)

A cut $(S, V \setminus S)$

Let $e = (u, v)$ be **a** lightest edge across $(S, V \setminus S)$

\exists MST T of $G : e \in T$




Cut Property (II)

A cut $(S, V \setminus S)$

Let $e = (u, v)$ be **a** lightest edge across $(S, V \setminus S)$

\exists MST T of $G : e \in T$




$$T' = T + \underbrace{\{e\}}_{\text{if } e \notin T} - \{e'\}$$

Cut Property (II)

A cut $(S, V \setminus S)$

Let $e = (u, v)$ be **a** lightest edge across $(S, V \setminus S)$

\exists MST T of $G : e \in T$



$$T' = \underbrace{T + \{e\}}_{\text{if } e \notin T} - \{e'\}$$

“a” \rightarrow “the” \implies “ \exists ” \rightarrow “ \forall ”

Application of Cut Property [Problem: 10.15 (3)]

$e = (u, v) \in G$ is a lightest edge $\implies e \in \exists$ MST of G

Application of Cut Property [Problem: 10.15 (4)]

$e = (u, v) \in G$ is the unique lightest edge $\implies e \in \forall$ MST

Application of Cut Property [Problem: 10.15 (3)]

$e = (u, v) \in G$ is a lightest edge $\implies e \in \exists$ MST of G

$$(S = \{u\}, V \setminus S)$$

Application of Cut Property [Problem: 10.15 (4)]

$e = (u, v) \in G$ is the unique lightest edge $\implies e \in \forall$ MST

Wrong Divide&Conquer Algorithm for MST [Problem: 10.21]


$$(V_1, V_2) : \left| |V_1| - |V_2| \right| \leq 1$$

$$T_1 + T_2 + \{e\} : e \text{ is a lightest edge across } (V_1, V_2)$$

Wrong Divide&Conquer Algorithm for MST [Problem: 10.21]

$$(V_1, V_2) : \left| |V_1| - |V_2| \right| \leq 1$$

$T_1 + T_2 + \{e\} : e$ is a lightest edge across (V_1, V_2)




Cycle Property

Cycle Property [Problem: 10.19(b)]

- ▶ Let C be any cycle in G
- ▶ Let $e = (u, v)$ be **a** maximum-weight edge in C


Then \exists MST T of $G : e \notin T$.



Cycle Property [Problem: 10.19(b)]

- ▶ Let C be any cycle in G
- ▶ Let $e = (u, v)$ be **a** maximum-weight edge in C

Then \exists MST T of $G : e \notin T$.




$$T' = T - \underbrace{\{e\}}_{\text{if } e \in T} + \{e'\}$$

Cycle Property [Problem: 10.19(b)]

- ▶ Let C be any cycle in G
- ▶ Let $e = (u, v)$ be **a** maximum-weight edge in C

Then \exists MST T of $G : e \notin T$.



$$T' = T - \underbrace{\{e\}}_{\text{if } e \in T} + \{e'\}$$

“a” \rightarrow “the” \Rightarrow “ \exists ” \rightarrow “ \forall ”

Anti-Kruskal algorithm [[Problem: 10.19 \(c\)](#)]

Reverse-delete algorithm ([wiki](#); [clickable](#))

Anti-Kruskal algorithm [Problem: 10.19 (c)]

Reverse-delete algorithm ([wiki](#); [clickable](#))

$$O(m \log n (\log \log n)^3)$$

Anti-Kruskal algorithm [Problem: 10.19 (c)]

Reverse-delete algorithm ([wiki](#); [clickable](#))

$$O(m \log n (\log \log n)^3)$$

Proof.

Cycle Property

Anti-Kruskal algorithm [Problem: 10.19 (c)]

Reverse-delete algorithm ([wiki](#); [clickable](#))

$$O(m \log n (\log \log n)^3)$$

Proof.

Cycle Property

$$T \subseteq F \implies \exists T' : T' \subseteq F - \{e\}$$



Anti-Kruskal algorithm [Problem: 10.19 (c)]

Reverse-delete algorithm ([wiki](#); [clickable](#))

$$O(m \log n (\log \log n)^3)$$

Proof.

Cycle Property

$$T \subseteq F \implies \exists T' : T' \subseteq F - \{e\}$$



*“On the Shortest Spanning Subtree of a Graph
and the Traveling Salesman Problem”*

— **Kruskal**, 1956.

Application of Cycle Property [Problem: 10.15 (1)]

$$G = (V, E), \quad |E| > |V| - 1$$

e : the unique maximum-weighted edge of G

$\implies ?$

$e \notin \text{any MST}$

Application of Cycle Property [Problem: 10.15 (1)]

$$G = (V, E), \quad |E| > |V| - 1$$

e : the unique maximum-weighted edge of G

$\implies ?$

$e \notin \text{any MST}$

Bridge

Application of Cycle Property [Problem: 10.15 (2)]

$$C \subseteq G, \quad e \in C$$

e : the unique maximum-weighted edge of C

\implies

$e \notin \text{any MST}$

Application of Cycle Property [Problem: 10.15 (2)]

$$C \subseteq G, \quad e \in C$$

e : the unique maximum-weighted edge of C

\implies

$e \notin \text{any MST}$

Cycle Property

Application of Cycle Property [Problem: 10.15 (5)]

$$C \subseteq G, \quad e \in C$$

e : the unique lightest edge of C

\Rightarrow ?

$$e \in \forall \text{ MST}$$


Application of Cycle Property [Problem: 10.15 (5)]

$$C \subseteq G, \quad e \in C$$

e : the unique lightest edge of C

\Rightarrow ?

$$e \in \forall \text{ MST}$$



Uniqueness of MST

Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights \implies Unique MST.

Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights \implies Unique MST.

By Contradiction.

Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights \implies Unique MST.

By Contradiction.

\exists MSTs $T_1 \neq T_2$

Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights \implies Unique MST.

By Contradiction.

\exists MSTs $T_1 \neq T_2$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$

Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights \implies Unique MST.

By Contradiction.

\exists MSTs $T_1 \neq T_2$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$

$$e = \min \Delta E$$

Uniqueness of MST [Problem: 10.18 (1)]

Distinct weights \implies Unique MST.

By Contradiction.


\exists MSTs $T_1 \neq T_2$

$$\Delta E = \{e \mid e \in T_1 \setminus T_2 \vee e \in T_2 \setminus T_1\}$$


$$e = \min \Delta E$$

$$e \in T_1 \setminus T_2 \text{ (w.l.o.g.)}$$

$$e \in T_1 \setminus T_2$$




$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$


$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$

$$\exists (e' \in C) \notin T_1$$


$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$

$$\exists (e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E$$


$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$

$$\exists(e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E \implies w(e') > w(e)$$

$$e \in T_1 \setminus T_2$$



$$T_2 + \{e\} \implies C$$

$$\exists(e' \in C) \notin T_1 \implies e' \in T_2 \setminus T_1 \implies e' \in \Delta E \implies w(e') > w(e)$$


$$T' = T_2 + \{e\} - \{e'\} \implies w(T') < w(T_2)$$

Condition for Uniqueness of MST [Problem: 10.18 (2)]

Unique MST $\not\Rightarrow$ Equal weights.

Condition for Uniqueness of MST [Problem: 10.18 (2)]

Unique MST $\not\Rightarrow$ Equal weights.




Unique MST [Problem: 10.21 (3)]

Unique MST $\not\Rightarrow$ Minimum-weight edge across any cut is unique.


Unique MST [Problem: 10.21 (3)]

Unique MST $\not\Rightarrow$ Minimum-weight edge across any cut is unique.



Unique MST [Problem: 10.21 (3)]

Unique MST $\not\Rightarrow$ Minimum-weight edge across any cut is unique.



Theorem (After-class Exercise)


Minimum-weight edge across any cut is unique \implies Unique MST.

Unique MST [Problem: 10.21 (3)]

Unique MST $\not\Rightarrow$ Maximum-weight edge in any cycle is unique.


Unique MST [Problem: 10.21 (3)]

Unique MST $\not\Rightarrow$ Maximum-weight edge in any cycle is unique.



Unique MST [Problem: 10.21 (3)]

Unique MST $\not\Rightarrow$ Maximum-weight edge in any cycle is unique.



Theorem (After-class Exercise)

Maximum-weight edge in any cycle is unique \implies Unique MST.



Unique MST [Problem: 10.21 (4)]

To decide whether a graph has a unique MST.

Unique MST [Problem: 10.21 (4)]


To decide whether a graph has a unique MST.

Ties in Prim's and Kruskal's algorithms

Unique MST [Problem: 10.21 (4)]

To decide whether a graph has a unique MST.


Ties in Prim's and Kruskal's algorithms



Unique MST [Problem: 10.21 (4)]

To decide whether a graph has a unique MST.

Ties in Prim's and Kruskal's algorithms




$$\underbrace{T}_{\text{Any MST}} + \underbrace{\{e\}, \forall e \notin T}_{\text{Cycle}}$$

Unique MST [Problem: 10.21 (4)]

To decide whether a graph has a unique MST.

Ties in Prim's and Kruskal's algorithms



$$\underbrace{T}_{\text{Any MST}} + \underbrace{\{e\}, \forall e \notin T}_{\text{Cycle}}$$

By Kruskal Algorithm.

Critical Edges [Problem: 10.12]

$$G \rightarrow T, \quad G' \triangleq G \setminus \{e\} \rightarrow T'$$
$$w(T') > w(T)$$

Critical Edges [Problem: 10.12]

$$G \rightarrow T, \quad G' \triangleq G \setminus \{e\} \rightarrow T'$$

$$w(T') > w(T)$$


To find all critical edges in $O(m \log m)$ time.

Critical Edges [Problem: 10.12]

$$G \rightarrow T, \quad G' \triangleq G \setminus \{e\} \rightarrow T'$$

$$w(T') > w(T)$$

To find all critical edges in $O(m \log m)$ time.




$$w(e) = 3 \quad w(e) = 2$$

Critical Edges [Problem: 10.12]

$$G \rightarrow T, \quad G' \triangleq G \setminus \{e\} \rightarrow T'$$

$$w(T') > w(T)$$

To find all critical edges in $O(m \log m)$ time.



By Kruskal Algorithm.


$$w(e) = 3 \quad w(e) = 2$$

Critical Edges [Problem: 10.12]

$$G \rightarrow T, \quad G' \triangleq G \setminus \{e\} \rightarrow T'$$

$$w(T') > w(T)$$

To find all critical edges in $O(m \log m)$ time.



By Kruskal Algorithm.

No missing: Check all cycles.


$$w(e) = 3 \quad w(e) = 2$$

Critical Edges [Problem: 10.12]

$$G \rightarrow T, \quad G' \triangleq G \setminus \{e\} \rightarrow T'$$

$$w(T') > w(T)$$

To find all critical edges in $O(m \log m)$ time.



By Kruskal Algorithm.

No missing: Check all cycles.

$O(m \log m)$

$$w(e) = 3 \quad w(e) = 2$$





Office 302

Mailbox: H016

hfwei@nju.edu.cn