

1-11 Set Theory (IV): Infinity

魏恒峰

hfwei@nju.edu.cn

2019 年 12 月 19 日



Finite Sets



Finite Sets



“关于有穷，我原以为我是懂的”

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n$$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n$$

Theorem (Pigeonhole Principle (UD Theorem 22.2))

$$f : \{1, \dots, m\} \rightarrow \{1, \dots, n\} \quad (m, n \in \mathbb{N}^+, m > n)$$

f is not one-to-one.

$A \setminus \{a\}$ (UD Problem 22.17)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.

Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$A \setminus \{a\}$ (UD Problem 22.17)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.
Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$A \setminus \{a\}$ (UD Problem 22.17)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.
Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$$f|_{A \setminus \{a\}} : A \setminus \{a\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\} \setminus \{f(a)\}$$

$A \setminus \{a\}$ (UD Problem 22.17)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.
Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$$f|_{A \setminus \{a\}} : A \setminus \{a\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\} \setminus \{f(a)\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n-1\}$$

(UD Problem 22.18)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 21.10). Show that $|B| \leq |A|$.

(UD Problem 22.18)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 21.10). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

(UD Problem 22.18)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 21.10). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

(UD Problem 22.18)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 21.10). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

(UD Problem 22.18)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 21.10). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

- (c) If two finite sets A and B satisfy $B \subseteq A$ and $|A| \leq |B|$, then $A = B$.

(UD Problem 22.18)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 21.10). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

- (c) If two finite sets A and B satisfy $B \subseteq A$ and $|A| \leq |B|$, then $A = B$.

By contradiction and (b).

$f : A \rightarrow A$ (UD Problem 22.21)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$f : A \rightarrow A$ (UD Problem 22.21)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

$f : A \rightarrow A$ (UD Problem 22.21)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

By Contradiction.

$f : A \rightarrow A$ (UD Problem 22.21)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

By Contradiction.

$$f : A \rightarrow A \setminus \{a\}$$

By Pigeonhole Principle.

$f : A \rightarrow A$ (UD Problem 22.21)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

\impliedby

By Contradiction.

$$f : A \rightarrow A \setminus \{a\}$$

By Pigeonhole Principle.

$f : A \rightarrow A$ (UD Problem 22.21)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

By Contradiction.

\impliedby

By Contradiction.

$$f : A \rightarrow A \setminus \{a\}$$

By Pigeonhole Principle.

$f : A \rightarrow A$ (UD Problem 22.21)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

By Contradiction.

$$f : A \rightarrow A \setminus \{a\}$$

By Pigeonhole Principle.

\impliedby

By Contradiction.

$$\sum_{y \in A} f^{-1}(\{y\}) > |A|$$

Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of pairwise disjoint finite sets whose union is countably infinite.

Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \qquad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \qquad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \qquad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \quad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

- (c) A countably infinite collection of **pairwise disjoint nonempty sets** whose union is finite.

Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \quad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

- (c) A countably infinite collection of **pairwise disjoint nonempty sets** whose union is finite.

$$|A| = n \implies$$

Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \quad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

- (c) A countably infinite collection of **pairwise disjoint nonempty sets** whose union is finite.

$$|A| = n \implies |\mathcal{P}(A)| = 2^n$$

UD Problem 23.3 (d)

Is it countable or uncountable?

$$A = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x + y = 1\}$$

UD Problem 23.3 (d)

Is it countable or uncountable?

$$A = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x + y = 1\}$$

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} A$$

UD Problem 23.3 (d)

Is it countable or uncountable?

$$A = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x + y = 1\}$$

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} A$$

$$f(x) = (x, 1 - x)$$

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$\begin{array}{l} s_1 = 0000000000\dots \\ s_2 = 1111111111\dots \\ s_3 = 0101010101\dots \\ s_4 = 1010101010\dots \\ s_5 = 1101010101\dots \\ s_6 = 0011010110\dots \\ s_7 = 10001000100\dots \\ s_8 = 0011001001\dots \\ s_9 = 11001100110\dots \\ s_{10} = 11011100101\dots \\ s_{11} = 11010100100\dots \\ \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \end{array}$$

$$s = 10111010011\dots$$

By Diagonal Argument.

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

$$f(x_0x_1\cdots) = \sum_{i=0}^{\infty} x_i 2^i$$

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

$$f(x_0x_1\cdots) = \sum_{i=0}^{\infty} x_i 2^i$$



Complex Numbers (UD Problem 24.16)

Prove that

$$|\mathbb{R}| = |\mathbb{C}|, \quad \mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$$

Complex Numbers (UD Problem 24.16)

Prove that

$$|\mathbb{R}| = |\mathbb{C}|, \quad \mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$$

$$|\mathbb{C}| = |\mathbb{R} \times \mathbb{R}|$$

Complex Numbers (UD Problem 24.16)

Prove that

$$|\mathbb{R}| = |\mathbb{C}|, \quad \mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$$

$$|\mathbb{C}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}|$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots)$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$



Was Cantor Surprised?

UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$

UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$

$$f(x) = (x, 0.5)$$

UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$

$$f(x) = (x, 0.5)$$

$$g : (0, 1) \times (0, 1) \rightarrow (0, 1)$$

UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$

$$f(x) = (x, 0.5)$$

$$g : (0, 1) \times (0, 1) \rightarrow (0, 1)$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$

Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \dots$$

Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \dots$$

$$f(0) = \frac{1}{2} \quad f(1) = \frac{1}{3}$$

Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \cdots$$

$$f(0) = \frac{1}{2} \quad f(1) = \frac{1}{3}$$

$$\forall n \geq 4 : f\left(\frac{1}{n-2}\right) = \frac{1}{n}$$

Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \cdots$$

$$f(0) = \frac{1}{2} \quad f(1) = \frac{1}{3}$$

$$\forall n \geq 4 : f\left(\frac{1}{n-2}\right) = \frac{1}{n}$$

$$f(x) = x, \text{ otherwise}$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$f(x) = \frac{x}{x+1}$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$f(x) = \frac{x}{x+1}$$

$$[0, 1] \approx (0, 1]$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$f(x) = \frac{x}{x+1}$$

$$[0, 1] \approx (0, 1]$$

$$f(0) = \frac{1}{2} \quad f\left(\frac{1}{2}\right) = \frac{2}{3} \quad f\left(\frac{2}{3}\right) = \frac{3}{4} \quad \cdots \quad f(x) = x$$

Thank
You!