

# 1-10 Set Theory (III): Functions

魏恒峰

hfwei@nju.edu.cn

2019 年 12 月 12 日



$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$x \in f^{-1}(Y) \iff f(x) \in Y$$

$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$x \in f^{-1}(Y) \iff f(x) \in Y$$

$$a \in A \implies f(a) \in f(A)$$

$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$x \in f^{-1}(Y) \iff f(x) \in Y$$

$$a \in A \implies f(a) \in f(A)$$

$$f(A_1 \cup A_2) = f(A_1) \cup f(A_2)$$

$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$x \in f^{-1}(Y) \iff f(x) \in Y$$

$$a \in A \implies f(a) \in f(A)$$

$$f(A_1 \cup A_2) = f(A_1) \cup f(A_2) \qquad f\left(\bigcup_{\alpha \in I} A_\alpha\right) = \bigcup \{f(A_\alpha) \mid \alpha \in I\}$$

$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$x \in f^{-1}(Y) \iff f(x) \in Y$$

$$a \in A \implies f(a) \in f(A)$$

$$f(A_1 \cup A_2) = f(A_1) \cup f(A_2) \qquad f\left(\bigcup_{\alpha \in I} A_\alpha\right) = \bigcup \{f(A_\alpha) \mid \alpha \in I\}$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$x \in f^{-1}(Y) \iff f(x) \in Y$$

$$a \in A \implies f(a) \in f(A)$$

$$f(A_1 \cup A_2) = f(A_1) \cup f(A_2) \qquad f\left(\bigcup_{\alpha \in I} A_\alpha\right) = \bigcup \{f(A_\alpha) \mid \alpha \in I\}$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2) \qquad f\left(\bigcap_{\alpha \in I} A_\alpha\right) \subseteq \bigcap \{f(A_\alpha) \mid \alpha \in I\}$$



## UD Problem 16.14

$$f : A \rightarrow B \quad g_1, g_2 : B \rightarrow A$$

$$f \circ g_1 = f \circ g_2$$

- (a) Show that if  $f$  is bijective, then  $g_1 = g_2$ .
- (b) If  $g_1 \circ f = g_2 \circ f$  and  $f$  is bijective, must  $g_1 = g_2$ ?

## UD Problem 16.14

$$f : A \rightarrow B \quad g_1, g_2 : B \rightarrow A$$

$$f \circ g_1 = f \circ g_2$$

- (a) Show that if  $f$  is bijective, then  $g_1 = g_2$ .
- (b) If  $g_1 \circ f = g_2 \circ f$  and  $f$  is bijective, must  $g_1 = g_2$ ?

$$f^{-1} \circ (f \circ g_1) = f^{-1} \circ (f \circ g_2)$$

## UD Problem 16.14

$$f : A \rightarrow B \quad g_1, g_2 : B \rightarrow A$$

$$f \circ g_1 = f \circ g_2$$

- (a) Show that if  $f$  is bijective, then  $g_1 = g_2$ .
- (b) If  $g_1 \circ f = g_2 \circ f$  and  $f$  is bijective, must  $g_1 = g_2$ ?

$$f^{-1} \circ (f \circ g_1) = f^{-1} \circ (f \circ g_2)$$

$$(g_1 \circ f) \circ f^{-1} = (g_2 \circ f) \circ f^{-1}$$

## Theorem

$$f \circ g_1 = f \circ g_2 \wedge f \text{ is injective} \implies g_1 = g_2.$$

## Theorem

$$f \circ g_1 = f \circ g_2 \wedge f \text{ is injective} \implies g_1 = g_2.$$

$$(f \circ g_1)(x) = (f \circ g_2)(x)$$

## Theorem

$$f \circ g_1 = f \circ g_2 \wedge f \text{ is injective} \implies g_1 = g_2.$$

$$\begin{aligned} (f \circ g_1)(x) &= (f \circ g_2)(x) \\ \implies f(g_1(x)) &= f(g_2(x)) \end{aligned}$$

## Theorem

$$f \circ g_1 = f \circ g_2 \wedge f \text{ is injective} \implies g_1 = g_2.$$

$$\begin{aligned} & (f \circ g_1)(x) = (f \circ g_2)(x) \\ \implies & f(g_1(x)) = f(g_2(x)) \\ \implies & g_1(x) = g_2(x) \quad (f \text{ is injective}) \end{aligned}$$

## Theorem

$$g_1 \circ f = g_2 \circ f \wedge f \text{ is surjective} \implies g_1 = g_2.$$



## Theorem

$$g_1 \circ f = g_2 \circ f \wedge f \text{ is surjective} \implies g_1 = g_2.$$

$$\forall b \in B : \exists a \in A : b = f(a)$$

## Theorem

$$g_1 \circ f = g_2 \circ f \wedge f \text{ is surjective} \implies g_1 = g_2.$$

$$\forall b \in B : \exists a \in A : b = f(a)$$

$$g_1(b) = g_1(f(a))$$

## Theorem

$$g_1 \circ f = g_2 \circ f \wedge f \text{ is surjective} \implies g_1 = g_2.$$

$$\forall b \in B : \exists a \in A : b = f(a)$$

$$g_1(b) = g_1(f(a)) = g_2(f(a))$$

## Theorem

$$g_1 \circ f = g_2 \circ f \wedge f \text{ is surjective} \implies g_1 = g_2.$$

$$\forall b \in B : \exists a \in A : b = f(a)$$

$$g_1(b) = g_1(f(a)) = g_2(f(a)) = g_2(b)$$

## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ .  
Prove that  $f$  is bijective.

## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ .  
Prove that  $f$  is bijective.

$f$  is onto

## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ .  
Prove that  $f$  is bijective.

$f$  is onto

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = y$$

## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ . Prove that  $f$  is bijective.

$f$  is onto

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = y$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f^3(x) = f^2(y)$$



## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ . Prove that  $f$  is bijective.

$f$  is onto

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = y$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f^3(x) = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = f^2(y)$$

## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ . Prove that  $f$  is bijective.

$f$  is onto

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = y$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f^3(x) = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : y = f^2(y)$$

## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ . Prove that  $f$  is bijective.

$f$  is onto

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = y$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f^3(x) = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : y = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : y = f^2(y)$$

## (Composition) UD Problem 16.22

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f$  is onto and  $f \circ f \circ f = f$ . Prove that  $f$  is bijective.

$f$  is onto

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = y$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f^3(x) = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : f(x) = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : \exists x \in \mathbb{R} : y = f^2(y)$$

$$\implies \forall y \in \mathbb{R} : y = f^2(y)$$

$$\implies f^2 = Id_{\mathbb{R}}$$

## Theorem

If  $f : X \rightarrow X$ , then

$$f^2 = Id_X \implies f \text{ is injective.}$$

## Theorem

If  $f : X \rightarrow X$ , then

$$f^2 = Id_X \implies f \text{ is injective.}$$

$$f(x_1) = f(x_2)$$

## Theorem

If  $f : X \rightarrow X$ , then

$$f^2 = Id_X \implies f \text{ is injective.}$$

$$\begin{aligned} f(x_1) &= f(x_2) \\ \implies f^2(x_1) &= f^2(x_2) \end{aligned}$$

## Theorem

If  $f : X \rightarrow X$ , then

$$f^2 = Id_X \implies f \text{ is injective.}$$

$$\begin{aligned} f(x_1) &= f(x_2) \\ \implies f^2(x_1) &= f^2(x_2) \\ \implies Id_X(x_1) &= Id_X(x_2) \end{aligned}$$



## Theorem

If  $f : X \rightarrow X$ , then

$$f^2 = Id_X \implies f \text{ is injective.}$$

$$\begin{aligned} f(x_1) &= f(x_2) \\ \implies f^2(x_1) &= f^2(x_2) \\ \implies Id_X(x_1) &= Id_X(x_2) \\ \implies x_1 &= x_2 \end{aligned}$$

## Image (UD Problem 17.22)

$$f : A \rightarrow B, \quad A_1, A_2 \subseteq A$$

- (i) If  $f(A_1) = f(A_2)$ , must  $A_1 = A_2$ ?
- (ii) When is  $f(A_1) = f(A_2) \implies A_1 = A_2$ ?

## Image (UD Problem 17.22)

$$f : A \rightarrow B, \quad A_1, A_2 \subseteq A$$

- (i) If  $f(A_1) = f(A_2)$ , must  $A_1 = A_2$ ?
- (ii) When is  $f(A_1) = f(A_2) \implies A_1 = A_2$ ?

$$a_1 \in A_1$$

## Image (UD Problem 17.22)

$$f : A \rightarrow B, \quad A_1, A_2 \subseteq A$$

- (i) If  $f(A_1) = f(A_2)$ , must  $A_1 = A_2$ ?
- (ii) When is  $f(A_1) = f(A_2) \implies A_1 = A_2$ ?

$$\begin{aligned} a_1 &\in A_1 \\ \implies f(a_1) &\in f(A_1) \end{aligned}$$

## Image (UD Problem 17.22)

$$f : A \rightarrow B, \quad A_1, A_2 \subseteq A$$

- (i) If  $f(A_1) = f(A_2)$ , must  $A_1 = A_2$ ?
- (ii) When is  $f(A_1) = f(A_2) \implies A_1 = A_2$ ?

$$a_1 \in A_1$$

$$\implies f(a_1) \in f(A_1)$$

$$\implies f(a_1) \in f(A_2)$$

## Image (UD Problem 17.22)

$$f : A \rightarrow B, \quad A_1, A_2 \subseteq A$$

- (i) If  $f(A_1) = f(A_2)$ , must  $A_1 = A_2$ ?
- (ii) When is  $f(A_1) = f(A_2) \implies A_1 = A_2$ ?

$$a_1 \in A_1$$

$$\implies f(a_1) \in f(A_1)$$

$$\implies f(a_1) \in f(A_2)$$

$$\implies \exists a_2 \in A_2 : f(a_2) = f(a_1)$$

## Image (UD Problem 17.22)

$$f : A \rightarrow B, \quad A_1, A_2 \subseteq A$$

- (i) If  $f(A_1) = f(A_2)$ , must  $A_1 = A_2$ ?
- (ii) When is  $f(A_1) = f(A_2) \implies A_1 = A_2$ ?

$$a_1 \in A_1$$

$$\implies f(a_1) \in f(A_1)$$

$$\implies f(a_1) \in f(A_2)$$

$$\implies \exists a_2 \in A_2 : f(a_2) = f(a_1)$$

$$\implies a_1 = a_2 \quad (\text{if } f \text{ is injective})$$

## Image (UD Problem 17.22)

$$f : A \rightarrow B, \quad A_1, A_2 \subseteq A$$

- (i) If  $f(A_1) = f(A_2)$ , must  $A_1 = A_2$ ?
- (ii) When is  $f(A_1) = f(A_2) \implies A_1 = A_2$ ?

$$\begin{aligned} & a_1 \in A_1 \\ \implies & f(a_1) \in f(A_1) \\ \implies & f(a_1) \in f(A_2) \\ \implies & \exists a_2 \in A_2 : f(a_2) = f(a_1) \\ \implies & a_1 = a_2 \quad (\text{if } f \text{ is injective}) \\ \implies & a_1 \in A_2 \end{aligned}$$



## (Inverse Image) UD Problem 17.23

$$f : A \rightarrow B, \quad B_1, B_2 \subseteq B$$

- (i) If  $f^{-1}(B_1) = f^{-1}(B_2)$ , must  $B_1 = B_2$ ?
- (ii) When is  $f^{-1}(B_1) = f^{-1}(B_2) \implies B_1 = B_2$ ?

## (Inverse Image) UD Problem 17.23

$$f : A \rightarrow B, \quad B_1, B_2 \subseteq B$$

- (i) If  $f^{-1}(B_1) = f^{-1}(B_2)$ , must  $B_1 = B_2$ ?
- (ii) When is  $f^{-1}(B_1) = f^{-1}(B_2) \implies B_1 = B_2$ ?

$$b_1 \in B_1$$

## (Inverse Image) UD Problem 17.23

$$f : A \rightarrow B, \quad B_1, B_2 \subseteq B$$

- (i) If  $f^{-1}(B_1) = f^{-1}(B_2)$ , must  $B_1 = B_2$ ?
- (ii) When is  $f^{-1}(B_1) = f^{-1}(B_2) \implies B_1 = B_2$ ?

$$\begin{aligned} & b_1 \in B_1 \\ \implies & \exists a_1 \in A : f(a_1) = b_1 \in B_1 \quad (\text{if } f \text{ is surjective}) \end{aligned}$$

## (Inverse Image) UD Problem 17.23

$$f : A \rightarrow B, \quad B_1, B_2 \subseteq B$$

- (i) If  $f^{-1}(B_1) = f^{-1}(B_2)$ , must  $B_1 = B_2$ ?
- (ii) When is  $f^{-1}(B_1) = f^{-1}(B_2) \implies B_1 = B_2$ ?

$$b_1 \in B_1$$

$$\implies \exists a_1 \in A : f(a_1) = b_1 \in B_1 \quad (\text{if } f \text{ is surjective})$$

$$\implies \exists a_1 \in A : a_1 \in f^{-1}(B_1)$$

## (Inverse Image) UD Problem 17.23

$$f : A \rightarrow B, \quad B_1, B_2 \subseteq B$$

- (i) If  $f^{-1}(B_1) = f^{-1}(B_2)$ , must  $B_1 = B_2$ ?
- (ii) When is  $f^{-1}(B_1) = f^{-1}(B_2) \implies B_1 = B_2$ ?

$$b_1 \in B_1$$

$$\implies \exists a_1 \in A : f(a_1) = b_1 \in B_1 \quad (\text{if } f \text{ is surjective})$$

$$\implies \exists a_1 \in A : a_1 \in f^{-1}(B_1)$$

$$\implies \exists a_1 \in A : a_1 \in f^{-1}(B_2)$$

## (Inverse Image) UD Problem 17.23

$$f : A \rightarrow B, \quad B_1, B_2 \subseteq B$$

- (i) If  $f^{-1}(B_1) = f^{-1}(B_2)$ , must  $B_1 = B_2$ ?
- (ii) When is  $f^{-1}(B_1) = f^{-1}(B_2) \implies B_1 = B_2$ ?

$$b_1 \in B_1$$

$$\implies \exists a_1 \in A : f(a_1) = b_1 \in B_1 \quad (\text{if } f \text{ is surjective})$$

$$\implies \exists a_1 \in A : a_1 \in f^{-1}(B_1)$$

$$\implies \exists a_1 \in A : a_1 \in f^{-1}(B_2)$$

$$\implies \exists a_1 \in A : f(a_1) \in B_2$$

## (Inverse Image) UD Problem 17.23

$$f : A \rightarrow B, \quad B_1, B_2 \subseteq B$$

- (i) If  $f^{-1}(B_1) = f^{-1}(B_2)$ , must  $B_1 = B_2$ ?
- (ii) When is  $f^{-1}(B_1) = f^{-1}(B_2) \implies B_1 = B_2$ ?

$$b_1 \in B_1$$

$$\implies \exists a_1 \in A : f(a_1) = b_1 \in B_1 \quad (\text{if } f \text{ is surjective})$$

$$\implies \exists a_1 \in A : a_1 \in f^{-1}(B_1)$$

$$\implies \exists a_1 \in A : a_1 \in f^{-1}(B_2)$$

$$\implies \exists a_1 \in A : f(a_1) \in B_2$$

$$\implies b_1 \in B_2 \quad (f(a_1) = b_1, \text{ since } f \text{ is a function})$$

## Monotonicity

Assume that  $F : \mathcal{P}(A) \rightarrow \mathcal{P}(A)$  and that  $F$  has the monotonicity property:

$$X \subseteq Y \subseteq A \implies F(X) \subseteq F(Y).$$

Define

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

- (a) Show that  $F(B) = B$  and  $F(C) = C$ .
- (b) Show that if  $F(X) = X$ , then  $B \subseteq X \subseteq C$ .



## Monotonicity

Assume that  $F : \mathcal{P}(A) \rightarrow \mathcal{P}(A)$  and that  $F$  has the monotonicity property:

$$X \subseteq Y \subseteq A \implies F(X) \subseteq F(Y).$$

Define

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

- (a) Show that  $F(B) = B$  and  $F(C) = C$ .
- (b) Show that if  $F(X) = X$ , then  $B \subseteq X \subseteq C$ .

$$F(X) = X$$

## Monotonicity

Assume that  $F : \mathcal{P}(A) \rightarrow \mathcal{P}(A)$  and that  $F$  has the monotonicity property:

$$X \subseteq Y \subseteq A \implies F(X) \subseteq F(Y).$$

Define

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

- (a) Show that  $F(B) = B$  and  $F(C) = C$ .
- (b) Show that if  $F(X) = X$ , then  $B \subseteq X \subseteq C$ .

$$F(X) = X \implies F(X) \subseteq X \wedge X \subseteq F(X)$$

## Monotonicity

Assume that  $F : \mathcal{P}(A) \rightarrow \mathcal{P}(A)$  and that  $F$  has the monotonicity property:

$$X \subseteq Y \subseteq A \implies F(X) \subseteq F(Y).$$

Define

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

- (a) Show that  $F(B) = B$  and  $F(C) = C$ .
- (b) Show that if  $F(X) = X$ , then  $B \subseteq X \subseteq C$ .

$$F(X) = X \implies F(X) \subseteq X \wedge X \subseteq F(X) \implies B \subseteq X \subseteq C$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$F(B) \subseteq B$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$F(B) \subseteq B$$

$$F(B) = F\left(\bigcap \{X \subseteq A \mid F(X) \subseteq X\}\right)$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$F(B) \subseteq B$$

$$\begin{aligned} F(B) &= F\left(\bigcap \{X \subseteq A \mid F(X) \subseteq X\}\right) \\ &\subseteq \bigcap \{F(X) : X \subseteq A \wedge F(X) \subseteq X\} \end{aligned}$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$F(B) \subseteq B$$

$$\begin{aligned} F(B) &= F\left(\bigcap \{X \subseteq A \mid F(X) \subseteq X\}\right) \\ &\subseteq \bigcap \{F(X) : X \subseteq A \wedge F(X) \subseteq X\} \\ &\subseteq \bigcap \{X : X \subseteq A \wedge F(X) \subseteq X\} \end{aligned}$$



$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$F(B) \subseteq B$$

$$\begin{aligned} F(B) &= F\left(\bigcap \{X \subseteq A \mid F(X) \subseteq X\}\right) \\ &\subseteq \bigcap \{F(X) : X \subseteq A \wedge F(X) \subseteq X\} \\ &\subseteq \bigcap \{X : X \subseteq A \wedge F(X) \subseteq X\} \\ &= B \end{aligned}$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$B \subseteq F(B)$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$B \subseteq F(B)$$

$$F(B) \subseteq B$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$B \subseteq F(B)$$

$$F(B) \subseteq B \implies F(F(B)) \subseteq F(B)$$

$$B = \bigcap \{X \subseteq A \mid F(X) \subseteq X\}$$

$$B \subseteq F(B)$$

$$F(B) \subseteq B \implies F(F(B)) \subseteq F(B) \implies B \subseteq F(B)$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$C \subseteq F(C)$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$C \subseteq F(C)$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}$$



$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$C \subseteq F(C)$$

$$\begin{aligned} C &= \bigcup \{X \subseteq A \mid X \subseteq F(X)\} \\ &\subseteq \bigcup \{F(X) \subseteq A \mid X \subseteq F(X)\} \end{aligned}$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$C \subseteq F(C)$$

$$\begin{aligned} C &= \bigcup \{X \subseteq A \mid X \subseteq F(X)\} \\ &\subseteq \bigcup \{F(X) \subseteq A \mid X \subseteq F(X)\} \\ &= F\left(\bigcup \{X \subseteq A \mid X \subseteq F(X)\}\right) \end{aligned}$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$C \subseteq F(C)$$

$$\begin{aligned} C &= \bigcup \{X \subseteq A \mid X \subseteq F(X)\} \\ &\subseteq \bigcup \{F(X) \subseteq A \mid X \subseteq F(X)\} \\ &= F\left(\bigcup \{X \subseteq A \mid X \subseteq F(X)\}\right) \\ &= F(C) \end{aligned}$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$F(C) \subseteq C$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$F(C) \subseteq C$$

$$C \subseteq F(C)$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$F(C) \subseteq C$$

$$C \subseteq F(C) \implies F(C) \subseteq F(F(C))$$

$$C = \bigcup \{X \subseteq A \mid X \subseteq F(X)\}.$$

$$F(C) \subseteq C$$

$$C \subseteq F(C) \implies F(C) \subseteq F(F(C)) \implies F(C) \subseteq C$$



Thank  
You!