

# 1-12 Partial Order and Lattice Theory

魏恒峰

hfwei@nju.edu.cn

2020 年 02 月 23 日





### SM Problem 14.44

Suppose the following are three consistent enumerations of an ordered set  $A = \{a, b, c, d\}$ :

$A_1 :$      $a$         $b$         $c$         $d$

$A_2 :$      $a$         $c$         $b$         $d$

$A_3 :$      $a$         $c$         $d$         $b$

Assuming the Hasse diagram  $D$  of  $A$  is **connected**, draw  $D$ .

### SM Problem 14.44

Suppose the following are three consistent enumerations of an ordered set  $A = \{a, b, c, d\}$ :

$$A_1 : \quad a \quad \textcolor{red}{b} \quad \textcolor{red}{c} \quad d$$


$$A_2 : \quad a \quad c \quad b \quad d$$



$$A_3 : \quad a \quad c \quad \textcolor{blue}{d} \quad \textcolor{blue}{b}$$



Assuming the Hasse diagram  $D$  of  $A$  is **connected**, draw  $D$ .

$$b \prec_{A_1} c \wedge c \prec_{A_2} b \implies b \parallel_A c$$




$$d \prec_{A_3} b \wedge b \prec_{A_2} d \implies b \parallel_A d$$









$$\# = 6$$




$\# = 6$





$\# = 6$



$\# = 3$



## Theorem



## Theorem

Let  $A$  be a nonempty finite set with  $|A| = n$  and let  $a \in A$ .  
Prove that  $A \setminus \{a\}$  is finite and  $|A \setminus \{a\}| = n - 1$ .





## Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of pairwise disjoint finite sets whose union is countably infinite.

## Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \qquad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$



## Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \qquad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

## Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \quad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

## Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \quad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

- (c) A countably infinite collection of **pairwise disjoint nonempty sets** whose union is finite.

## Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \quad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

- (c) A countably infinite collection of **pairwise disjoint nonempty sets** whose union is finite.

$$|A| = n \implies$$

## Set Union (UD Problem 23.1)

Give an example, if possible, of

- (a) A countably infinite collection of **pairwise disjoint finite sets** whose union is countably infinite.

$$\forall n \in \mathbb{N} : A_n = \{n\} \quad \bigcup_{n \in \mathbb{N}} A_n = \mathbb{N}$$

- (b) A countably infinite collection of nonempty sets whose union is finite.

$$\forall n \in \mathbb{N} : A_n = \{1\} \quad \bigcup_{n \in \mathbb{N}} A_n = \{1\}$$

- (c) A countably infinite collection of **pairwise disjoint nonempty sets** whose union is finite.

$$|A| = n \implies |\mathcal{P}(A)| = 2^n$$

### UD Problem 23.3 (d)

Is it countable or uncountable?

$$A = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x + y = 1\}$$

### UD Problem 23.3 (d)

Is it countable or uncountable?

$$A = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x + y = 1\}$$

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} A$$

### UD Problem 23.3 (d)

Is it countable or uncountable?

$$A = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x + y = 1\}$$

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} A$$

$$f(x) = (x, 1 - x)$$



## Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

## Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

By Diagonal Argument.

## Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

## Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

$$f(x_0x_1\cdots) = \sum_{i=0}^{\infty} x_i 2^i$$

## Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

$$f(x_0x_1\cdots) = \sum_{i=0}^{\infty} x_i 2^i$$

## Complex Numbers (UD Problem 24.16)

Prove that

$$|\mathbb{R}| = |\mathbb{C}|, \quad \mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$$

## Complex Numbers (UD Problem 24.16)

Prove that

$$|\mathbb{R}| = |\mathbb{C}|, \quad \mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$$

$$|\mathbb{C}| = |\mathbb{R} \times \mathbb{R}|$$

## Complex Numbers (UD Problem 24.16)

Prove that

$$|\mathbb{R}| = |\mathbb{C}|, \quad \mathbb{C} = \{a + bi : a, b \in \mathbb{R}\}$$

$$|\mathbb{C}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}|$$



$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots)$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$

$$\mathbb{R} \times \mathbb{R} \approx \mathbb{R}$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$

Was Cantor Surprised?

## UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

## UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

## UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

## UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$



## UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$

$$f(x) = (x, 0.5)$$

## UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$

$$f(x) = (x, 0.5)$$

$$g : (0, 1) \times (0, 1) \rightarrow (0, 1)$$

## UD Problem 24.15

$$(0, 1) \approx (0, 1) \times (0, 1)$$

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : X \rightarrow Y \wedge g : Y \rightarrow X \implies \exists \text{ bijection } h : X \rightarrow Y$$

$$f : (0, 1) \rightarrow (0, 1) \times (0, 1)$$

$$f(x) = (x, 0.5)$$

$$g : (0, 1) \times (0, 1) \rightarrow (0, 1)$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$

## Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

## Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \dots$$

## Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \dots$$

$$f(0) = \frac{1}{2} \quad f(1) = \frac{1}{3}$$

## Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \dots$$

$$f(0) = \frac{1}{2} \quad f(1) = \frac{1}{3}$$

$$\forall n \geq 4 : f\left(\frac{1}{n-2}\right) = \frac{1}{n}$$

## Bijections (UD Problem 21.21)

$$[0, 1] \approx (0, 1)$$

$$0, 1, \quad \frac{1}{2}, \frac{1}{3}, \quad \frac{1}{4}, \frac{1}{5} \dots$$

$$f(0) = \frac{1}{2} \quad f(1) = \frac{1}{3}$$

$$\forall n \geq 4 : f\left(\frac{1}{n-2}\right) = \frac{1}{n}$$

$$f(x) = x, \text{ otherwise}$$



$$(-\infty, \infty) \approx (0, \infty)$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$f(x) = \frac{x}{x+1}$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$f(x) = \frac{x}{x+1}$$

$$[0, 1] \approx (0, 1]$$

$$(-\infty, \infty) \approx (0, \infty)$$

$$f(x) = e^x$$

$$(0, \infty) \approx (0, 1)$$

$$f(x) = \frac{x}{x+1}$$

$$[0, 1] \approx (0, 1]$$

$$f(0) = \frac{1}{2} \quad f\left(\frac{1}{2}\right) = \frac{2}{3} \quad f\left(\frac{2}{3}\right) = \frac{3}{4} \quad \cdots \quad f(x) = x$$

Thank  
You!