

score: 400

500

3-2 Greedy Algorithms

(How to justify your greed?)


Hengfeng Wei

hfwei@nju.edu.cn


September 29, 2018



"GREED IS GOOD."



“GREED IS GOOD.”



“BUT”

*“Greedy algorithms without **proofs** are considered wrong.”*

Proof: Inductive Exchange Argument

Proof: Inductive Exchange Argument

Theorem (Greedy-Choice Property)

There is an optimal solution that makes the greedy choice.

Theorem (Optimal Substructure Property)

A problem exhibits optimal substructure if an optimal solution to the problem contains within it optimal solutions to subproblems.

Live with Your Deadlines (Additional Problem)

Live with Your Deadlines (Additional Problem)



Live with Your Deadlines (Additional Problem)

$HW_i : (Length : t_i, \ Deadline : d_i)$



4

$OT : 2$

$PS : 5$

$OJ : 6$

$PA : 8$

Live with Your Deadlines (Additional Problem)

$HW_i : (Length : t_i, \ Deadline : d_i)$



4

$OT : 2$

$PS : 5$

$OJ : 6$

$PA : 8$

$Finish\ time : f_i = s_i + t_i$

$\text{Lateness} : l_i = f_i - d_i \quad (0 \text{ if } f_i \leq d_i)$

Live with Your Deadlines (Additional Problem)

$HW_i : (Length : t_i, \ Deadline : d_i)$



4

$OT : 2$

$PS : 5$

$OJ : 6$

$PA : 8$

$Finish\ time : f_i = s_i + t_i$

$\text{Lateness} : l_i = f_i - d_i \quad (0 \text{ if } f_i \leq d_i)$

$$\min (L \triangleq \max_i l_i)$$

What are your greedy strategies?

**“I love deadlines.
I like the
whooshing
sound they
make as they
fly by. ”**

Douglas Adams

Shortest Job First: $\min t_i$

Shortest Job First: $\min t_i$

Urgentest Job First: $\min(d_i - t_i)$

$$t_1 = 1, d_1 = 100$$

$$t_2 = 10, d_2 = 10$$

Shortest Job First: $\min t_i$

$$t_1 = 1, d_1 = 100$$

$$t_2 = 10, d_2 = 10$$

Urgentest Job First: $\min(d_i - t_i)$

$$t_1 = 1, d_1 = 2$$

$$t_2 = 10, d_2 = 10$$

Earliest Deadline First



Proof: Inductive Exchange Argument.

$OPT : J_1^*, J_2^*, \dots, J_n^*$

$G : J_1, J_2, \dots, J_n, \quad (d_i \leq d_j, \forall i \leq j)$

$\exists J_i : d_{J_i} < d_{J_1^*}$

$$d_{J_i} < d_{J_1^*}$$

J_1^*		J_i^*		J_i		
---------	--	---------	--	-------	--	--

J_i		J_i^*		J_1^*		
-------	--	---------	--	---------	--	--

$$d_{J_i} < d_{J_1^*}$$

J_1^*		J_i^*		J_i		
---------	--	---------	--	-------	--	--

J_i		J_i^*		J_1^*		
-------	--	---------	--	---------	--	--

$$l'_{J_1^*} = f'_{J_1^*} - d_{J_1^*} = f_{J_i} - d_{J_1^*} < f_{J_i} - d_{J_i} = l_{J_i}$$

$$d_{J_i} < d_{J_1^*}$$

J_1^*		J_i^*		J_i		
---------	--	---------	--	-------	--	--

J_i		J_i^*		J_1^*		
-------	--	---------	--	---------	--	--

$$l'_{J_1^*} = f'_{J_1^*} - d_{J_1^*} = f_{J_i} - d_{J_1^*} < f_{J_i} - d_{J_i} = l_{J_i}$$

$$l'_{J_i^*} = f'_{J_i^*} - d_{J_i^*} < f_{J_i} - d_{J_i^*} \leq f_{J_i} - d_{J_i} = l_{J_i}$$

Theorem

All schedules with no idle time and no inversions have the same maximum lateness.

Theorem (Optimal Substructure Property)

A problem exhibits *optimal substructure* if an optimal solution to the problem contains within it optimal solutions to subproblems.

Theorem (Optimal Substructure Property)

A problem exhibits *optimal substructure* if an optimal solution to the problem contains within it optimal solutions to subproblems.

$$L(s, S)$$

The optimal lateness obtainable from scheduling the jobs in S

which are available to start at the common start time s .

Theorem (Optimal Substructure Property)

A problem exhibits *optimal substructure* if an optimal solution to the problem contains within it optimal solutions to subproblems.

$$L(s, S)$$

The optimal lateness obtainable from scheduling the jobs in S

which are available to start at the common start time s .

What is the first job to schedule?

$$L(s, S) = \min_{1 \leq i \leq n} \left(\max (L(0, \{J_i\}), L(t_i, J \setminus \{J_i\})) \right)$$

$$J = \{J_1, J_2, J_3\}$$

$$t_1 = 6, t_2 = 2, t_3 = 2$$

$$d_1 = 2, d_2 = 7, d_3 = 8$$

$$J = \{J_1, J_2, J_3\}$$

$$t_1 = 6, t_2 = 2, t_3 = 2$$

$$d_1 = 2, d_2 = 7, d_3 = 8$$

$$\mathcal{O} : J_1, J_3, J_2$$

$$L(0, \{J_1, J_2, J_3\}) = 4$$

$$J = \{J_1, J_2, J_3\}$$

$$t_1 = 6, t_2 = 2, t_3 = 2$$

$$d_1 = 2, d_2 = 7, d_3 = 8$$

$$\mathcal{O} : J_1, J_3, J_2$$

$$\mathcal{O}' : J_2, J_3$$

$$L(0, \{J_1, J_2, J_3\}) = 4$$

$$L(6, \{J_2, J_3\}) = 2$$

$$J = \{J_1, J_2, J_3\}$$

$$t_1 = 6, t_2 = 2, t_3 = 2$$

$$d_1 = 2, d_2 = 7, d_3 = 8$$

$$\mathcal{O} : J_1, J_3, J_2$$

$$\mathcal{O}' : J_2, J_3$$

$$L(0, \{J_1, J_2, J_3\}) = 4$$

$$L(6, \{J_2, J_3\}) = 2$$

$$\mathcal{O}' \not\subset \mathcal{O}$$

$$J = \{J_1, J_2, J_3\}$$

$$t_1 = 6, t_2 = 2, t_3 = 2$$

$$d_1 = 2, d_2 = 7, d_3 = 8$$

$$\mathcal{O} : J_1, J_3, J_2$$

$$\mathcal{O}' : J_2, J_3$$

$$L(0, \{J_1, J_2, J_3\}) = 4$$

$$L(6, \{J_2, J_3\}) = 2$$

$$\mathcal{O}' \not\subset \mathcal{O}$$

$$\exists \mathcal{O} \triangleq J_1, J_2, J_3 : \mathcal{O}' \subset \mathcal{O}$$

Theorem

The greedy solution has no idle time and no inversions.

Theorem

The greedy solution has no idle time and no inversions.

Theorem

There is an optimal schedule with no idle time.

Theorem

The greedy solution has no idle time and no inversions.

Theorem

There is an optimal schedule with no idle time.

Theorem

There is an optimal schedule with no idle time and no inversions.

Theorem

The greedy solution has no idle time and no inversions.

Theorem

There is an optimal schedule with no idle time.

Theorem

There is an optimal schedule with no idle time and no inversions.

By Exchange Argument.

Theorem

The greedy solution has no idle time and no inversions.

Theorem

There is an optimal schedule with no idle time.

Theorem

There is an optimal schedule with no idle time and no inversions.

By Exchange Argument.

Theorem

All schedules with no idle time and no inversions have the same maximum lateness.

Theorem

There is an optimal schedule with no idle time and no inversions.

Theorem

There is an optimal schedule with no idle time and no inversions.

Immediate Inversion

Theorem

There is an optimal schedule with no idle time and no inversions.



Immediate Inversion

Theorem

There is an optimal schedule with no idle time and no inversions.

Immediate Inversion

$$d_j < d_i$$




Theorem

There is an optimal schedule with no idle time and no inversions.

Immediate Inversion

$$d_j < d_i$$

$$l'_j \leq l_j$$




Theorem

There is an optimal schedule with no idle time and no inversions.

Immediate Inversion

$$d_j < d_i$$

$$l'_j \leq l_j$$




$$l'_i = f_j - d_i < f_j - d_j = l_j$$




Fractional Knapsack Problem (Problem 16.2-1)

Prove that the fractional knapsack problem has the greedy-choice property.



Fractional Knapsack Problem (Problem 16.2-1)


Prove that the fractional knapsack problem has the greedy-choice property.



\exists an optimal solution which contains the greedy choice.

Fractional Knapsack Problem (Problem 16.2-1)

Prove that the fractional knapsack problem has the greedy-choice property.




\exists an optimal solution which contains the greedy choice.

$$\frac{v_1}{w_1} \geq \frac{v_2}{w_2} \geq \dots \geq \frac{v_n}{w_n}$$

Fractional Knapsack Problem (Problem 16.2-1)

Prove that the fractional knapsack problem has the greedy-choice property.



\exists an optimal solution which contains the greedy choice.

$$\frac{v_1}{w_1} \geq \frac{v_2}{w_2} \geq \dots \geq \frac{v_n}{w_n}$$

As most item of $\frac{v_1}{w_1}$ as possible.

Change-Making Problem (Problem 16-1 (a))



Making change for v cents using the fewest number of coins.

Change-Making Problem (Problem 16-1 (a))



25



10



5



1

Making change for v cents using the fewest number of coins.

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

$$G : v = 25 \times A + 10 \times B + 5 \times C + 1 \times D$$

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

$$\mathsf{G} : v = 25 \times A + 10 \times B + 5 \times C + 1 \times D$$

(A^*, B^*, C^*, D^*) vs. (A, B, C, D)

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

$$\mathsf{G} : v = 25 \times A + 10 \times B + 5 \times C + 1 \times D$$

(A^*, B^*, C^*, D^*) vs. (A, B, C, D)

$$A^* = A$$

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

Lemma (Properties of OPT)

- (I) $B^* \leq 2$
- (II) $C^* \leq 1$
- (III) $D^* \leq 4$
- (IV) $\neg(B^* = 2 \wedge C^* = 1)$

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

Lemma (Properties of OPT)

- (I) $B^* \leq 2$
- (II) $C^* \leq 1$
- (III) $D^* \leq 4$
- (IV) $\neg(B^* = 2 \wedge C^* = 1)$

Theorem

$$10 \times B^* + 5 \times C^* + 1 \times D^* \leq 24$$

$$5 \times C^* + 1 \times D^* \leq 9$$

$$1 \times D^* \leq 4$$

$$\text{OPT} : v = 25 \times A^* + 10 \times B^* + 5 \times C^* + 1 \times D^*$$

Lemma (Properties of OPT)

- (I) $B^* \leq 2$
- (II) $C^* \leq 1$
- (III) $D^* \leq 4$
- (IV) $\neg(B^* = 2 \wedge C^* = 1)$

Theorem

$$10 \times B^* + 5 \times C^* + 1 \times D^* \leq 24$$

$$5 \times C^* + 1 \times D^* \leq 9$$

$$1 \times D^* \leq 4$$

$$(A^*, B^*, C^*, D^*) = (A, B, C, D)$$

Change-Making Problem (Problem 16-1 (c))



Change-Making Problem (Problem 16-1 (c))



$$\{1, 3, 4\}, \quad v = 6$$

Change-Making Problem (Problem 16-1 (c))



$$\{1, 3, 4\}, \quad v = 6$$

G : 4, 1, 1 **vs.** OPT : 3, 3

Change-Making Problem (Problem 16-1 (c))



$$\{1, 3, 4\}, \quad v = 6$$

$$G : 4, 1, 1 \quad vs. \quad OPT : 3, 3$$

Why does the previous proof fail here?

Change-Making Problem (Problem 16-1 (b))

$$c^0, c^1, c^2, \dots, c^k, \quad c > 1, k \geq 1$$

Change-Making Problem (Problem 16-1 (b))

$$c^0, c^1, c^2, \dots, c^k, \quad c > 1, k \geq 1$$

$$\text{OPT} : v = \sum_{i=0}^{i=k} c^i \color{red}{a_i^*}$$

Change-Making Problem (Problem 16-1 (b))

$$c^0, c^1, c^2, \dots, c^k, \quad c > 1, k \geq 1$$

$$\text{OPT} : v = \sum_{i=0}^{i=k} c^i \color{red}{a_i^*}$$

$$\text{G} : v = \sum_{i=0}^{i=k} c^i \color{blue}{a_i}$$

Change-Making Problem (Problem 16-1 (b))

$$c^0, c^1, c^2, \dots, c^k, \quad c > 1, k \geq 1$$

$$\text{OPT} : v = \sum_{i=0}^{i=k} c^i \color{red}{a_i^*}$$


$$\text{G} : v = \sum_{i=0}^{i=k} c^i \color{blue}{a_i}$$

$$\color{red}{a_i^*} = \color{blue}{a_i}$$


Canonical Coin Systems



Minimum Shots Problem (Additional Problem)



Minimum Shots Problem (Additional Problem)




Assumption:

Possible to shoot a ray that does not intersect any balloons.







Office 302

Mailbox: H016

hfwei@nju.edu.cn