


# 1-10 Set Theory (III): Functions

魏恒峰

hfwei@nju.edu.cn

2019 年 12 月 10 日





# Functions

# Functions



PROOF!

# Definition of Functions

$$R \subseteq A \times B$$

is a *relation* from  $A$  to  $B$

## Definition (Function)

$R \subseteq A \times B$  is a *function* from  $A$  to  $B$  if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

## Definition (Function)

$R \subseteq A \times B$  is a *function* from  $A$  to  $B$  if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

$$f : A \rightarrow B$$



## Definition (Function)

$R \subseteq A \times B$  is a *function* from  $A$  to  $B$  if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

$$f : A \rightarrow B$$

$$\text{dom}(f) = A \quad \text{cod}(f) = B$$

$$\text{ran}(f) = f(A) \subseteq B$$

## Definition (Function)

$R \subseteq A \times B$  is a *function* from  $A$  to  $B$  if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

$$f : A \rightarrow B$$

$$\text{dom}(f) = A \quad \text{cod}(f) = B$$

$$\text{ran}(f) = f(A) \subseteq B$$

$$f : a \mapsto b$$

$$f(a) \triangleq b$$

## Definition (Function)

$R \subseteq A \times B$  is a *function* from  $A$  to  $B$  if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

## Definition (Function)

$R \subseteq A \times B$  is a *function* from  $A$  to  $B$  if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

$$\forall a \in A :$$

$$\forall a \in A : \exists b \in B : (a, b) \in f$$

## Definition (Function)

$R \subseteq A \times B$  is a *function* from  $A$  to  $B$  if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

$$\forall a \in A :$$

$$\forall a \in A : \exists b \in B : (a, b) \in f$$

$$\exists! b \in B :$$

$$\forall b, b' \in B : (a, b) \in f \wedge (a, b') \in f \implies b = b'$$

## Definition

The *set* of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

## Definition

The *set* of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$Y^X = \{f \in \mathcal{P}(X \times Y) \mid f : X \rightarrow Y\}$$

## Definition

The *set* of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$Y^X = \{f \in \mathcal{P}(X \times Y) \mid f : X \rightarrow Y\}$$

$X$  and  $Y$  are finite sets with  $x$  and  $y$  elements, respectively.

$$|X| = x \quad |Y| = y, \quad |Y^X| =$$



## Definition

The *set* of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$Y^X = \{f \in \mathcal{P}(X \times Y) \mid f : X \rightarrow Y\}$$

$X$  and  $Y$  are finite sets with  $x$  and  $y$  elements, respectively.

$$|X| = x \quad |Y| = y, \quad |Y^X| = y^x$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset =$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

$$\emptyset^\emptyset = \{\emptyset\}$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

$$\emptyset^\emptyset = \{\emptyset\}$$

$$\forall X \neq \emptyset : \emptyset^X =$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

$$\emptyset^\emptyset = \{\emptyset\}$$

$$\forall X \neq \emptyset : \emptyset^X = \emptyset$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$2^X = \{0, 1\}^X \cong \mathcal{P}(X)$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$2^X = \{0, 1\}^X \cong \mathcal{P}(X)$$

$$\mathcal{P}(\{\text{🍏 🍌}\}) = \left\{ \left\{ \begin{array}{l} \text{🍏 🍌} \\ \text{🍏} \\ \text{🍌} \\ \end{array} \right\} \right\} \cong \left\{ \begin{array}{ll} \text{in} & \text{in} \\ \text{in} & \text{out} \\ \text{out} & \text{in} \\ \text{out} & \text{out} \end{array} \right\}$$



## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$Q$  : Is there a set consisting of all functions?

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$Q$  : Is there a set consisting of all functions?

## Theorem

*There is no set consisting of all functions.*

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$Q$  : Is there a set consisting of all functions?

## Theorem

*There is no set consisting of all functions.*

Suppose **by contradiction** that  $A$  is the set of all functions.

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$Q$  : Is there a set consisting of all functions?

## Theorem

*There is no set consisting of all functions.*

Suppose **by contradiction** that  $A$  is the set of all functions.

For every set  $X$ , there exists a function  $I_X : \{X\} \rightarrow \{X\}$ .

## Definition

The set of all functions from  $X$  to  $Y$ :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$Q$  : Is there a set consisting of all functions?

## Theorem

*There is no set consisting of all functions.*

Suppose **by contradiction** that  $A$  is the set of all functions.

For every set  $X$ , there exists a function  $I_X : \{X\} \rightarrow \{X\}$ .

$$\bigcup_{I_X \in A} \text{dom}(I_X)$$

# Functions as Sets

## Axiom (Axiom of Extensionality)

$$\forall A \forall B \forall x : (x \in A \iff x \in B) \iff A = B.$$



## Axiom (Axiom of Extensionality)

$$\forall A \forall B \forall x : (x \in A \iff x \in B) \iff A = B.$$

## Theorem (The Principle of Functional Extensionality)

*f, g are functions :*

$$f = g \iff \text{dom}(f) = \text{dom}(g) \wedge \left( \forall x \in \text{dom}(f) : f(x) = g(x) \right)$$

## Axiom (Axiom of Extensionality)

$$\forall A \forall B \forall x : (x \in A \iff x \in B) \iff A = B.$$

## Theorem (The Principle of Functional Extensionality)

*f, g are functions :*

$$f = g \iff \text{dom}(f) = \text{dom}(g) \wedge \left( \forall x \in \text{dom}(f) : f(x) = g(x) \right)$$

$$\forall f \forall g \forall (a, b) : ((a, b) \in f \iff (a, b) \in g) \iff f = g.$$

## Axiom (Axiom of Extensionality)

$$\forall A \forall B \forall x : (x \in A \iff x \in B) \iff A = B.$$

## Theorem (The Principle of Functional Extensionality)

*f, g are functions :*

$$f = g \iff \text{dom}(f) = \text{dom}(g) \wedge \left( \forall x \in \text{dom}(f) : f(x) = g(x) \right)$$

$$\forall f \forall g \forall (a, b) : ((a, b) \in f \iff (a, b) \in g) \iff f = g.$$

It may be that  $\text{cod}(f) \neq \text{cod}(g)$ .

$$f : A \rightarrow B \quad g : C \rightarrow D$$

$Q$  : Is  $f \cap g$  a function?

$$f : A \rightarrow B \quad g : C \rightarrow D$$

$Q$  : Is  $f \cap g$  a function?

Theorem (Intersection of Functions)

$$f \cap g : (A \cap C) \rightarrow (B \cap D)$$

$$f : A \rightarrow B \quad g : C \rightarrow D$$

$Q$  : Is  $f \cup g$  a function?

$$f : A \rightarrow B \quad g : C \rightarrow D$$

$Q$  : Is  $f \cup g$  a function?

Theorem (Union of Functions)

$$f \cup g : (A \cup C) \rightarrow (B \cup D) \iff \forall x \in \text{dom}(f) \cap \text{dom}(g) : f(x) = g(x)$$

$$f : A \rightarrow B \quad g : C \rightarrow D$$

$Q$  : Is  $f \cup g$  a function?

Theorem (Union of Functions)

$$f \cup g : (A \cup C) \rightarrow (B \cup D) \iff \forall x \in \text{dom}(f) \cap \text{dom}(g) : f(x) = g(x)$$

UD Problem 14.3 (g)

$$f : \mathbb{Q} \rightarrow \mathbb{R}$$

$$f(x) = \begin{cases} x + 1 & \text{if } x \in 2\mathbb{Z} \\ x - 1 & \text{if } x \in 3\mathbb{Z} \\ 2 & \text{otherwise} \end{cases}$$



$$f : A \rightarrow B \quad g : C \rightarrow D$$

$Q$  : Is  $f \cup g$  a function?

Theorem (Union of Functions)

$$f \cup g : (A \cup C) \rightarrow (B \cup D) \iff \forall x \in \text{dom}(f) \cap \text{dom}(g) : f(x) = g(x)$$

UD Problem 14.3 (g)

$$f : \mathbb{Q} \rightarrow \mathbb{R}$$

$$f(x) = \begin{cases} x + 1 & \text{if } x \in 2\mathbb{Z} \\ x - 1 & \text{if } x \in 3\mathbb{Z} \\ 2 & \text{otherwise} \end{cases}$$

$$x \in 6\mathbb{Z}$$

$$D : \mathbb{R} \rightarrow \mathbb{R}$$

$$D(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$$

Dirichlet Function

# Special Functions (*-jectivity*)

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B \quad f : A \rightarrowtail B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B \quad f : A \rightarrowtail B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

For Proof:

► To prove that  $f$  *is* 1-1:

$$\forall a_1, a_2 \in A : f(a_1) = f(a_2) \implies a_1 = a_2$$

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B \quad f : A \rightarrowtail B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

For Proof:

► To prove that  $f$  *is* 1-1:

$$\forall a_1, a_2 \in A : f(a_1) = f(a_2) \implies a_1 = a_2$$

► To show that  $f$  *is not* 1-1:

$$\exists a_1, a_2 \in A : a_1 \neq a_2 \wedge f(a_1) = f(a_2)$$

## Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B$$

$$\text{ran}(f) = B$$



## Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B \quad f : A \twoheadrightarrow B$$

$$\text{ran}(f) = B$$

## Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B \quad f : A \twoheadrightarrow B$$

$$\text{ran}(f) = B$$

For Proof:

► To prove that  $f$  *is* onto:

$$\forall b \in B \left( \exists a \in A : f(a) = b \right)$$

## Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B \quad f : A \twoheadrightarrow B$$

$$\text{ran}(f) = B$$

For Proof:

► To prove that  $f$  *is* onto:

$$\forall b \in B \left( \exists a \in A : f(a) = b \right)$$

► To show that  $f$  *is not* onto:

$$\exists b \in B \left( \forall a \in A : f(a) \neq b \right)$$

Definition (Bijective (one-to-one correspondence) 一一对应)

$$f : A \rightarrow B$$

1-1 & onto

Definition (Bijective (one-to-one correspondence) 一一对应)

$$f : A \rightarrow B \quad f : A \overset{1-1}{\underset{\text{onto}}{\longleftrightarrow}} B$$

1-1 & onto

# Functions as Relations

$$f|_X \quad f(A) \quad f^{-1}(B) \quad f^{-1} \quad f \circ g$$

## Definition (Restriction)

The *restriction* of a function  $f$  to  $X$  is the **function**:

$$f|_X = \{(x, y) \in f \mid x \in X\}$$

## Definition (Restriction)

The *restriction* of a function  $f$  to  $X$  is the **function**:

$$f|_X = \{(x, y) \in f \mid x \in X\}$$

$$f : A \rightarrow B$$



## Definition (Restriction)

The *restriction* of a function  $f$  to  $X$  is the **function**:

$$f|_X = \{(x, y) \in f \mid x \in X\}$$

$$f : A \rightarrow B$$

$$f|_X : A \cap X \rightarrow B$$

$$f|_X(x) = f(x), \forall x \in A \cap X$$

## Definition (Image)

The *image* of  $X$  under a function  $f$  is the **set**

$$f(X) = \{b \mid \exists a \in X : (a, b) \in f\}$$

## Definition (Inverse Image)

The *inverse image* of  $Y$  under a function  $f$  is the **set**

$$f^{-1}(Y) = \{a \mid \exists b \in Y : (a, b) \in f\}$$

## Definition (Image)

The *image* of  $X$  under a function  $f$  is the **set**

$$f(X) = \{b \mid \exists a \in X : (a, b) \in f\}$$

## Definition (Inverse Image)

The *inverse image* of  $Y$  under a function  $f$  is the **set**

$$f^{-1}(Y) = \{a \mid \exists b \in Y : (a, b) \in f\}$$

$X \subseteq \text{dom}(f)$ ,  $Y \subseteq \text{ran}(f)$  are not necessary

## Definition (Image)

The *image* of  $X$  under a function  $f$  is the **set**

$$f(X) = \{b \mid \exists a \in X : (a, b) \in f\}$$

## Definition (Inverse Image)

The *inverse image* of  $Y$  under a function  $f$  is the **set**

$$f^{-1}(Y) = \{a \mid \exists b \in Y : (a, b) \in f\}$$

$X \subseteq \text{dom}(f)$ ,  $Y \subseteq \text{ran}(f)$  are not necessary

$f$  may not be **invertible** in  $f^{-1}(Y)$

$$y \in f(X) \iff \exists x \in \text{dom}(f) \cap X : y = f(x)$$

$$y \in f(X) \iff \exists x \in X : y = f(x)$$

$$x \in f^{-1}(Y) \iff f(x) \in Y$$

## Theorem (Properties of $f$ and $f^{-1}$ (UD Theorem 17.7))

$$f : A \rightarrow B \quad A_1, A_2 \subseteq A, \quad B_1, B_2 \subseteq B$$

(i)  $f$  preserves only  $\subseteq$  and  $\cup$ :

$$(1) \quad A_1 \subseteq A_2 \implies f(A_1) \subseteq f(A_2)$$

$$(2) \quad f(A_1 \cup A_2) = f(A_1) \cup f(A_2)$$

$$(3) \quad f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$(4) \quad f(A_1 \setminus A_2) \supseteq f(A_1) \setminus f(A_2)$$

(ii)  $f^{-1}$  preserves  $\subseteq, \cup, \cap$ , and  $\setminus$ :

$$(5) \quad B_1 \subseteq B_2 \implies f^{-1}(B_1) \subseteq f^{-1}(B_2)$$

$$(6) \quad f^{-1}(B_1 \cup B_2) = f^{-1}(B_1) \cup f^{-1}(B_2)$$

$$(7) \quad f^{-1}(B_1 \cap B_2) = f^{-1}(B_1) \cap f^{-1}(B_2)$$

$$(8) \quad f^{-1}(B_1 \setminus B_2) = f^{-1}(B_1) \setminus f^{-1}(B_2)$$

## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$b \in f(A_1 \cap A_2)$$



## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$b \in f(A_1 \cap A_2)$$

$$\implies \exists a \in A_1 \cap A_2 \cap A : b = f(a)$$

## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$b \in f(A_1 \cap A_2)$$

$$\implies \exists a \in A_1 \cap A_2 : b = f(a)$$

$$\implies \exists a \in A : a \in A_1 \wedge a \in A_2 \wedge b = f(a)$$

## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$b \in f(A_1 \cap A_2)$$

$$\implies \exists a \in A_1 \cap A_2 : b = f(a)$$

$$\implies \exists a \in A : a \in A_1 \wedge a \in A_2 \wedge b = f(a)$$

$$\implies \exists a \in A \cap A_1 : b = f(a) \wedge \exists a \in A \cap A_2 : b = f(a)$$

## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$b \in f(A_1 \cap A_2)$$

$$\implies \exists a \in A_1 \cap A_2 : b = f(a)$$

$$\implies \exists a \in A : a \in A_1 \wedge a \in A_2 \wedge b = f(a)$$

$$\implies \exists a \in A \cap A_1 : b = f(a) \wedge \exists a \in A \cap A_2 : b = f(a)$$

$$\implies b \in f(A_1) \cap f(A_2)$$

## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$b \in f(A_1 \cap A_2)$$

$$\implies \exists a \in A_1 \cap A_2 : b = f(a)$$

$$\implies \exists a \in A : a \in A_1 \wedge a \in A_2 \wedge b = f(a)$$

$$\implies \exists a \in A \cap A_1 : b = f(a) \wedge \exists a \in A \cap A_2 : b = f(a)$$

$$\implies b \in f(A_1) \cap f(A_2)$$

*Q* : When does  $f(A_1 \cap A_2) = f(A_1) \cap f(A_2)$  hold?

## Theorem (UD Problem 17.5)

$$f : A \rightarrow B$$

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

$$b \in f(A_1 \cap A_2)$$

$$\implies \exists a \in A_1 \cap A_2 : b = f(a)$$

$$\implies \exists a \in A : a \in A_1 \wedge a \in A_2 \wedge b = f(a)$$

$$\implies \exists a \in A \cap A_1 : b = f(a) \wedge \exists a \in A \cap A_2 : b = f(a)$$

$$\implies b \in f(A_1) \cap f(A_2)$$

$Q$  : When does  $f(A_1 \cap A_2) = f(A_1) \cap f(A_2)$  hold?

$f$  is injective.

Theorem (Properties of  $f$  and  $f^{-1}$  (UD Theorem 17.7))

$$f : A \rightarrow B$$

(iii)  $f$  and  $f^{-1}$ :

$$(9) \quad A_0 \subseteq A \implies A_0 \subseteq f^{-1}(f(A_0))$$

$$(10) \quad B_0 \supseteq f(f^{-1}(B_0))$$

Theorem (Properties of  $f$  and  $f^{-1}$  (UD Theorem 17.7))

$$f : A \rightarrow B$$

(iii)  $f$  and  $f^{-1}$ :

$$(9) \quad A_0 \subseteq A \implies A_0 \subseteq f^{-1}(f(A_0))$$

$$(10) \quad B_0 \supseteq f(f^{-1}(B_0))$$

Theorem (UD Problem 17.8)

$$A_0 \subseteq A \implies A_0 \subseteq f^{-1}(f(A_0))$$



## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

$$b \in f(f^{-1}(B_0))$$

## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

$$\begin{aligned} b &\in f(f^{-1}(B_0)) \\ \implies \exists a \in f^{-1}(B_0) : b &= f(a) \end{aligned}$$

## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

$$b \in f(f^{-1}(B_0))$$

$$\implies \exists a \in f^{-1}(B_0) : b = f(a)$$

$$\implies \exists a \in A : f(a) \in B_0 \wedge b = f(a)$$

## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

$$b \in f(f^{-1}(B_0))$$

$$\implies \exists a \in f^{-1}(B_0) : b = f(a)$$

$$\implies \exists a \in A : f(a) \in B_0 \wedge b = f(a)$$

$$\implies b \in B_0$$

## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

$$b \in f(f^{-1}(B_0))$$

$$\implies \exists a \in f^{-1}(B_0) : b = f(a)$$

$$\implies \exists a \in A : f(a) \in B_0 \wedge b = f(a)$$

$$\implies b \in B_0$$

*Q:* When does  $B_0 = f(f^{-1}(B_0))$  hold?

## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

$$b \in f(f^{-1}(B_0))$$

$$\implies \exists a \in f^{-1}(B_0) : b = f(a)$$

$$\implies \exists a \in A : f(a) \in B_0 \wedge b = f(a)$$

$$\implies b \in B_0$$

*Q:* When does  $B_0 = f(f^{-1}(B_0))$  hold?

*f* is surjective and  $B_0 \subseteq B$ .

## Theorem

$$f : A \rightarrow B$$

$$B_0 \supseteq f(f^{-1}(B_0))$$

$$b \in f(f^{-1}(B_0))$$

$$\implies \exists a \in f^{-1}(B_0) : b = f(a)$$

$$\implies \exists a \in A : f(a) \in B_0 \wedge b = f(a)$$

$$\implies b \in B_0$$


*Q:* When does  $B_0 = f(f^{-1}(B_0))$  hold?

$f$  is surjective and  $B_0 \subseteq B$ .

$$B_0 \subseteq \text{ran}(f)$$



# Function Composition



## Definition (Composition)

$$f : A \rightarrow B \quad g : C \rightarrow D$$

$$\text{ran}(f) \subseteq C$$

The *composite function*  $g \circ f : A \rightarrow D$  is defined as

$$(g \circ f)(x) = g(f(x))$$

## Definition (Composition)

$$f : A \rightarrow B \quad g : C \rightarrow D$$

$$\text{ran}(f) \subseteq C$$

The *composite function*  $g \circ f : A \rightarrow D$  is defined as

$$(g \circ f)(x) = g(f(x))$$

Why not “ $\exists b$ ” as below?

## Definition (Composition)

The *composition* of relations  $R$  and  $S$  is the relation

$$R \circ S = \{(a, c) \mid \exists b : (a, b) \in S \wedge (b, c) \in R\}$$

## Theorem (Associative Property for Composition)

$$f : A \rightarrow B \quad g : B \rightarrow C \quad h : C \rightarrow D$$

$$h \circ (g \circ f) = (h \circ g) \circ f$$

## Theorem (Associative Property for Composition)

$$f : A \rightarrow B \quad g : B \rightarrow C \quad h : C \rightarrow D$$

$$h \circ (g \circ f) = (h \circ g) \circ f$$

Proof.

## Theorem (Associative Property for Composition)

$$f : A \rightarrow B \quad g : B \rightarrow C \quad h : C \rightarrow D$$

$$h \circ (g \circ f) = (h \circ g) \circ f$$

Proof.

(i)

$$\text{dom}(h \circ (g \circ f)) = \text{dom}((h \circ g) \circ f)$$

(ii)

$$(h \circ (g \circ f))(x) = ((h \circ g) \circ f)(x)$$



## Theorem (UD Theorem 16.7)

$$f : A \rightarrow B \quad g : B \rightarrow C$$

- (i) *If  $f, g$  are injective, then  $g \circ f$  is injective.*
- (ii) *If  $f, g$  are surjective, then  $g \circ f$  is surjective.*
- (iii) *If  $f, g$  are bijective, then  $g \circ f$  is bijective.*

## Theorem (UD Theorem 16.7)

$$f : A \rightarrow B \quad g : B \rightarrow C$$

- (i) *If  $f, g$  are injective, then  $g \circ f$  is injective.*
- (ii) *If  $f, g$  are surjective, then  $g \circ f$  is surjective.*
- (iii) *If  $f, g$  are bijective, then  $g \circ f$  is bijective.*

Proof for (i).

$$\forall a_1, a_2 \in A : ((g \circ f)(a_1) = (g \circ f)(a_2) \implies a_1 = a_2)$$





## Theorem (UD Theorem 16.7)

$$f : A \rightarrow B \quad g : B \rightarrow C$$

- (i) *If  $f, g$  are injective, then  $g \circ f$  is injective.*
- (ii) *If  $f, g$  are surjective, then  $g \circ f$  is surjective.*
- (iii) *If  $f, g$  are bijective, then  $g \circ f$  is bijective.*

Proof for (i).

$$\forall a_1, a_2 \in A : ((g \circ f)(a_1) = (g \circ f)(a_2) \implies a_1 = a_2)$$



Proof for (ii).

$$\forall c \in C : (\exists a \in A : (g \circ f)(a) = c)$$



## Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow C$$

- (i) *If  $g \circ f$  is surjective, then  $g$  is surjective.*
- (ii) *If  $g \circ f$  is injective, then  $f$  is injective.*


## Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow C$$

- (i) *If  $g \circ f$  is surjective, then  $g$  is surjective.*
- (ii) *If  $g \circ f$  is injective, then  $f$  is injective.*

You can also prove it by contradiction.

# Inverse Functions



## Definition (Inverse)

Let  $f : A \rightarrow B$  be a **bijective** function.

The *inverse* of  $f$  is the **function**  $f^{-1} : B \rightarrow A$  defined by

$$f^{-1}(b) = a \iff f(a) = b.$$

## Definition (Inverse)

Let  $f : A \rightarrow B$  be a **bijective** function.

The *inverse* of  $f$  is the **function**  $f^{-1} : B \rightarrow A$  defined by

$$f^{-1}(b) = a \iff f(a) = b.$$



信息量太大

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$f$  is invertible  $\iff f$  is bijective.



## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$f$  is invertible  $\iff f$  is bijective.

$f$  is invertible  $\implies f$  is bijective

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$f$  is invertible  $\iff f$  is bijective.

$f$  is invertible  $\implies f$  is bijective

$g$  is a function  $\implies f$  is injective

$\text{dom}(g) = Y \implies f$  is surjective

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$f$  is invertible  $\iff f$  is bijective.

$f$  is invertible  $\implies f$  is bijective

$f$  is bijective  $\implies f$  is invertible

$g$  is a function  $\implies f$  is injective

$\text{dom}(g) = Y \implies f$  is surjective

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$f$  is invertible  $\iff f$  is bijective.

$f$  is invertible  $\implies f$  is bijective

$g$  is a function  $\implies f$  is injective

$\text{dom}(g) = Y \implies f$  is surjective

$f$  is bijective  $\implies f$  is invertible

To show that  $g$  defined above is indeed a function from  $Y$  to  $X$ .

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$g : Y \rightarrow X$  is *unique*.

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$g : Y \rightarrow X$  is *unique*.

By Contradiction

## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$g : Y \rightarrow X$  is *unique*.

By Contradiction

$$f^{-1} \triangleq g$$



## Definition (Invertible)

$f : X \rightarrow Y$  is *invertible* if there exists  $g : Y \rightarrow X$  such that

$$f(x) = y \iff g(y) = x.$$

## Theorem

$g : Y \rightarrow X$  is *unique*.

By Contradiction

$$f^{-1} \triangleq g$$

$$f(x) = y \iff f^{-1}(y) = x$$

## Theorem (UD Theorem 16.4)

*$f : A \rightarrow B$  is bijective*

- (i)  $f \circ f^{-1} = I_B$
- (ii)  $f^{-1} \circ f = I_A$
- (iii)  $f^{-1}$  is bijective.
- (iv)  $g : B \rightarrow A \wedge f \circ g = I_B \implies g = f^{-1}$
- (v)  $g : B \rightarrow A \wedge g \circ f = I_A \implies g = f^{-1}$

## Theorem (UD Theorem 16.4)

*$f : A \rightarrow B$  is bijective*

- (i)  $f \circ f^{-1} = I_B$
- (ii)  $f^{-1} \circ f = I_A$
- (iii)  $f^{-1}$  is bijective.
- (iv)  $g : B \rightarrow A \wedge f \circ g = I_B \implies g = f^{-1}$
- (v)  $g : B \rightarrow A \wedge g \circ f = I_A \implies g = f^{-1}$

The ways to find/check  $f^{-1}$ .

## Theorem (UD Theorem 16.4)

*$f : A \rightarrow B$  is bijective*

- (i)  $f \circ f^{-1} = I_B$
- (ii)  $f^{-1} \circ f = I_A$
- (iii)  $f^{-1}$  is bijective.
- (iv)  $g : B \rightarrow A \wedge f \circ g = I_B \implies g = f^{-1}$
- (v)  $g : B \rightarrow A \wedge g \circ f = I_A \implies g = f^{-1}$

The ways to find/check  $f^{-1}$ .

$$g = f^{-1} \circ (f \circ g) = f^{-1} \circ I_B = f^{-1}$$

## Theorem (Inverse of Composition (UD Theorem 16.6))

$f : A \rightarrow B$   $g : B \rightarrow C$  are bijective

- (i)  $g \circ f$  is bijective
- (ii)  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$

Proof for (ii).

It suffices to check either one of the following identities:

$$(f^{-1} \circ g^{-1}) \circ (g \circ f) = I_A$$

$$(g \circ f) \circ (f^{-1} \circ g^{-1}) = I_C$$



## Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow A$$

$$(iii) \quad f \circ g = I_B \wedge g \circ f = I_A \implies g = f^{-1}$$

Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow A$$

$$(iii) \quad f \circ g = I_B \wedge g \circ f = I_A \implies g = f^{-1}$$

You need to check **both** identities.

### Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow A$$

$$(iii) \quad f \circ g = I_B \wedge g \circ f = I_A \implies g = f^{-1}$$

You need to check **both** identities.

### Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow C$$

- (i) *If  $g \circ f$  is surjective, then  $g$  is surjective.*
- (ii) *If  $g \circ f$  is injective, then  $f$  is injective.*



Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow A$$

$$(iii) \quad f \circ g = I_B \wedge g \circ f = I_A \implies g = f^{-1}$$

You need to check **both** identities.

Theorem (UD Theorem 16.8)

$$f : A \rightarrow B \quad g : B \rightarrow C$$

- (i) *If  $g \circ f$  is surjective, then  $g$  is surjective.*
- (ii) *If  $g \circ f$  is injective, then  $f$  is injective.*

First show that  $f$  is bijective, and then use Theorem 16.4.

Thank  
You!