


1-11 Set Theory (IV): Infinity


魏恒峰

hfwei@nju.edu.cn


2019 年 12 月 17 日








Georg Cantor (1845 – 1918)



Georg Cantor (1845 – 1918)



Leopold Kronecker
(1823 – 1891)




Georg Cantor (1845 – 1918)



Leopold Kronecker
(1823 – 1891)



Henri Poincaré
(1854 – 1912)



Georg Cantor (1845 – 1918)



Leopold Kronecker
(1823 – 1891)




Henri Poincaré
(1854 – 1912)



Ludwig Wittgenstein

◀ ◻ ▶ ◀ ◻ ▶ (1889 – 1951) ≡ 🔍 🔍 🔍



Georg Cantor (1845 – 1918)



David Hilbert (1862 – 1943)



Leopold Kronecker
(1823 – 1891)



Henri Poincaré
(1854 – 1912)



Ludwig Wittgenstein
(1889 – 1951)

*From his paradise that Cantor with us unfolded, we hold our
breath in awe; knowing, we shall not be expelled.*

— *David Hilbert*

“没有人能把我们从 Cantor 创造的乐园中驱逐出去”

*From his paradise that Cantor with us unfolded, we hold our
breath in awe; knowing, we shall not be expelled.*

— *David Hilbert*

“没有人能把我们从 Cantor 创造的乐园中驱逐出去”








“das wesen der mathematik liegt in ihrer freiheit”



“das wesen der mathematik liegt in ihrer freiheit”

“The essence of mathematics lies in its freedom”


Before Cantor








公理: “整体大于部分”



Galileo Galilei (1564 – 1642)



“关于两门新科学的对话” (1638)



Galileo Galilei (1564 – 1642)

“关于两门新科学的对话” (1638)

“用我们有限的心智来讨论无限...”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”



吓得我吃了一鲸

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”



吓得我吃了一鲸

说到底，“等于”、“大于”和“小于”诸性质不能用于无限，而只能用于有限的数量。
— Galileo Galilei

$$S_1 = \{1, 2, 3, \dots, n, \dots\}$$

$$S_2 = \{1, 4, 9, \dots, n^2, \dots\}$$

$$|S_1| = |S_2| \quad S_2 \subset S_1$$

“部分等于全体”



吓得我吃了一鲸

说到底，“等于”、“大于”和“小于”诸性质不能用于无限，而只能用于有限的数量。
— Galileo Galilei

无穷数是不可能的。
— Gottfried Wilhelm Leibniz

这些证明一开始就期望那些数要具有有穷数的一切性质，或者甚至于把有穷数的性质强加于无穷。

相反，这些无穷数，如果它们能够以任何形式被理解的话，倒是由于它们与有穷数的对应，它们必须具有完全新的数量特征。

这些性质完全依赖于事物的本性， \dots 而并非来自我们的主观任意性或我们的偏见。

— Georg Cantor (1885)

Definition (Dedekind-infinite & Dedekind-finite (Dedekind, 1888))


A set A is *Dedekind-infinite* if there is a bijective function from A onto some proper subset B of A .

A set is *Dedekind-finite* if it is not Dedekind-infinite.

Definition (Dedekind-infinite & Dedekind-finite (Dedekind, 1888))

A set A is *Dedekind-infinite* if there is a bijective function from A onto some proper subset B of A .

A set is *Dedekind-finite* if it is not Dedekind-infinite.



We will prove this as a **theorem** in our theory of infinity.



We have not defined “finite” and “infinite”!!!

Comparing Sets

Comparing Sets



Comparing Sets



Function



Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

$\overline{\overline{A}}$ (two *abstractions*)

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

$\overline{\overline{A}}$ (two *abstractions*)

Abstract from elements: $\{1, 2, 3\}$ vs. $\{a, b, c\}$

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

$\overline{\overline{A}}$ (two *abstractions*)

Abstract from elements: $\{1, 2, 3\}$ vs. $\{a, b, c\}$

Abstract from order: $\{1, 2, 3, \dots\}$ vs. $\{1, 3, 5, \dots, 2, 4, 6, \dots\}$

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Q : Is “ \approx ” an equivalence relation?

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Q : Is “ \approx ” an equivalence relation?

Theorem ()

For any sets A, B, C :

- (a) $A \approx B$
- (b) $A \approx B \implies B \approx A$
- (c) $A \approx B \wedge B \approx C \implies A \approx C$

Definition ($|A| = |B|$ ($A \approx B$) (1878))

A and B are *equipotent* if there exists a *bijection* from A to B .

Q : Is “ \approx ” an equivalence relation?

Theorem (The “Equivalence Concept” of Equipotent)

For any sets A, B, C :

- (a) $A \approx B$
- (b) $A \approx B \implies B \approx A$
- (c) $A \approx B \wedge B \approx C \implies A \approx C$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n.$$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n.$$

$$|X| = |\{0, 1, \dots, n-1\}|$$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n.$$

$$|X| = |\{0, 1, \dots, n-1\}|$$

Theorem (UD Theorem 22.6)

*Let A be a finite set. There is a **unique** $n \in \mathbb{N}$ such that $A \approx \{0, 1, \dots, n-1\}$.*

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem (UD Theorem 22.3)

\mathbb{N} is infinite.

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem (UD Theorem 22.3)

\mathbb{N} is infinite.

By Contradiction.

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem (UD Theorem 22.3)

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem (UD Theorem 22.3)

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem (UD Theorem 22.3)

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

$$g \triangleq f|_{\{0,1,\dots,n\}} : \{0, 1, \dots, n\} \rightarrow \{0, 1, \dots, n-1\}$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem (UD Theorem 22.3)

\mathbb{N} is infinite.

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

$$g \triangleq f|_{\{0,1,\dots,n\}} : \{0, 1, \dots, n\} \rightarrow \{0, 1, \dots, n-1\}$$

$$g \text{ is not 1-1} \implies f \text{ is not 1-1}$$

Definition (Infinite)

X is infinite if it is not finite:

$$\forall n \in \mathbb{N} : |X| \neq n.$$

Theorem (UD Theorem 22.3)

\mathbb{N} is infinite. (So are \mathbb{Z} , \mathbb{Q} , \mathbb{R} .)

By Contradiction.

$$\exists n \in \mathbb{N} : |\mathbb{N}| = n.$$

$$\exists f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \{0, 1, \dots, n-1\}$$

$$g \triangleq f|_{\{0,1,\dots,n\}} : \{0, 1, \dots, n\} \rightarrow \{0, 1, \dots, n-1\}$$

$$g \text{ is not 1-1} \implies f \text{ is not 1-1}$$

Definition (Infinite)

For any set X ,

Countably Infinite

$$|X| = |\mathbb{N}| \triangleq \aleph_0$$

Countable

(finite \vee countably infinite)

Uncountable


(\neg countable)

(infinite) \wedge (\neg (countably infinite))

\aleph_0


Theorem (\mathbb{Z} is Countable.)

$$|\mathbb{Z}| = |\mathbb{N}|$$



Theorem (\mathbb{Z} is Countable.)

$$|\mathbb{Z}| = |\mathbb{N}|$$



0 1 -1 2 -2 ...

Theorem (\mathbb{Q} is Countable. (Cantor 1873-11; Published in 1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$

Theorem (\mathbb{Q} is Countable. (Cantor 1873-11; Published in 1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$

$|\mathbb{Q}| = |\mathbb{N}|$ (UD Problem 23.12)


$$q \in \mathbb{Q}^+ : a/b \ (a, b \in \mathbb{N}^+)$$

Theorem (\mathbb{Q} is Countable. (Cantor 1873-11; Published in 1874))

$$|\mathbb{Q}| = |\mathbb{N}|$$


$|\mathbb{Q}| = |\mathbb{N}|$ (UD Problem 23.12)

$$q \in \mathbb{Q}^+ : a/b \ (a, b \in \mathbb{N}^+)$$




Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)

$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$



Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)


$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$



$$\pi : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$$

Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)

$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$




$$\pi : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$$

$$\pi(k_1, k_2) = \frac{1}{2}(k_1 + k_2)(k_1 + k_2 + 1) + k_2$$

Theorem ($\mathbb{N} \times \mathbb{N}$ is Countable.)

$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$



$$\pi : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$$

$$\pi(k_1, k_2) = \frac{1}{2}(k_1 + k_2)(k_1 + k_2 + 1) + k_2$$

Cantor Pairing Function

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

*The Cartesian product of **finitely many** countable sets is countable.*

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

*The Cartesian product of **finitely many** countable sets is countable.*

$$\mathbb{N}^n \quad \text{vs.} \quad \mathbb{N}^{\mathbb{N}}$$

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

*The Cartesian product of **finitely many** countable sets is countable.*

$$\mathbb{N}^n \quad \text{vs.} \quad \mathbb{N}^{\mathbb{N}}$$

$$\pi^{(n)} : \mathbb{N}^n \rightarrow \mathbb{N}$$

Theorem (\mathbb{N}^n is Countable.)

$$|\mathbb{N}^n| = |\mathbb{N}|$$

Theorem

*The Cartesian product of **finitely many** countable sets is countable.*

$$\mathbb{N}^n \quad \text{vs.} \quad \mathbb{N}^{\mathbb{N}}$$

$$\pi^{(n)} : \mathbb{N}^n \rightarrow \mathbb{N}$$

$$\pi^{(n)}(k_1, \dots, k_{n-1}, k_n) = \pi(\pi^{(n-1)}(k_1, \dots, k_{n-1}), k_n)$$

Theorem

*Any **finite** union of countable sets is countable.*

Theorem

*Any **finite** union of countable sets is countable.*

$$A = \{a_n \mid n \in \mathbb{N}\} \quad B = \{b_n \mid n \in \mathbb{N}\} \quad C = \{c_n \mid n \in \mathbb{N}\}$$

Theorem

Any *finite* union of countable sets is countable.

$$A = \{a_n \mid n \in \mathbb{N}\} \quad B = \{b_n \mid n \in \mathbb{N}\} \quad C = \{c_n \mid n \in \mathbb{N}\}$$


$$a_0 \quad b_0 \quad c_0 \quad a_1 \quad b_1 \quad c_1 \cdots$$

Theorem

*The union of **countably many** countable sets is countable.*

Theorem


The union of *countably many* countable sets is countable.



Counting by Diagonals.

Theorem

The union of *countably many* countable sets is countable.



Counting by Diagonals.

We need Axiom of (Countable) Choice!

\aleph_0

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

**VERY
IMPORTANT**

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$



Different “Sizes” of Infinity

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$



Different “Sizes” of Infinity

Cantor’s Diagonal Argument (1890)

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} \mathbb{N}$$

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} \mathbb{N}$$

3.14159...
1.41421...
1.73205...
2.23606...
2.71828...
0.14285...



3.43625...



2.32514...

Theorem (\mathbb{R} is Uncountable. (Cantor 1873-12; Published in 1874))

$$|\mathbb{R}| \neq |\mathbb{N}|$$

By Contradiction.

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} \mathbb{N}$$

3.14159...
1.41421...
1.73205...
2.23606...
2.71828...
0.14285...



3.43625...



2.32514...

By Diagonal Argument.

Theorem (Cantor's Theorem (1891))

$$|A| \neq |2^A|$$

Theorem (Cantor's Theorem (1891))

$$|A| \neq |2^A|$$

Theorem (Cantor Theorem (ES Theorem 24.4))

If $f : A \rightarrow 2^A$, then f is not onto.

Theorem (Cantor's Theorem (1891))

$$|A| \neq |2^A|$$

Theorem (Cantor Theorem (ES Theorem 24.4))

If $f : A \rightarrow 2^A$, then f is not onto.

Proof. Let A be a set and let $f : A \rightarrow 2^A$. To show that f is not onto, we must find a $B \in 2^A$ (i.e., $B \subseteq A$) for which there is no $a \in A$ with $f(a) = B$. In other words, B is a set that f “misses.” To this end, let

$$B = \{x \in A : x \notin f(x)\}.$$

We claim there is no $a \in A$ with $f(a) = B$.

Suppose, for the sake of contradiction, there is an $a \in A$ such that $f(a) = B$. We ponder: Is $a \in B$?

- If $a \in B$, then, since $B = f(a)$, we have $a \in f(a)$. So, by definition of B , $a \notin f(a)$; that is, $a \notin B. \Rightarrow \Leftarrow$
- If $a \notin B = f(a)$, then, by definition of B , $a \in B. \Rightarrow \Leftarrow$

Both $a \in B$ and $a \notin B$ lead to contradictions, and hence our supposition [there is an $a \in A$ with $f(a) = B$] is false, and therefore f is not onto. ■

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.




Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

$$2^A = \mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

$$2^A = \mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Onto

$$\forall B \in 2^A : \left(\exists a \in A : f(a) = B \right)$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

$$2^A = \mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Onto

$$\forall B \in 2^A : (\exists a \in A : f(a) = B)$$

Not Onto

$$\exists B \in 2^A : (\forall a \in A : f(a) \neq B)$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

► Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$


Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

► Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.


$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

- ▶ Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$

- ▶ By contradiction (\forall):

$$\exists a \in A : f(a) = B.$$



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$


- ▶ Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$

- ▶ By contradiction (\forall):

$$\exists a \in A : f(a) = B.$$

$$Q : a \in B?$$



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.


$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

- Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$

- By contradiction (\forall):

$$\exists a \in A : f(a) = B.$$



$$Q : a \in B?$$

$$a \in B \iff a \notin B$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Diagonal Argument .

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Diagonal Argument .

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Diagonal Argument .

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Diagonal Argument .

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...

$$B = \{0, 1, 1, 0, 1\}$$



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Diagonal Argument (以下仅适用于可数集合 A).

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...

$$B = \{0, 1, 1, 0, 1\}$$



Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$\begin{array}{l} s_1 = 0000000000\dots \\ s_2 = 1111111111\dots \\ s_3 = 0101010101\dots \\ s_4 = 1010101010\dots \\ s_5 = 1101010101\dots \\ s_6 = 0011010110\dots \\ s_7 = 10001000100\dots \\ s_8 = 0011001001\dots \\ s_9 = 11001100110\dots \\ s_{10} = 11011100101\dots \\ s_{11} = 11010100100\dots \\ \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \end{array}$$

$$s = 10111010011\dots$$

By Diagonal Argument.

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

$$f(x_0x_1\cdots) = \sum_{i=0}^{\infty} x_i 2^i$$

Infinite Sequences of 0's and 1's (UD Problem 23.4)

Is the set of all infinite sequences of 0's and 1's finite, countably infinite, or uncountable?

$$f : \{\{0, 1\}^*\} \rightarrow \mathbb{N}$$

$$f(x_0x_1\cdots) = \sum_{i=0}^{\infty} x_i 2^i$$



Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$f(x) = \tan \frac{(2x - 1)\pi}{2}$$

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$f(x) = \tan \frac{(2x - 1)\pi}{2}$$

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots)$$

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^{n \in \mathbb{N}}|$$

Proof.

$$f(x) = \tan \frac{(2x - 1)\pi}{2}$$

$$|(0, 1)| = |(-\frac{\pi}{2}, \frac{\pi}{2})| = |\mathbb{R}|$$

$$(x = 0.a_1a_2a_3\cdots, y = 0.b_1b_2b_3\cdots) \mapsto 0.a_1b_1a_2b_2a_3b_3\cdots$$



Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^n|$$

“Je le vois, mais je ne le crois pas !”

“I see it, but I don't believe it !”

— Cantor's letter to Dedekind (1877).

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^n|$$

“Je le vois, mais je ne le crois pas !”

“I see it, but I don't believe it !”

— Cantor's letter to Dedekind (1877).

Q : Then, what is “dimension”?

Theorem ($|\mathbb{R}|$ (Cantor 1877))

$$|(0, 1)| = |\mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^n|$$

“Je le vois, mais je ne le crois pas !”


“I see it, but I don't believe it !”

— Cantor's letter to Dedekind (1877).

Q : Then, what is “dimension”?

Theorem (Brouwer (Topological Invariance of Dimension))

*There is no **continuous** bijections between \mathbb{R}^m and \mathbb{R}^n for $m \neq n$.*



Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

bijection $f : A \rightarrow f(A)$ ($\subseteq B$)

Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

bijection $f : A \rightarrow f(A) (\subseteq B)$

Q : What about onto function $f : A \rightarrow B$?

Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

bijection $f : A \rightarrow f(A)$ ($\subseteq B$)

Q : What about onto function $f : A \rightarrow B$?

$|B| \leq |A|$

Definition ($|A| \leq |B|$)

$|A| \leq |B|$ if there exists an *one-to-one* function f from A into B .

bijection $f : A \rightarrow f(A)$ ($\subseteq B$)

Q : What about onto function $f : A \rightarrow B$?

$|B| \leq |A|$ (Axiom of Choice)

Definition ($|A| < |B|$)

$$|A| < |B| \iff |A| \leq |B| \wedge |A| \neq |B|$$

Definition ($|A| < |B|$)

$$|A| < |B| \iff |A| \leq |B| \wedge |A| \neq |B|$$

$$|\mathbb{N}| < |\mathbb{R}|$$

$$|X| < |2^X|$$

Definition (Countable Revisited)

X is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

Definition (Countable Revisited)

X is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable (UD Exercise 22.5))

X is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

Definition (Countable Revisited)

X is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable (UD Exercise 22.5))

X is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

X is countable iff

$$|X| \leq |\mathbb{N}|.$$

Definition (Countable Revisited)

X is countable:

$$\exists n \in \mathbb{N} : |X| = n \vee |X| = |\mathbb{N}|$$

Theorem (Proof for Countable (UD Exercise 22.5))

X is countable iff there exists a *one-to-one* function

$$f : X \rightarrow \mathbb{N}.$$

X is countable iff

$$|X| \leq |\mathbb{N}|.$$

Subsets of Countable Set (UD Problem 22.6; UD Corollary 22.4)

Every subset B of a countable set A is countable.

Set Union (UD Problem 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.

Set Union (UD Problem 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.
- (b) a countably infinite collection of nonempty sets whose union is finite.

Set Union (UD Problem 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.
- (b) a countably infinite collection of nonempty sets whose union is finite.

$$\left(\{A_i : i \in \mathbb{R}\} \mid A_i = \{1\} \right) = \{\{1\}\}$$

Set Union (UD Problem 22.1)

Give an example, if possible, of

- (c) a countably infinite collection of *pairwise disjoint* nonempty sets whose union is finite.
- (b) a countably infinite collection of nonempty sets whose union is finite.

$$\left(\{A_i : i \in \mathbb{R}\} \mid A_i = \{1\} \right) = \{\{1\}\}$$

$$|A| = n \implies |2^A| = 2^n$$

Slope (UD Problem 22.2 (e))

(e) the set of all lines with rational slopes

Slope (UD Problem 22.2 (e))

(e) the set of all lines with rational slopes

$$(\mathbb{Q}, \mathbb{R})$$

Slope (UD Problem 22.2 (e))

(e) the set of all lines with rational slopes

$$(\mathbb{Q}, \mathbb{R})$$

$$|\mathbb{R}| \leq |\mathbb{Q} \times \mathbb{R}| \leq |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}|$$

Q : Is “ \leq ” a partial order?

Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder–Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder–Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$


$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$

Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$




Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$




Q : Is “ \leq ” a partial order?

Theorem (Cantor-Schröder-Bernstein (1887))

$$|X| \leq |Y| \wedge |Y| \leq |X| \implies |X| = |Y|$$

$$\exists \text{ one-to-one } f : A \rightarrow B \wedge g : B \rightarrow A \implies \exists \text{ bijection } h : A \rightarrow B$$



Q : Is “ \leq ” a total order?

Q : Is “ \leq ” a total order?


Theorem (PCC)

Principle of Cardinal Comparability (PCC) \iff Axiom of Choice

Finite Sets



Finite Sets



“关于有穷，我原以为我是懂的”

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n$$

Definition (Finite)

X is finite if

$$\exists n \in \mathbb{N} : |X| = n$$

Theorem (Pigeonhole Principle (UD Theorem 21.2))

$$f : \{1, \dots, m\} \rightarrow \{1, \dots, n\} \quad (m, n \in \mathbb{N}^+, m > n)$$

f is not one-to-one.

$A \setminus \{a\}$ (UD Problem 21.15)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.
Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$A \setminus \{a\}$ (UD Problem 21.15)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.
Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$A \setminus \{a\}$ (UD Problem 21.15)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.
Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$$f|_{A \setminus \{a\}} : A \setminus \{a\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\} \setminus \{f(a)\}$$

$A \setminus \{a\}$ (UD Problem 21.15)

Let A be a nonempty finite set with $|A| = n$ and let $a \in A$.
Prove that $A \setminus \{a\}$ is finite and $|A \setminus \{a\}| = n - 1$.

$$f : A \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\}$$

$$f|_{A \setminus \{a\}} : A \setminus \{a\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n\} \setminus \{f(a)\} \xrightarrow[\text{onto}]{1-1} \{1, \dots, n-1\}$$

$|A| \leq |B|$ (UD Problem 21.17)

A and B are finite sets and $f : A \rightarrow B$ is one-to-one.

Show that $|A| \leq |B|$.

$|A| \leq |B|$ (UD Problem 21.17)

A and B are finite sets and $f : A \rightarrow B$ is one-to-one.
Show that $|A| \leq |B|$.



$|A| \leq |B|$ (UD Problem 21.17)

A and B are finite sets and $f : A \rightarrow B$ is one-to-one.

Show that $|A| \leq |B|$.



By contradiction and the pigeonhole principle.

(UD Problem 21.16)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 20.11). Show that $|B| \leq |A|$.

(UD Problem 21.16)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 20.11). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

(UD Problem 21.16)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 20.11). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

(UD Problem 21.16)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 20.11). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

(UD Problem 21.16)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 20.11). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

- (c) If two finite sets A and B satisfy $B \subseteq A$ and $|A| \leq |B|$, then $A = B$.

(UD Problem 21.16)

- (a) A is a finite set and $B \subseteq A$. We showed that B is finite (Corollary 20.11). Show that $|B| \leq |A|$.

one-to-one $f : B \rightarrow A$

- (b) A is a finite set and $B \subseteq A$. Show that if $B \neq A$, then $|B| < |A|$.

$$\exists a : a \in A \wedge a \notin B \quad f : B \rightarrow A \setminus \{a\} \quad |B| \leq |A \setminus \{a\}|$$

- (c) If two finite sets A and B satisfy $B \subseteq A$ and $|A| \leq |B|$, then $A = B$.

By contradiction and (b).

Cardinality of $|\text{ran}(f)|$ (UD Problem 21.18)

Let A and B be sets with A finite.

$$f : A \rightarrow B$$

Prove that $|\text{ran}(f)| \leq |A|$.

Cardinality of $|\text{ran}(f)|$ (UD Problem 21.18)

Let A and B be sets with A finite.

$$f : A \rightarrow B$$

Prove that $|\text{ran}(f)| \leq |A|$.

one-to-one $g : \text{ran}(f) \rightarrow A$

Cardinality of $|\text{ran}(f)|$ (UD Problem 21.18)

Let A and B be sets with A finite.

$$f : A \rightarrow B$$

Prove that $|\text{ran}(f)| \leq |A|$.

one-to-one $g : \text{ran}(f) \rightarrow A$

(No Axiom of Choice Here)

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

By contradiction.

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\implies

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\Leftarrow

\Rightarrow

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

$$\iff$$

$$\implies$$

$$\forall y \in A \exists x \in A : y = f(x)$$

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\Leftarrow

\Rightarrow

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$$\forall y \in A \exists x \in A : y = f(x)$$

$$\forall y, \text{ choose } x : (g : g(y) = x)$$

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\Leftarrow

\Rightarrow

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$$\forall y \in A \exists x \in A : y = f(x)$$

$$\forall y, \text{ choose } x : (g : g(y) = x)$$

g is bijective.

$f : A \rightarrow A$ (UD Problem 21.19)

Let A be a finite set.

$$f : A \rightarrow A$$

Prove that

$$f \text{ is one-to-one} \iff f \text{ is onto.}$$

\Leftarrow

\Rightarrow

By contradiction.

$$f' : A \rightarrow A \setminus \{a\}$$

$$\forall y \in A \exists x \in A : y = f(x)$$

$$\forall y, \text{ choose } x : (g : g(y) = x)$$

g is bijective.

$$f(g(y)) = f(x) = y \implies f = g^{-1}$$

Dangerous Knowledge (BBC 2007)



Continuum Hypothesis (CH):

$$c = \aleph_1$$

Continuum Hypothesis (CH):

$$\boxed{c = \aleph_1}$$

$$c = 2^{\aleph_0} = \aleph_1$$

Continuum Hypothesis (CH):

$$\boxed{c = \aleph_1}$$

$$c = 2^{\aleph_0} = \aleph_1$$

👉 Dangerous Knowledge (22:20)

Continuum Hypothesis (CH):

$$\boxed{c = \aleph_1}$$

$$c = 2^{\aleph_0} = \aleph_1$$

👉 Dangerous Knowledge (22:20)

Independence from ZFC:

Kurt Gödel (1940) CH cannot be disproved from ZF.

Paul Cohen (1964) CH cannot be proven from the ZFC axioms.

Thank
You!



Office 302

Mailbox: H016

hfwei@nju.edu.cn