


1-10 Set Theory (III): Functions

魏恒峰

hfwei@nju.edu.cn

2019 年 12 月 10 日





Functions

Functions



Functions



PROOF! PROOF! PROOF!

Definition of Functions

$$R \subseteq A \times B$$

is a *relation* from A to B

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

$$f : A \rightarrow B$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

$$f : A \rightarrow B$$

$$\text{dom}(f) = A \quad \text{cod}(f) = B$$

$$\text{ran}(f) = f(A) \subseteq B$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

$$f : A \rightarrow B$$

$$\text{dom}(f) = A \quad \text{cod}(f) = B$$

$$\text{ran}(f) = f(A) \subseteq B$$

$$f : a \mapsto b$$

$$f(a) \triangleq b$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

$$\forall a \in A :$$

$$\forall a \in A : \exists b \in B : (a, b) \in f$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

$$\forall a \in A :$$

$$\forall a \in A : \exists b \in B : (a, b) \in f$$

$$\exists! b \in B :$$

$$\forall b, b' \in B : (a, b) \in f \wedge (a, b') \in f \implies b = b'$$

$$D : \mathbb{R} \rightarrow \mathbb{R}$$

$$D(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$$

Dirichlet Function

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$Y^X = \{f \in \mathcal{P}(X \times Y) \mid f : X \rightarrow Y\}$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$Y^X = \{f \in \mathcal{P}(X \times Y) \mid f : X \rightarrow Y\}$$

X and Y are finite sets with x and y elements, respectively.

$$|X| = x \quad |Y| = y, \quad |Y^X| =$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$Y^X = \{f \in \mathcal{P}(X \times Y) \mid f : X \rightarrow Y\}$$

X and Y are finite sets with x and y elements, respectively.

$$|X| = x \quad |Y| = y, \quad |Y^X| = y^x$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset =$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

$$\emptyset^\emptyset = \{\emptyset\}$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

$$\emptyset^\emptyset = \{\emptyset\}$$

$$\forall X \neq \emptyset : \emptyset^X =$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$\forall Y : Y^\emptyset = \{\emptyset\}$$

$$\emptyset^\emptyset = \{\emptyset\}$$

$$\forall X \neq \emptyset : \emptyset^X = \emptyset$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

$$2^X = \{0, 1\}^X \cong \mathcal{P}(X)$$

$$\mathcal{P}(\{\text{🍏} \text{🍌}\}) = \left\{ \left\{ \begin{array}{l} \text{🍏} \text{🍌} \\ \text{🍏} \\ \text{🍌} \\ \end{array} \right\} \right\} \cong \left\{ \begin{array}{ll} \text{in} & \text{in} \\ \text{in} & \text{out} \\ \text{out} & \text{in} \\ \text{out} & \text{out} \end{array} \right\}$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

Q : Is there a set consisting of all functions?

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

Q : Is there a set consisting of all functions?

Theorem

There is no set consisting of all functions.

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

Q : Is there a set consisting of all functions?

Theorem

There is no set consisting of all functions.

Suppose *by contradiction* that A is the set of all functions.

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

Q : Is there a set consisting of all functions?

Theorem

There is no set consisting of all functions.

Suppose **by contradiction** that A is the set of all functions.

For every set X , there exists a function $I_X : \{X\} \rightarrow \{X\}$.

Definition

The *set* of all functions from X to Y :

$$Y^X = \{f \mid f : X \rightarrow Y\}$$

Q : Is there a set consisting of all functions?

Theorem

There is no set consisting of all functions.

Suppose *by contradiction* that A is the set of all functions.

For every set X , there exists a function $I_X : \{X\} \rightarrow \{X\}$.

$$\bigcup_{I_X \in A} \text{dom}(I_X)$$

Functions as Sets

Axiom (Axiom of Extensionality)

$$\forall A : \forall B : \forall x : (x \in A \iff x \in B) \iff A = B.$$

Axiom (Axiom of Extensionality)

$$\forall A : \forall B : \forall x : (x \in A \iff x \in B) \iff A = B.$$

Theorem (The Principle of Functional Extensionality)

f, g are functions :

$$f = g \iff \text{dom}(f) = \text{dom}(g) \wedge \left(\forall x \in \text{dom}(f) : f(x) = g(x) \right)$$

Axiom (Axiom of Extensionality)

$$\forall A : \forall B : \forall x : (x \in A \iff x \in B) \iff A = B.$$

Theorem (The Principle of Functional Extensionality)

f, g are functions :

$$f = g \iff \text{dom}(f) = \text{dom}(g) \wedge \left(\forall x \in \text{dom}(f) : f(x) = g(x) \right)$$

It may be that $\text{cod}(f) \neq \text{cod}(g)$.

$$f : A \rightarrow B \quad g : C \rightarrow D$$

Q : Is $f \cap g$ a function?

$$f : A \rightarrow B \quad g : C \rightarrow D$$

Q : Is $f \cap g$ a function?

Theorem (Intersection of Functions)

$$f \cap g : (A \cap C) \rightarrow (B \cap D)$$

$$f : A \rightarrow B \quad g : C \rightarrow D$$

Q : Is $f \cup g$ a function?

$$f : A \rightarrow B \quad g : C \rightarrow D$$

Q : Is $f \cup g$ a function?

Theorem (Union of Functions)

$$f \cup g : (A \cup C) \rightarrow (B \cup D) \iff \forall x \in \text{dom}(f) \cap \text{dom}(g) : f(x) = g(x)$$

$$f : A \rightarrow B \quad g : C \rightarrow D$$

Q : Is $f \cup g$ a function?

Theorem (Union of Functions)

$$f \cup g : (A \cup C) \rightarrow (B \cup D) \iff \forall x \in \text{dom}(f) \cap \text{dom}(g) : f(x) = g(x)$$

UD Problem 14.3 (g)

$$f : \mathbb{Q} \rightarrow \mathbb{R}$$

$$f(x) = \begin{cases} x + 1 & \text{if } x \in 2\mathbb{Z} \\ x - 1 & \text{if } x \in 3\mathbb{Z} \\ 2 & \text{otherwise} \end{cases}$$

$$f : A \rightarrow B \quad g : C \rightarrow D$$

Q : Is $f \cup g$ a function?

Theorem (Union of Functions)

$$f \cup g : (A \cup C) \rightarrow (B \cup D) \iff \forall x \in \text{dom}(f) \cap \text{dom}(g) : f(x) = g(x)$$

UD Problem 14.3 (g)

$$f : \mathbb{Q} \rightarrow \mathbb{R}$$

$$f(x) = \begin{cases} x + 1 & \text{if } x \in 2\mathbb{Z} \\ x - 1 & \text{if } x \in 3\mathbb{Z} \\ 2 & \text{otherwise} \end{cases}$$

$$x \in 6\mathbb{Z}$$

UD Problem 14.5

$$f : \mathcal{P}(\mathbb{R}) \rightarrow \mathbb{Z}$$

$$f(A) = \begin{cases} \min(A \cap \mathbb{N}) & \text{if } A \cap \mathbb{N} \neq \emptyset \\ -1 & \text{if } A \cap \mathbb{N} = \emptyset \end{cases}$$

UD Problem 14.5

$$f : \mathcal{P}(\mathbb{R}) \rightarrow \mathbb{Z}$$

$$f(A) = \begin{cases} \min(A \cap \mathbb{N}) & \text{if } A \cap \mathbb{N} \neq \emptyset \\ -1 & \text{if } A \cap \mathbb{N} = \emptyset \end{cases}$$

By the *Well-Ordering Principle* of \mathbb{N}

Special Functions (*-jectivity*)

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B \quad f : A \rightarrowtail B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B \quad f : A \rightarrowtail B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

For Proof:

► To prove that f *is* 1-1:

$$\forall a_1, a_2 \in A : f(a_1) = f(a_2) \implies a_1 = a_2$$

Definition (Injective (one-to-one; 1-1) 单射函数)

$$f : A \rightarrow B \quad f : A \rightarrowtail B$$

$$\forall a_1, a_2 \in A : a_1 \neq a_2 \implies f(a_1) \neq f(a_2)$$

For Proof:

► To prove that f *is* 1-1:

$$\forall a_1, a_2 \in A : f(a_1) = f(a_2) \implies a_1 = a_2$$

► To show that f *is not* 1-1:

$$\exists a_1, a_2 \in A : a_1 \neq a_2 \wedge f(a_1) = f(a_2)$$

Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B$$

$$\text{ran}(f) = B$$

Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B \quad f : A \twoheadrightarrow B$$

$$\text{ran}(f) = B$$

Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B \quad f : A \twoheadrightarrow B$$

$$\text{ran}(f) = B$$

For Proof:

► To prove that f *is* onto:

$$\forall b \in B \left(\exists a \in A : f(a) = b \right)$$

Definition (Surjective (onto) 满射函数)

$$f : A \rightarrow B \quad f : A \twoheadrightarrow B$$

$$\text{ran}(f) = B$$

For Proof:

► To prove that f *is* onto:

$$\forall b \in B \left(\exists a \in A : f(a) = b \right)$$

► To show that f *is not* onto:

$$\exists b \in B \left(\forall a \in A : f(a) \neq b \right)$$

Definition (Bijective (one-to-one correspondence) 一一对应)

$$f : A \rightarrow B$$

1-1 & onto

Definition (Bijective (one-to-one correspondence) 一一对应)

$$f : A \rightarrow B \quad f : A \overset{1-1}{\underset{\text{onto}}{\longleftrightarrow}} B$$

1-1 & onto

Theorem (Cantor Theorem (ES Theorem 24.4))

If $f : A \rightarrow 2^A$, then f is not onto.

Theorem (Cantor Theorem (ES Theorem 24.4))

If $f : A \rightarrow 2^A$, then f is not onto.

Proof. Let A be a set and let $f : A \rightarrow 2^A$. To show that f is not onto, we must find a $B \in 2^A$ (i.e., $B \subseteq A$) for which there is no $a \in A$ with $f(a) = B$. In other words, B is a set that f “misses.” To this end, let

$$B = \{x \in A : x \notin f(x)\}.$$

We claim there is no $a \in A$ with $f(a) = B$.

Suppose, for the sake of contradiction, there is an $a \in A$ such that $f(a) = B$.

We ponder: Is $a \in B$?

- If $a \in B$, then, since $B = f(a)$, we have $a \in f(a)$. So, by definition of B , $a \notin f(a)$; that is, $a \notin B \Rightarrow \Leftarrow$
- If $a \notin B = f(a)$, then, by definition of B , $a \in B \Rightarrow \Leftarrow$

Both $a \in B$ and $a \notin B$ lead to contradictions, and hence our supposition [there is an $a \in A$ with $f(a) = B$] is false, and therefore f is not onto. ■

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.




Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

$$2^A = \mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

$$2^A = \mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Onto

$$\forall B \in 2^A : (\exists a \in A : f(a) = B)$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

Understanding this problem:

$$A = \{1, 2, 3\}$$

$$2^A = \mathcal{P}(A) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$$

Onto

$$\forall B \in 2^A : (\exists a \in A : f(a) = B)$$

Not Onto

$$\exists B \in 2^A : (\forall a \in A : f(a) \neq B)$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

► Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$


Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

► Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.


$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

- ▶ Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$

- ▶ By contradiction (\forall):

$$\exists a \in A : f(a) = B.$$



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.


$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

- ▶ Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$

- ▶ By contradiction (\forall):

$$\exists a \in A : f(a) = B.$$



$$Q : a \in B?$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.


$$\exists B \in 2^A : \left(\forall a \in A : f(a) \neq B \right)$$

- Constructive proof (\exists):

$$B = \{a \in A \mid a \notin f(a)\}$$

- By contradiction (\forall):

$$\exists a \in A : f(a) = B.$$



$$Q : a \in B?$$

$$a \in B \iff a \notin B$$

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

对角线论证 (Cantor's diagonal argument) .

Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

对角线论证 (Cantor's diagonal argument) .

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

对角线论证 (Cantor's diagonal argument) .

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

对角线论证 (Cantor's diagonal argument) .

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...

$$B = \{0, 1, 1, 0, 1\}$$



Theorem (Cantor Theorem)

If $f : A \rightarrow 2^A$, then f is not onto.

对角线论证 (Cantor's diagonal argument) (以下仅适用于可数集合 A).

a	$f(a)$					
	1	2	3	4	5	...
1	1	1	0	0	1	...
2	0	0	0	0	0	...
3	1	0	0	1	0	...
4	1	1	1	1	1	...
5	0	1	0	1	0	...
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	...

$$B = \{0, 1, 1, 0, 1\}$$



Thank
You!