


1-10 Set Theory (III): Functions

魏恒峰

hfwei@nju.edu.cn

2019 年 12 月 10 日





Function

Function



Function



PROOF! PROOF! PROOF!

Definition of Functions

$$R \subseteq A \times B$$

is a *relation* from A to B

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

Notations:

$$f : A \rightarrow B$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

Notations:

$$f : A \rightarrow B$$

$$\text{dom}(f) = A \quad \text{ran}(f) = f(A) \subseteq B \triangleq \text{cod}(f)$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

Notations:

$$f : A \rightarrow B$$

$$\text{dom}(f) = A \quad \text{ran}(f) = f(A) \subseteq B \triangleq \text{cod}(f)$$

$$f : a \mapsto b \triangleq f(a)$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

$$\forall a \in A :$$

$$\forall a \in A : \exists b \in B : (a, b) \in f$$

Definition (Function)

$R \subseteq A \times B$ is a *function* from A to B if

$$\forall a \in A : \exists! b \in B : (a, b) \in f.$$

For Proof:

$$\forall a \in A :$$

$$\forall a \in A : \exists b \in B : (a, b) \in f$$

$$\exists! b \in B :$$

$$\forall b, b' \in B : (a, b) \in f \wedge (a, b') \in f \implies b = b'$$

$$D : \mathbb{R} \rightarrow \mathbb{R}$$

$$D(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$$

Dirichlet Function

UD Problem 14.3 (g)

$$f : \mathbb{Q} \rightarrow \mathbb{R}$$

$$f(x) = \begin{cases} x + 1 & \text{if } x \in 2\mathbb{Z} \\ x - 1 & \text{if } x \in 3\mathbb{Z} \\ 2 & \text{otherwise} \end{cases}$$

UD Problem 14.3 (g)

$$f : \mathbb{Q} \rightarrow \mathbb{R}$$

$$f(x) = \begin{cases} x + 1 & \text{if } x \in 2\mathbb{Z} \\ x - 1 & \text{if } x \in 3\mathbb{Z} \\ 2 & \text{otherwise} \end{cases}$$

$$x = 6$$

UD Problem 14.5

$$f : \mathcal{P}(\mathbb{R}) \rightarrow \mathbb{Z}$$

$$f(A) = \begin{cases} \min(A \cap \mathbb{N}) & \text{if } A \cap \mathbb{N} \neq \emptyset \\ -1 & \text{if } A \cap \mathbb{N} = \emptyset \end{cases}$$

UD Problem 14.5

$$f : \mathcal{P}(\mathbb{R}) \rightarrow \mathbb{Z}$$

$$f(A) = \begin{cases} \min(A \cap \mathbb{N}) & \text{if } A \cap \mathbb{N} \neq \emptyset \\ -1 & \text{if } A \cap \mathbb{N} = \emptyset \end{cases}$$

By the *Well-Ordering Principle* of \mathbb{N}

Axiom (Axiom of Extensionality)

$$\forall A : \forall B : \forall x : (x \in A \iff x \in B) \iff A = B.$$

Axiom (Axiom of Extensionality)

$$\forall A : \forall B : \forall x : (x \in A \iff x \in B) \iff A = B.$$

Theorem (The Principle of Functional Extensionality)

f, g are functions :

$$f = g \iff \text{dom}(f) = \text{dom}(g) \wedge \left(\forall x \in \text{dom}(f) : f(x) = g(x) \right)$$

Axiom (Axiom of Extensionality)

$$\forall A : \forall B : \forall x : (x \in A \iff x \in B) \iff A = B.$$

Theorem (The Principle of Functional Extensionality)

f, g are functions :

$$f = g \iff \text{dom}(f) = \text{dom}(g) \wedge \left(\forall x \in \text{dom}(f) : f(x) = g(x) \right)$$

It may be that $\text{cod}(f) \neq \text{cod}(g)$.

Definition (Intersection, Union)

$$f_1, f_2 : A \rightarrow B$$

- (i) Q : Is $f_1 \cup f_2$ a function from A to B ?
- (ii) Q : Is $f_1 \cap f_2$ a function from A to B ?

Thank
You!