Getting to fit models in ${\bf R}$

Alexandre Courtiol

Leibniz Institute of Zoo and Wildlife Research

June 2018







Leibniz Institute for Zoo and Wildlife Research

IN THE FORSCHUNGSVERBUND BERLIN E.V.

Alexandre Courtiol (IZW) Getting to fit models in R June 2018

- Introduction
- 2 Linear Models
- Generalised Linear Models
- Mixed Models
- 5 General Additive (Mixed) Models

2/27

What is a linear model?

A statistical model represents, often in considerably idealized form, the data-generating process (https://en.wikipedia.org/wiki/Statistical_model).

What is a linear model?

A statistical model represents, often in considerably idealized form, the data-generating process (https://en.wikipedia.org/wiki/Statistical_model).

In a linear model, the data-generating process is assumed to be a linear function: it is constructed from a set of terms by multiplying each term by a constant (a model parameter) and adding the results.

Alexandre Courtiol (IZW) Getting to fit models in R June 2018

What is a linear model?

A statistical model represents, often in considerably idealized form, the data-generating process (https://en.wikipedia.org/wiki/Statistical_model).

In a linear model, the data-generating process is assumed to be a linear function: it is constructed from a set of terms by multiplying each term by a constant (a model parameter) and adding the results.

R allows to fit efficiently and easily all main kinds of linear models:

- classical linear models (t-test, correlation, linear regression, ANOVA, ANCOVA): LM
- generalized linear models (logistic regression, Poisson regression...): GLM
- linear mixed-effects models: LMM
- generalized linear mixed-effects models: GLMM
- general additive models & general additive mixed models: GAM & GAMM

Linear models in R

R is very rich in terms of capabilities to fit linear models due to an increasing number of dedicated packages!

For now, no other software seems to be remotely as good (prognostic: only Julia or Python may change that within a decade but I find it unlikely).

Linear models in R

R is very rich in terms of capabilities to fit linear models due to an increasing number of dedicated packages!

For now, no other software seems to be remotely as good (prognostic: only Julia or Python may change that within a decade but I find it unlikely).

Models	Packages for fitting	Helper packages
LM	none; spaMM	car; visreg
GLM	none; spaMM	car; DHARMa; visreg
LMM	lme4; spaMM; glmmTMB	DHARMa; pbkrtest
GLMM	lme4; spaMM; glmmTMB	DHARMa; pbkrtest
GAM	mgcv	visreg
GAMM	mgcv	

Note: those are my personal favorite ones but they are plenty more out there.

Useful books dealing with linear models in ${\bf R}$



Note: it is also useful to look at books not focussed on R!

Preparing data for (G)LM(M) & GA(M)

To maximize the chances of success prepare your data as follow:

- one row = one observation (if repeated measures, use several rows!)
- qualitative variables of class factor (check the levels, drop unused ones, set the reference properly)
- no NA (models can somewhat deal with them but it is a major source of headackes)

- Introduction
- 2 Linear Models
- Generalised Linear Models
- Mixed Models
- General Additive (Mixed) Models

- Introduction
- Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions

Generalised Linear Models

- introduction & inputs
- outputs
- tests
- predictions
- assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- General Additive (Mixed) Models

June 2018

The linear model

$$y_i = \hat{\beta}_0 + \hat{\beta}_1 \times x_{1,i} + \hat{\beta}_2 \times x_{2,i} + \dots + \hat{\beta}_p \times x_{p,i} + \varepsilon_i$$

- y_i = the observations to explain / response variable / dependent variable
- $x_{i,i}$ = constants derived from the predictors / explanatory variables / independent variables
- $\hat{\beta}_i$ = the (model parameter / regression coefficient) estimates
- ε_i = the residuals (i.e. the estimates for the errors)

outputs

- Introduction
- Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- General Additive (Mixed) Models

- Introduction
- Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- General Additive (Mixed) Models

Getting started with \boldsymbol{R}

- Introduction
- 2 Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

- Introduction
 - 2 Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- **5** General Additive (Mixed) Models

- Introduction
- 2 Linear Models
- Generalised Linear Models
- Mixed Models
- General Additive (Mixed) Models

June 2018

Getting started with ${\bf R}$

- Introduction
- 2 Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

Alexandre Courtiol (IZW) Getting to fit models in R

- Introduction
- 2 Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- **5** General Additive (Mixed) Models

Getting started with \boldsymbol{R}

- Introduction
 - 2 Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

- Introduction
 - Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- **5** General Additive (Mixed) Models

Alexandre Courtiol (IZW) Getting to fit models in R

Getting started with \boldsymbol{R}

- Introduction
 - Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

Alexandre Courtiol (IZW) Getting to fit models in R

- Mixed Models
- General Additive (Mixed) Models

Getting started with \boldsymbol{R}

- Introduction
- 2 Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

- Introduction
 - Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

Getting to fit models in R

- Introduction
 - Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

- Introduction
- Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- General Additive (Mixed) Models

- Introduction
 - Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

- Introduction
 - Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Generalised Linear Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
- Mixed Models
 - introduction & inputs
 - outputs
 - tests
 - predictions
 - assumptions
 - advanced MM
- 5 General Additive (Mixed) Models

June 2018

- Introduction
- 2 Linear Models
- Generalised Linear Models
- Mixed Models
- 5 General Additive (Mixed) Models