

Blood Pressure Tracker

WEEKLY LOG WITH SITTING & STANDING READINGS

No one should face healthcare alone.
The Cancer Lady, RN • thecancerladyrn.com

Name:

Week of:

Day / Date	Time	Sitting Sys / Dia	Standing Sys / Dia	Pulse	Notes (meds, symptoms)
Mon	AM				
	PM				
Tue	AM				
	PM				
Wed	AM				
	PM				
Thu	AM				
	PM				
Fri	AM				
	PM				
Sat	AM				
	PM				
Sun	AM				
	PM				

TIP: Take the sitting reading after 5 min of rest, then stand and measure again after 1 min. A big drop when standing (20+ systolic) can cause dizziness or falls — note it and tell your provider.

No one should face healthcare alone.

Courtney M. Bell, MS, RN • The Cancer Lady, RN • thecancerladyrn.com

This tool is for tracking only and does not replace medical advice. Share your log with your provider.

Blood Pressure Tracker

HOW TO MEASURE • UNDERSTANDING YOUR NUMBERS

No one should face healthcare alone.
The Cancer Lady, RN • thecancerladyrn.com

How to Measure Correctly

1. Rest quietly for 5 minutes before measuring. No caffeine, exercise, or smoking for 30 minutes prior.
2. Sit with back supported, feet flat on the floor, legs uncrossed. Empty your bladder first.
3. Rest your arm on a table so the cuff is at heart level. Place the cuff on bare skin, not over clothing.
4. Don't talk during the reading. Stay still and relaxed.
5. Take two readings one minute apart, at the same times each day (morning and evening). Record both.
6. For a standing (orthostatic) check: measure sitting, then stand and measure again after 1 minute.

Understanding Your Numbers

Category	Systolic (top)		Diastolic (bottom)
Normal	Below 120	and	Below 80
Elevated	120 – 129	and	Below 80
High – Stage 1	130 – 139	or	80 – 89
High – Stage 2	140 or higher	or	90 or higher
Crisis – call now	180 or higher	and/or	120 or higher

Categories based on American Heart Association guidance. Your personal target may differ — especially if you are older or frail, where a slightly higher number can be safer. Always confirm YOUR target with your provider.

When to Call for Help

Call your provider if:

- Your readings are consistently above your target for several days.
- You feel dizzy or faint when standing (a possible orthostatic drop).

Seek emergency care (911) if:

- Your BP is 180/120 or higher AND you have chest pain, trouble breathing, weakness, trouble speaking, or vision changes. These can be signs of a heart attack or stroke.

References

American Heart Association/American College of Cardiology. (2025). 2025 guideline for the prevention, detection, evaluation, and management of high blood pressure in adults. Hypertension. <https://professional.heart.org/en/science-news/2025-high-blood-pressure-guideline>
Mol, A., Maier, A. B., van Wezel, R. J. A., & Meskers, C. G. M. (2024). Hypertension and orthostatic hypotension in the elderly: A challenging balance. The Lancet Regional Health – Europe, 48, 101154. <https://doi.org/10.1016/j.lanepe.2024.101154>

No one should face healthcare alone.

Courtney M. Bell, MS, RN • The Cancer Lady, RN • thecancerladyrn.com

This tool is for tracking only and does not replace medical advice. Share your log with your provider.