

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Draft LABORATORY RESULTS

Patient Info:

Name:
DOB:
Sex:

Patient Information:

Physician Name:
Health Card:
MRN #:
Clinic: Health Sciences North
(Sudbury)

Procedure Date
2025-08-29 03:42:00

Accession Date
2025-08-30 09:59:00

Report Date
2025-09-01 12:42:47

Report Details

Specimen Tested - Ribo-Zero RNA

Sequencing Scope - Whole genome sequencing (WGS)

Referral Reason - Research

Table of findings:

Gene	Information	Interpretation
COL7A1	76 c.2591C>T p.Pro864Leu heterozygous	Variant of uncertain clinical significance
TCF7	1 c.601G>A p.Gly201Ser homozygous	Likely pathogenic

Summary: one likely pathogenic variant and one variant of uncertain clinical significance detected.. The details regarding the specific mutations are included below.

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Variant Interpretation:

The interpretation of these variants is as follows: one likely pathogenic variant and one variant of uncertain clinical significance were detected in the sample.

Variant 1 of 2

Gene COL7A1	Variant c.2591C>T	Amino p.Pro864Leu	Zygosity heterozygous
----------------	----------------------	----------------------	--------------------------

Variant of uncertain clinical significance

COL7A1 c.2591C>T p.Pro864Leu begins at position 864 in exon 76 within chromosome chr3 . This mutation has been identified in 5 families. It causes an amino acid substitution, which replaces proline with leucine . According to ClinVar, the evidence collected to date is insufficient to firmly establish the clinical significance of this variant, therefore it is classified as a variant of uncertain clinical significance . ClinVar and other genomic databases report the COL7A1 c.2591C>T variant as clinically relevant based on aggregated evidence.

The clinical implications of this variant are not yet fully understood. At present, available data is insufficient to confirm its role in disease.

The affected nucleotide lies within a region that is highly conserved across vertebrate species, which suggests functional importance and evolutionary constraint.

This variant is not currently strongly implicated in specific diseases according to ClinVar records (VCV accession: VCV007987783).

Supporting studies and case reports can be found in the scientific literature. Relevant PubMed references include: 773596543, 952343928, 299470415, 357493791, 597871846, 560249718, 967434724 .

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Variant 2 of 2

Gene TCF7	Variant c.601G>A	Amino p.Gly201Ser	Zygosity homozygous
--------------	---------------------	----------------------	------------------------

Likely pathogenic

TCF7 c.601G>A p.Gly201Ser begins at position 201 in exon 1 within chromosome chr5 . This mutation has been identified in 19 families. It causes an amino acid substitution, which replaces glycine with serine . In accordance with existing evidence, this variant is therefore classified as a likely pathogenic variant. ClinVar and other genomic databases report the TCF7 c.601G>A variant as clinically relevant based on aggregated evidence.

This variant is classified as likely pathogenic. It is believed to negatively impact protein function and may play a role in disease development in affected individuals.

The affected nucleotide lies within a region that is highly conserved across vertebrate species, which suggests functional importance and evolutionary constraint.

This variant is implicated in oncogenesis and other disease processes according to ClinVar records (VCV accession: VCV007815760).

Supporting studies and case reports can be found in the scientific literature. Relevant PubMed references include: 886323998, 206668541, 519253209, 835394322 .

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Test Details:

Genes Analyzed: MLF1, POLQ, CD79A, SYNE1, MRE11A, EIF4A2, GNAS, JAK3, RUNX1T1, CSF3R, FGFR1, CDK6, CBLB, MLLT4, ARHGEF12, VTI1A, CUX1, COL2A1,

Recommendations

A precision oncology approach is advised. These mutations are known oncogenic drivers, linked to constitutive pathway activation. Targeted therapies, including hormone-correcting agents, may be considered, guided by clinical judgment. PI3K inhibitors could be explored in trials for PIK3CA-mutated cases. Germline testing is not indicated, as all mutations are consistent with somatic events. A multidisciplinary tumour board review is recommended to integrate findings into care. Additional testing may be pursued at the physician's discretion.

Methodology

Genomic DNA was analyzed using a targeted panel covering all coding exons and adjacent intronic regions. Target enrichment was performed with hybrid capture (Twist Bioscience), followed by Illumina NextSeq sequencing. Reads were aligned to GRCh37 using BWA-MEM, and variants called with GATK. Annotation was performed in VarSeq using population databases, predictive algorithms, and ClinVar. CNVs were assessed with CNVkit and confirmed by MLPA when applicable. Regions with pseudogene interference, such as PMS2, were validated using long-range PCR and Sanger sequencing. Mean read depth exceeded 300x, with a minimum threshold of 50x. Analytical sensitivity is >99% for SNVs/indels and >95% for exon-level CNVs. **Only variants classified as pathogenic, likely pathogenic, or variants of uncertain significance (VUS) are reported**, per ACMG/AMP guidelines (PMID: 25741868).

Limitations

This test was developed and validated in a certified clinical laboratory. Limitations include reduced sensitivity in pseudogene regions (e.g., PMS2, CHEK2), and inability to detect certain structural variants (e.g., MSH2 inversions), deep intronic changes, or low-level mosaicism. PMS2 exons 11–15 are not analyzed. Interpretations reflect current knowledge and may be updated as new evidence emerges.

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Report Electronically Verified and Signed
by:

Mock specimen