

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Draft LABORATORY RESULTS

Patient Info:

Name:
DOB:
Sex:

Patient Information:

Physician Name:
Health Card:
MRN #:
Clinic: Children's Hospital of
Eastern Ontario (Ottawa)

Procedure Date
2023-01-22 04:54:00

Accession Date
2023-01-23 00:13:04

Report Date
2023-01-23 19:45:32

Report Details

Specimen Tested - Other RNA fractions

Sequencing Scope - Whole exome sequencing (WES)

Referral Reason - Research

Table of findings:

Gene	Information	Interpretation
CYP21A2	15 c.1408G>C p.Val470Leu homozygous	Likely pathogenic

Summary: One likely pathogenic variant detected.. The details regarding the specific mutations are included below.

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Variant Interpretation:

The interpretation of these variants is as follows: One likely pathogenic variant was detected in the sample.

Variant 1 of 1

Gene	Variant	Amino	Zygoty
CYP21A2	c.1408G>C	p.Val470Leu	homozygous

Likely pathogenic

CYP21A2 c.1408G>C p.Val470Leu begins at position 470 in exon 15 within chromosome chr6 . This mutation has been identified in 14 families. It causes an amino acid substitution, which replaces valine with leucine . In accordance with existing evidence, this variant is therefore classified as a likely pathogenic variant. ClinVar and other genomic databases report the CYP21A2 c.1408G>C variant as clinically relevant based on aggregated evidence.

This variant is classified as likely pathogenic. It is believed to negatively impact protein function and may play a role in disease development in affected individuals.

The affected nucleotide lies within a region that is highly conserved across vertebrate species, which suggests functional importance and evolutionary constraint.

This variant is implicated in oncogenesis and other disease processes according to ClinVar records (VCV accession: VCV004392353).

Supporting studies and case reports can be found in the scientific literature. Relevant PubMed references include: 195265484, 432975403, 600580030, 620254642 .

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Test Details:

Genes Analyzed: SMAD2, EPHB4, MLLT1, PPARG, KDSR, SET, HOXA9, HNRNP, HOXC11, MARK4,

Recommendations

A precision oncology approach is advised. These mutations are known oncogenic drivers, linked to constitutive pathway activation. Targeted therapies, including hormone-correcting agents, may be considered, guided by clinical judgment. PI3K inhibitors could be explored in trials for PIK3CA-mutated cases. Germline testing is not indicated, as all mutations are consistent with somatic events. A multidisciplinary tumour board review is recommended to integrate findings into care. Additional testing may be pursued at the physician's discretion.

Methodology

Genomic DNA was analyzed using a targeted panel covering all coding exons and adjacent intronic regions. Target enrichment was performed with hybrid capture (Twist Bioscience), followed by Illumina NextSeq sequencing. Reads were aligned to GRCh37 using BWA-MEM, and variants called with GATK. Annotation was performed in VarSeq using population databases, predictive algorithms, and ClinVar. CNVs were assessed with CNVkit and confirmed by MLPA when applicable. Regions with pseudogene interference, such as PMS2, were validated using long-range PCR and Sanger sequencing. Mean read depth exceeded 300x, with a minimum threshold of 50x. Analytical sensitivity is >99% for SNVs/indels and >95% for exon-level CNVs. **Only variants classified as pathogenic, likely pathogenic, or variants of uncertain significance (VUS) are reported**, per ACMG/AMP guidelines (PMID: 25741868).

Limitations

This test was developed and validated in a certified clinical laboratory. Limitations include reduced sensitivity in pseudogene regions (e.g., PMS2, CHEK2), and inability to detect certain structural variants (e.g., MSH2 inversions), deep intronic changes, or low-level mosaicism. PMS2 exons 11–15 are not analyzed. Interpretations reflect current knowledge and may be updated as new evidence emerges.

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Report Electronically Verified and Signed
by:

Mock specimen