

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Draft LABORATORY RESULTS

Patient Info:

Name:
DOB:
Sex:

Patient Information:

Physician Name:
Health Card:
MRN #:
Clinic: Women's College Hospi-
tal (Toronto)

Procedure Date
2025-06-29 04:03:00

Accession Date
2025-06-30 03:05:20

Report Date
2025-06-30 22:36:11

Report Details

Specimen Tested - Other RNA fractions

Sequencing Scope - Whole transcriptome sequencing (WTS)

Referral Reason - Clinical

Table of findings:

Gene	Information	Interpretation
TNR	20 c.590A>T p.Lys197Met heterozy- gous	Variant of uncertain clinical significance

Summary: One variant of uncertain clinical significance detected.. The de-
tails regarding the specific mutations are included below.

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Variant Interpretation:

The interpretation of these variants is as follows: One variant of uncertain clinical significance was detected in the sample.

Variant 1 of 1

Gene TNR	Variant c.590A>T	Amino p.Lys197Met	Zygoty heterozygous
-------------	---------------------	----------------------	------------------------

Variant of uncertain clinical significance

TNR c.590A>T p.Lys197Met begins at position 197 in exon 20 within chromosome chr1 . This mutation has been identified in 7 families. It causes an amino acid substitution, which replaces lysine with methionine . According to ClinVar, the evidence collected to date is insufficient to firmly establish the clinical significance of this variant, therefore it is classified as a variant of uncertain clinical significance . ClinVar and other genomic databases report the TNR c.590A>T variant as clinically relevant based on aggregated evidence.

The clinical implications of this variant are not yet fully understood. At present, available data is insufficient to confirm its role in disease.

The affected nucleotide lies within a region that is highly conserved across vertebrate species, which suggests functional importance and evolutionary constraint.

This variant is not currently strongly implicated in specific diseases according to ClinVar records (VCV accession: VCV000441145).

Supporting studies and case reports can be found in the scientific literature. Relevant PubMed references include: 279200251, 869656436, 963386221, 499707129, 491006281 .

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Test Details:

Genes Analyzed: KMT2C, SPOP, ETV5, FNBP1, ELN, ZNF91, WIF1, MLLT4,

Recommendations

A precision oncology approach is advised. These mutations are known oncogenic drivers, linked to constitutive pathway activation. Targeted therapies, including hormone-correcting agents, may be considered, guided by clinical judgment. PI3K inhibitors could be explored in trials for PIK3CA-mutated cases. Germline testing is not indicated, as all mutations are consistent with somatic events. A multidisciplinary tumour board review is recommended to integrate findings into care. Additional testing may be pursued at the physician's discretion.

Methodology

Genomic DNA was analyzed using a targeted panel covering all coding exons and adjacent intronic regions. Target enrichment was performed with hybrid capture (Twist Bioscience), followed by Illumina NextSeq sequencing. Reads were aligned to GRCh37 using BWA-MEM, and variants called with GATK. Annotation was performed in VarSeq using population databases, predictive algorithms, and ClinVar. CNVs were assessed with CNVkit and confirmed by MLPA when applicable. Regions with pseudogene interference, such as PMS2, were validated using long-range PCR and Sanger sequencing. Mean read depth exceeded 300x, with a minimum threshold of 50x. Analytical sensitivity is >99% for SNVs/indels and >95% for exon-level CNVs. **Only variants classified as pathogenic, likely pathogenic, or variants of uncertain significance (VUS) are reported**, per ACMG/AMP guidelines (PMID: 25741868).

Limitations

This test was developed and validated in a certified clinical laboratory. Limitations include reduced sensitivity in pseudogene regions (e.g., PMS2, CHEK2), and inability to detect certain structural variants (e.g., MSH2 inversions), deep intronic changes, or low-level mosaicism. PMS2 exons 11–15 are not analyzed. Interpretations reflect current knowledge and may be updated as new evidence emerges.

Ontario Cancer Hospital

Division of Tumor
Sequencing and
Diagnostics

123 Main St. West,
Toronto, ON, A4B5G9

info@och.on.ca
fax: 416-456-7890
phone: 437-416-6470

Report Electronically Verified and Signed
by:

Mock specimen