

Table A.5g: First-Year Treatment Effects: Experimental versus Observational Estimates

	(1)	(2)	(3)	(4)	(5)	(6)
	Experimental (IV)			Observational (OLS)		
Outcome Variable	No Controls	Strata FEs	Post-Lasso	No Controls	Strata FEs	Post-Lasso
C. Health Status and Behaviors [survey]						
Ever screened [survey]	0.060*** (0.014) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	0.073*** (0.011) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Physically active [survey]	0.023 (0.026) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	0.020 (0.021) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Trying to be active [survey]	0.008 (0.021) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	0.052*** (0.017) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Current smoker (cigarettes) [survey]	-0.035** (0.014) <i>N</i> =3,566	N/A N/A N/A	N/A N/A N/A	-0.033*** (0.010) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Drinker [survey]	-0.018 (0.025) <i>N</i> =3,565	N/A N/A N/A	N/A N/A N/A	0.010 (0.020) <i>N</i> =2,409	N/A N/A N/A	N/A N/A N/A
Heavy drinker [survey]	-0.004 (0.012) <i>N</i> =3,563	N/A N/A N/A	N/A N/A N/A	-0.003 (0.009) <i>N</i> =2,408	N/A N/A N/A	N/A N/A N/A
Chronic condition [survey]	-0.005 (0.024) <i>N</i> =3,565	N/A N/A N/A	N/A N/A N/A	0.033* (0.019) <i>N</i> =2,409	N/A N/A N/A	N/A N/A N/A
Excellent or v. good health [survey]	-0.007 (0.027) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	-0.015 (0.021) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Not poor health [survey]	-0.006 (0.005) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	0.009* (0.005) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Physical problems [survey]	-0.010 (0.027) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	0.025 (0.021) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Lots of energy [survey]	0.060** (0.025) <i>N</i> =3,566	N/A N/A N/A	N/A N/A N/A	-0.030 (0.020) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Bad emotional health [survey]	0.026 (0.025) <i>N</i> =3,566	N/A N/A N/A	N/A N/A N/A	-0.003 (0.020) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Overweight [survey]	0.014 (0.027) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	0.031 (0.021) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
High BP/cholesterol/glucose [survey]	0.008 (0.025) <i>N</i> =3,567	N/A N/A N/A	N/A N/A N/A	0.030 (0.020) <i>N</i> =2,410	N/A N/A N/A	N/A N/A N/A
Sedentary [survey]	0.002 (0.027) <i>N</i> =3,565	N/A N/A N/A	N/A N/A N/A	0.074*** (0.021) <i>N</i> =2,408	N/A N/A N/A	N/A N/A N/A

Notes: This table replicates Appendix Table A.5g from Jones, Molitor, and Reif (QJE 2019) using public-use data. N/A indicates a censored value.