

School of Mathematics and Statistics

MT5836 Galois Theory

Problem Sheet VII: Radical extensions; solution of equations by radicals;
soluble groups

1. Find a normal radical extension of \mathbb{Q} that contains $\mathbb{Q}(\sqrt[3]{2})$.
2. Find three radical extensions of \mathbb{Q} all containing $\mathbb{Q}(\sqrt{2})$ such that the Galois groups are distinct.
3. Let $f(X) = X^3 - 3X + 1$ and let K be the splitting field of $f(X)$ over \mathbb{Q} . Show that $[K : \mathbb{Q}] = 3$ and find a radical extension of \mathbb{Q} containing K .
Show that K is not itself a radical extension of \mathbb{Q} .
4. Show that $X^5 - 6X + 3$ is not soluble by radicals over \mathbb{Q} .
5. Let F be a field of characteristic zero. Show that a polynomial of the form $X^4 + bX^2 + c$ is soluble by radicals over F .
6. Let G be a soluble group with a chain of subgroups

$$G = G_0 \geq G_1 \geq G_2 \geq \dots \geq G_d = \mathbf{1}$$

where, for $i = 1, 2, \dots, d$, G_i is a normal subgroup of G_{i-1} and G_{i-1}/G_i is abelian.

- (a) If H is a subgroup of G , show that $H \cap G_i$ is a normal subgroup of $H \cap G_{i-1}$ and that $(H \cap G_{i-1})/(H \cap G_i)$ is isomorphic to a subgroup of G_{i-1}/G_i for each i . [Hint: Second Isomorphism Theorem.]
Deduce that subgroups of soluble groups are soluble.
 - (b) If A, B and C are subgroups of G with $A \leq B$, show that $A(B \cap C) = AC \cap B$. [This result is known as the *Modular Law*.]
 - (c) If N is a normal subgroup of G , show that $G_i N/N$ is a normal subgroup of $G_{i-1} N/N$ and that $(G_{i-1} N/N)/(G_i N/N)$ is isomorphic to a quotient of G_{i-1}/G_i for each i . [Hint: Use the Second and Third Isomorphism Theorems and the Modular Law.]
Deduce that quotients of soluble groups are soluble.
7. Let G be a group and N be a normal subgroup of G .
 - (a) If G/N is soluble, show that there is a chain of subgroups

$$G = G_0 \geq G_1 \geq \dots \geq G_k = N$$
 such that G_i is a normal subgroup of G_{i-1} and G_{i-1}/G_i is abelian for $i = 1, 2, \dots, k$. [Hint: Correspondence Theorem.]
 - (b) Deduce that if G/N and N are soluble, then G is soluble.