MATH2017 Problem Set 1 Solutions: Limits of functions, differentiability

- 1. (a) $\exists K \in \mathbb{R}, \forall x \in D, x \leq K$.
 - (b) $\exists y \in D, \forall x \in D, x \leq y$.
 - (c) $\forall y \in D, \exists x \in D, x > y.$
- 2. I claim that $a_n \to 3/2$. Proof: Let $\varepsilon > 0$ be given. Then there exists $N \in \mathbb{Z}^+$ such that $N > 5/(2\varepsilon)$. Now, for all $n \geq N$,

$$|a_n - \frac{3}{2}| = \frac{5}{4n - 2} \le \frac{5}{4n - 2n} = \frac{5}{2n} \le \frac{5}{2N} < \varepsilon.$$

3. For all $m > n \ge 2$,

$$|a_{m} - a_{n}| = \frac{1}{(n+1)!} + \frac{1}{(n+2)!} + \dots + \frac{1}{m!}$$

$$= \frac{1}{(n+1)!} \left(1 + \frac{1}{n+2} + \frac{1}{(n+2)(n+3)} + \dots + \frac{1}{(n+2)(n+3) \dots m} \right)$$

$$< \frac{1}{(n+1)!} \left(1 + \frac{1}{2} + \frac{1}{2^{2}} + \dots + \frac{1}{2^{m-1}} \right)$$

$$\leq \frac{2}{(n+1)!} < \frac{2}{n}.$$

So, let $\varepsilon > 0$ be given. Then choose any $N \in \mathbb{Z}^+$ greater than both 1 and $2/\varepsilon$. Then, for all $m > n \ge N$,

$$|a_m - a_n| < \frac{2}{n} \le \frac{2}{N} < \varepsilon.$$

Hence (a_n) is Cauchy.

4. Let $\varepsilon > 0$ be given. Then take $\delta = 4\sqrt{2}\varepsilon > 0$. Then, for all $x \in [0, \infty)$ such that $0 < |x - 2| < \delta$,

$$\left| \frac{\sqrt{x} - \sqrt{2}}{x - 2} - \frac{1}{2\sqrt{2}} \right| = \left| \frac{(\sqrt{x} - \sqrt{2})(\sqrt{x} + \sqrt{2})}{(x - 2)(\sqrt{x} + \sqrt{2})} - \frac{1}{2\sqrt{2}} \right| = \left| \frac{1}{\sqrt{x} + \sqrt{2}} - \frac{1}{2\sqrt{2}} \right|$$

$$= \left| \frac{\sqrt{x} - \sqrt{2}}{2\sqrt{2}(\sqrt{x} + \sqrt{2})} \right| = \frac{1}{2\sqrt{2}} \left| \frac{x - 2}{(\sqrt{x} + \sqrt{2})^2} \right|$$

$$< \frac{1}{4\sqrt{2}} |x - 2| < \frac{1}{4\sqrt{2}} \delta = \varepsilon.$$

This means that the function $f:[0,\infty)\to\mathbb{R}, f(x)=\sqrt{x}$ is differentiable at 2, with derivative $\frac{1}{2\sqrt{2}}$.

5. We must show that

$$\lim_{x \to 1} \frac{f(x) - f(1)}{x - 1}$$

exists. I claim that it does, and equals 4. Proof: Given any $\varepsilon > 0$, let

$$\delta = \min\{1, \varepsilon/11\}.$$

Then for all $x \in \mathbb{R}$ with $0 < |x - 1| < \delta$,

$$\left| \frac{f(x) - f(1)}{x - 1} - 4 \right| = \left| \frac{x^4 - 1}{x - 1} - 4 \right|$$

$$= \left| (x + 1)(x^2 + 1) - 4 \right|$$

$$= \left| x^3 + x^2 + x - 3 \right| = \left| (x - 1)(x^2 + 2x + 3) \right|$$

$$\leq \left| x - 1 \right| (x^2 + 2|x| + 3) \quad \text{by the triangle inequality}$$

$$\leq \left| x - 1 \right| (4 + 4 + 3) \quad \text{since } |x - 1| < 1, \text{ so } |x| < 2$$

$$< 11\delta \leq \varepsilon.$$