

The L^2 geometry of the moduli space of vortices on the two-sphere in the dissolving limit

Martin Speight (Leeds)

Rene García Lara (Universidad Autonoma de Yucatan)

22/8/22

Rene →



Vortices on the sphere

- Hermitian line bundle L over $\Sigma = (S^2, g_\Sigma)$, degree n

$$E(\phi, A) = \frac{1}{2} \|d_A \phi\|_{L^2}^2 + \frac{1}{2} \|F_A\|_{L^2}^2 + \frac{1}{8} \|\tau - |\phi|^2\|_{L^2}^2$$

Vortices on the sphere

- ▶ Hermitian line bundle L over $\Sigma = (S^2, g_\Sigma)$, degree n

$$E(\phi, A) = \frac{1}{2} \|d_A \phi\|_{L^2}^2 + \frac{1}{2} \|F_A\|_{L^2}^2 + \frac{1}{8} \|\tau - |\phi|^2\|_{L^2}^2$$

- ▶ Bogomolny bound: $E \geq \tau\pi n$, equality \Leftrightarrow

$$\bar{\partial}_A \phi = 0 \quad (V1)$$

$$*F_A = \frac{1}{2}(\tau - |\phi|^2) \quad (V2)$$

Vortices on the sphere

- ▶ Hermitian line bundle L over $\Sigma = (S^2, g_\Sigma)$, degree n

$$E(\phi, A) = \frac{1}{2} \|d_A \phi\|_{L^2}^2 + \frac{1}{2} \|F_A\|_{L^2}^2 + \frac{1}{8} \|\tau - |\phi|^2\|_{L^2}^2$$

- ▶ Bogomolny bound: $E \geq \tau\pi n$, equality \Leftrightarrow

$$\bar{\partial}_A \phi = 0 \quad (V1)$$

$$*F_A = \frac{1}{2}(\tau - |\phi|^2) \quad (V2)$$

- ▶ Bradlow bound: $\int_\Sigma (V2)$:

$$2\pi n = \frac{1}{2} \tau |\Sigma| - \frac{1}{2} \|\phi\|_{L^2}^2$$

$$\|\phi\|_{L^2}^2 = \tau |\Sigma| - 4\pi n =: \varepsilon > 0$$

Vortices on the sphere

- ▶ Hermitian line bundle L over $\Sigma = (S^2, g_\Sigma)$, degree n

$$E(\phi, A) = \frac{1}{2} \|d_A \phi\|_{L^2}^2 + \frac{1}{2} \|F_A\|_{L^2}^2 + \frac{1}{8} \|\tau - |\phi|^2\|_{L^2}^2$$

- ▶ Bogomolny bound: $E \geq \tau\pi n$, equality \Leftrightarrow

$$\bar{\partial}_A \phi = 0 \quad (V1)$$

$$*F_A = \frac{1}{2}(\tau - |\phi|^2) \quad (V2)$$

- ▶ Bradlow bound: $\int_\Sigma (V2)$:

$$2\pi n = \frac{1}{2} \tau |\Sigma| - \frac{1}{2} \|\phi\|_{L^2}^2$$

$$\|\phi\|_{L^2}^2 = \tau |\Sigma| - 4\pi n =: \varepsilon > 0$$

The moduli space (Bradlow)

$$\bar{\partial}_A \phi = 0 \quad (V1), \quad *F_A = \frac{1}{2}(\tau - |\phi|^2) \quad (V2)$$

$$M_n = \{\text{solns } (\phi, A) \text{ of (V1), (V2)}\} / \text{gauge group}$$

The moduli space (Bradlow)

$$\bar{\partial}_A \phi = 0 \quad (V1), \quad *F_A = \frac{1}{2}(\tau - |\phi|^2) \quad (V2)$$

$$M_n = \{\text{solns } (\phi, A) \text{ of (V1), (V2)}\} / \text{gauge group}$$

$\varepsilon < 0$	$\varepsilon = 0$	$\varepsilon > 0$
$M_n = \emptyset$	$M_n = \{pt\}$	$M_n \equiv \mathbb{C}P^n$

The moduli space (Bradlow)

$$\bar{\partial}_A \phi = 0 \quad (V1), \quad *F_A = \frac{1}{2}(\tau - |\phi|^2) \quad (V2)$$

$$M_n = \{\text{solns } (\phi, A) \text{ of (V1), (V2)}\} / \text{gauge group}$$

$\varepsilon < 0$	$\varepsilon = 0$	$\varepsilon > 0$
$M_n = \emptyset$	$M_n = \{pt\}$	$M_n \equiv \mathbb{C}P^n$

► $\varepsilon = 0$: $\phi = 0$, $A = \hat{A}$, const curv


The moduli space (Bradlow)

$$\bar{\partial}_A \phi = 0 \quad (V1), \quad *F_A = \frac{1}{2}(\tau - |\phi|^2) \quad (V2)$$

$$M_n = \{\text{solns } (\phi, A) \text{ of } (V1), (V2)\} / \text{gauge group}$$

$\varepsilon < 0$	$\varepsilon = 0$	$\varepsilon > 0$
$M_n = \emptyset$	$M_n = \{pt\}$	$M_n \equiv \mathbb{C}P^n$

- ▶ $\varepsilon = 0$: $\phi = 0$, $A = \hat{A}$, const curv
- ▶ $\varepsilon > 0$: $[(\phi, A)]$ uniquely determined by **divisor** (ϕ)



$$\Leftrightarrow p(z) = a_0 z^n + a_1 z^{n-1} + \cdots + a_n$$

$$\Leftrightarrow [a_0, a_1, \dots, a_n]$$

The L^2 metric on M_n

- ▶ Any curve of solutions $(\phi(t), A(t))$ of (V1), (V2) defines a tangent vector to M_n at $[(\phi(0), A(0))]$, v say

The L^2 metric on M_n

- ▶ Any curve of solutions $(\phi(t), A(t))$ of (V1), (V2) defines a tangent vector to M_n at $[(\phi(0), A(0))]$, v say
- ▶ Length of v ? Project $(\dot{\phi}(0), \dot{A}(0))$ $L^2 \perp$ gauge orbit through $(\phi(0), A(0))$

$$g_{L^2}(v, v) = \|P(\dot{\phi}(0), \dot{A}(0))\|_{L^2}^2$$

The L^2 metric on M_n

- ▶ Any curve of solutions $(\phi(t), A(t))$ of (V1), (V2) defines a tangent vector to M_n at $[(\phi(0), A(0))]$, v say
- ▶ Length of v ? Project $(\dot{\phi}(0), \dot{A}(0))$ $L^2 \perp$ gauge orbit through $(\phi(0), A(0))$

$$g_{L^2}(v, v) = \|P(\dot{\phi}(0), \dot{A}(0))\|_{L^2}^2$$

- ▶ Controls low energy vortex dynamics (Manton, Stuart)

The L^2 metric on M_n

- ▶ Any curve of solutions $(\phi(t), A(t))$ of (V1), (V2) defines a tangent vector to M_n at $[(\phi(0), A(0))]$, v say
- ▶ Length of v ? Project $(\dot{\phi}(0), \dot{A}(0))$ $L^2 \perp$ gauge orbit through $(\phi(0), A(0))$

$$g_{L^2}(v, v) = \|P(\dot{\phi}(0), \dot{A}(0))\|_{L^2}^2$$

- ▶ Controls low energy vortex dynamics (Manton, Stuart)
- ▶ Kähler. $[\omega_{L^2}]$ known explicitly (Baptista)

$$|M_n| = \frac{\pi^n \varepsilon^n}{n!}$$

The L^2 metric on M_n

- ▶ Any curve of solutions $(\phi(t), A(t))$ of (V1), (V2) defines a tangent vector to M_n at $[(\phi(0), A(0))]$, v say
- ▶ Length of v ? Project $(\dot{\phi}(0), \dot{A}(0))$ $L^2 \perp$ gauge orbit through $(\phi(0), A(0))$

$$g_{L^2}(v, v) = \|P(\dot{\phi}(0), \dot{A}(0))\|_{L^2}^2$$

- ▶ Controls low energy vortex dynamics (Manton, Stuart)
- ▶ Kähler. $[\omega_{L^2}]$ known explicitly (Baptista)

$$|M_n| = \frac{\pi^n \varepsilon^n}{n!}$$

- ▶ Rescale: $g_\varepsilon := \varepsilon^{-1} g_{L^2}$

The L^2 metric on M_n

- ▶ Any curve of solutions $(\phi(t), A(t))$ of (V1), (V2) defines a tangent vector to M_n at $[(\phi(0), A(0))]$, v say
- ▶ Length of v? Project $(\dot{\phi}(0), \dot{A}(0))$ $L^2 \perp$ gauge orbit through $(\phi(0), A(0))$

$$g_{L^2}(v, v) = \|P(\dot{\phi}(0), \dot{A}(0))\|_{L^2}^2$$

- ▶ Controls low energy vortex dynamics (Manton, Stuart)
- ▶ Kähler. $[\omega_{L^2}]$ known explicitly (Baptista)

$$|M_n| = \frac{\pi^n \varepsilon^n}{n!}$$

- ▶ Rescale: $g_\varepsilon := \varepsilon^{-1} g_{L^2}$
- ▶ Baptista-Manton conjecture: $\lim_{\varepsilon \rightarrow 0} g_\varepsilon = \text{Fubini-Study metric}$

The conjecture

- Equip L with hol structure $\bar{\partial}_L = \bar{\partial}_{\hat{A}}$

$$H^0(L) = \{\phi \in \Gamma(L) : \bar{\partial}_{\hat{A}}\phi = 0\} \equiv \mathbb{C}^{n+1}$$

The conjecture

- ▶ Equip L with hol structure $\bar{\partial}_L = \bar{\partial}_{\hat{A}}$

$$H^0(L) = \{\phi \in \Gamma(L) : \bar{\partial}_{\hat{A}}\phi = 0\} \equiv \mathbb{C}^{n+1}$$

- ▶ $S = \{\hat{\phi} \in H^0(L) : \|\hat{\phi}\|_{L^2} = 1\}$, unit sphere

$$\pi : S \rightarrow (\mathbb{P}(H^0(L)), g_{FS}), \quad \hat{\phi} \mapsto \{c\hat{\phi}\}$$

g_{FS} **defined** s.t. π a Riem sub

The conjecture

- ▶ Equip L with hol structure $\bar{\partial}_L = \bar{\partial}_{\hat{A}}$

$$H^0(L) = \{\phi \in \Gamma(L) : \bar{\partial}_{\hat{A}}\phi = 0\} \equiv \mathbb{C}^{n+1}$$

- ▶ $S = \{\hat{\phi} \in H^0(L) : \|\hat{\phi}\|_{L^2} = 1\}$, unit sphere

$$\pi : S \rightarrow (\mathbb{P}(H^0(L)), g_{FS}), \quad \hat{\phi} \mapsto \{c\hat{\phi}\}$$

g_{FS} **defined** s.t. π a Riem sub

- ▶ $f : M_n \rightarrow \mathbb{P}(H^0(L)), [(\phi, A)] \mapsto \{\psi \in H^0(L) : (\psi) = (\phi)\}$

The conjecture

- ▶ Equip L with hol structure $\bar{\partial}_L = \bar{\partial}_{\hat{A}}$

$$H^0(L) = \{\phi \in \Gamma(L) : \bar{\partial}_{\hat{A}}\phi = 0\} \equiv \mathbb{C}^{n+1}$$

- ▶ $S = \{\hat{\phi} \in H^0(L) : \|\hat{\phi}\|_{L^2} = 1\}$, unit sphere

$$\pi : S \rightarrow (\mathbb{P}(H^0(L)), g_{FS}), \quad \hat{\phi} \mapsto \{c\hat{\phi}\}$$

g_{FS} **defined** s.t. π a Riem sub

- ▶ $f : M_n \rightarrow \mathbb{P}(H^0(L)), [(\phi, A)] \mapsto \{\psi \in H^0(L) : (\psi) = (\phi)\}$
- ▶ “The” Fubini-Study metric: $g_0 := f^*g_{FS}$

The conjecture

- ▶ Equip L with hol structure $\bar{\partial}_L = \bar{\partial}_{\hat{A}}$

$$H^0(L) = \{\phi \in \Gamma(L) : \bar{\partial}_{\hat{A}}\phi = 0\} \equiv \mathbb{C}^{n+1}$$

- ▶ $S = \{\hat{\phi} \in H^0(L) : \|\hat{\phi}\|_{L^2} = 1\}$, unit sphere

$$\pi : S \rightarrow (\mathbb{P}(H^0(L)), g_{FS}), \quad \hat{\phi} \mapsto \{c\hat{\phi}\}$$

g_{FS} **defined** s.t. π a Riem sub

- ▶ $f : M_n \rightarrow \mathbb{P}(H^0(L)), [(\phi, A)] \mapsto \{\psi \in H^0(L) : (\psi) = (\phi)\}$
- ▶ “The” Fubini-Study metric: $g_0 := f^*g_{FS}$
- ▶ Baptista-Manton conjecture: $\lim_{\varepsilon \rightarrow 0} g_\varepsilon = g_0$
- ▶ Surprising? Massive gain in symmetry

The theorem

Theorem (JMS, RGL 2022) In the limit $\varepsilon \searrow 0$, g_ε converges in C^0 topology to g_0 .

The theorem

Theorem (JMS, RGL 2022) In the limit $\varepsilon \searrow 0$, g_ε converges in C^0 topology to g_0 .

More precisely:

There exists $C > 0$ such that, for all $v \in TM_n$ and all $\varepsilon \in (0, 1)$

$$|g_\varepsilon(v, v) - g_0(v, v)| \leq C\varepsilon g_0(v, v)$$

The proof

- ▶ Given divisor D , exists $\hat{\phi} \in S \subset H^0(L)$ with $(\hat{\phi}) = D$
Unique up to $\hat{\phi} \mapsto e^{ic} \hat{\phi}$

The proof

- ▶ Given divisor D , exists $\widehat{\phi} \in S \subset H^0(L)$ with $(\widehat{\phi}) = D$
Unique up to $\widehat{\phi} \mapsto e^{ic} \widehat{\phi}$
- ▶ Pseudovortex: $(\sqrt{\varepsilon} \widehat{\phi}, \widehat{A})$
Satisfies (V1) vacuously. Satisfies $\int_{\Sigma} (V2)$

The proof

- ▶ Given divisor D , exists $\widehat{\phi} \in S \subset H^0(L)$ with $(\widehat{\phi}) = D$
Unique up to $\widehat{\phi} \mapsto e^{ic} \widehat{\phi}$
- ▶ Pseudovortex: $(\sqrt{\varepsilon} \widehat{\phi}, \widehat{A})$
Satisfies (V1) vacuously. Satisfies $\int_{\Sigma} (V2)$
- ▶ Deform this to obtain true vortex with $(\phi) = D$:

$$(\phi, A) = (\sqrt{\varepsilon} \widehat{\phi} e^{u/2}, \widehat{A} - \frac{1}{2} * du)$$

for some smooth $u : \Sigma \rightarrow \mathbb{R}$.

The proof

- ▶ Given divisor D , exists $\widehat{\phi} \in S \subset H^0(L)$ with $(\widehat{\phi}) = D$
Unique up to $\widehat{\phi} \mapsto e^{ic} \widehat{\phi}$
- ▶ Pseudovortex: $(\sqrt{\varepsilon} \widehat{\phi}, \widehat{A})$
Satisfies (V1) vacuously. Satisfies $\int_{\Sigma} (V2)$
- ▶ Deform this to obtain true vortex with $(\phi) = D$:

$$(\phi, A) = (\sqrt{\varepsilon} \widehat{\phi} e^{u/2}, \widehat{A} - \frac{1}{2} * du)$$

for some smooth $u : \Sigma \rightarrow \mathbb{R}$.

- ▶ Satisfies (V1) automatically
- ▶ Satisfies (V2) iff

$$\Delta u - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\widehat{\phi}|^2 e^u = 0.$$

The proof

- ▶ Given divisor D , exists $\widehat{\phi} \in S \subset H^0(L)$ with $(\widehat{\phi}) = D$
Unique up to $\widehat{\phi} \mapsto e^{ic} \widehat{\phi}$
- ▶ Pseudovortex: $(\sqrt{\varepsilon} \widehat{\phi}, \widehat{A})$
Satisfies (V1) vacuously. Satisfies $\int_{\Sigma} (V2)$
- ▶ Deform this to obtain true vortex with $(\phi) = D$:

$$(\phi, A) = (\sqrt{\varepsilon} \widehat{\phi} e^{u/2}, \widehat{A} - \frac{1}{2} * du)$$

for some smooth $u : \Sigma \rightarrow \mathbb{R}$.

- ▶ Satisfies (V1) automatically
- ▶ Satisfies (V2) iff

$$\Delta u - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\widehat{\phi}|^2 e^u = 0.$$

- ▶ Energy estimate, elliptic estimate, Sobolev \Rightarrow

$$\|u\|_{C^0} \leq C\varepsilon.$$

Vortices are uniformly well approximated by pseudovortices
(for small ε)

Unpacking that a little...

- Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

Unpacking that a little...

- Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\Delta u_0 - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\hat{\phi}|^2 e^{\bar{u}} e^{u_0} = 0.$$

- To bound $\|u\|_{C^0}$ it suffices to bound

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\int_{\Sigma} \left(\Delta u_0 - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\widehat{\phi}|^2 e^{\bar{u}} e^{u_0} \right) = 0.$$

- ▶ To bound $\|u\|_{C^0}$ it suffices to bound

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$0 - \varepsilon + \varepsilon \int_{\Sigma} |\widehat{\phi}|^2 e^{\bar{u}} e^{u_0} = 0.$$

- ▶ To bound $\|u\|_{C^0}$ it suffices to bound

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ To bound $\|u\|_{C^0}$ it suffices to bound

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound $\|u_0\|_{C^0}$

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound $\|u_0\|_{H^2}$

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound $\|\Delta u_0\|_{H^2}$

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound $\|\Delta u\|_{L^2}$

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\hat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\hat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound $\|F_A - F_{\hat{A}}\|_{L^2}$

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound $\|F_A\|_{L^2}$

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$
- ▶ To bound $\|u\|_{C^0}$ it suffices to bound E

Unpacking that a little...

- ▶ Sobolev: $\|u\|_{C^0} \leq C\|u\|_{H^2}$

$$\|u\|_{H^2}^2 = \int_{\Sigma} u^2 + |du|^2 + |\nabla du|^2$$

- ▶ SEE: for all smooth u with $\int_{\Sigma} u = 0$,

$$\|u\|_{H^2} \leq C\|\Delta u\|_{L^2}$$

- ▶ Decompose $u = u_0 + \bar{u}$ where $\bar{u} = |\Sigma|^{-1} \int_{\Sigma} u$

$$\bar{u} = \ln \left(\int_{\Sigma} |\widehat{\phi}|^2 e^{u_0} \right).$$

- ▶ Monotonicity, $\|\widehat{\phi}\|_{L^2} = 1$: $\|\bar{u}\|_{C^0} \leq \|u_0\|_{C^0}$

- ▶ To bound $\|u\|_{C^0}$ it suffices to bound $E = \tau\pi n$

Unpacking that a little...

► So... $\|u\|_{C^0} \leq C$

Unpacking that a little...

- ▶ So... $\|u\|_{C^0} \leq C$
- ▶ But

$$\Delta u_0 - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\hat{\phi}|^2 e^u = 0$$

Unpacking that a little...

- ▶ So... $\|u\|_{C^0} \leq C$
- ▶ But

$$\Delta u_0 - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\hat{\phi}|^2 e^u = 0$$
$$\Rightarrow \|\Delta u_0\|_{L^2} \leq C\varepsilon$$

Unpacking that a little...

- ▶ So... $\|u\|_{C^0} \leq C$
- ▶ But

$$\Delta u_0 - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\hat{\phi}|^2 e^u = 0$$

$$\Rightarrow \|\Delta u_0\|_{L^2} \leq C\varepsilon$$

$$\Rightarrow \|u\|_{C^0} \leq C\varepsilon$$

Unpacking that a little...

- ▶ So... $\|u\|_{C^0} \leq C$
- ▶ But

$$\Delta u_0 - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\hat{\phi}|^2 e^u = 0$$

$$\Rightarrow \|\Delta u_0\|_{L^2} \leq C\varepsilon$$

$$\Rightarrow \|u\|_{C^0} \leq C\varepsilon$$

- ▶ Vortices are uniformly well approximated by pseudovortices (for small ε)

Unpacking that a little...

- ▶ So... $\|u\|_{C^0} \leq C$
- ▶ But

$$\Delta u_0 - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\hat{\phi}|^2 e^u = 0$$

$$\Rightarrow \|\Delta u_0\|_{L^2} \leq C\varepsilon$$

$$\Rightarrow \|u\|_{C^0} \leq C\varepsilon$$

- ▶ Vortices are uniformly well approximated by pseudovortices (for small ε)
- ▶ Now we need to estimate the *metric*

The proof (cont)

- Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\hat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\hat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ WLOG $\langle i\hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\hat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ WLOG $\langle i\hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ Then $g_0(v, v) = \|\dot{\hat{\phi}}(0)\|_{L^2}^2$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\hat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ WLOG $\langle i\hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ Then $g_0(v, v) = \|\dot{\hat{\phi}}(0)\|_{L^2}^2$
- ▶ $(\dot{\phi}, \dot{A})$ controlled by \dot{u}

$$\Delta u - \frac{\varepsilon}{|\Sigma|} + \varepsilon |\hat{\phi}|^2 e^u = 0.$$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\hat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ WLOG $\langle i\hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ Then $g_0(v, v) = \|\dot{\hat{\phi}}(0)\|_{L^2}^2$
- ▶ $(\dot{\phi}, \dot{A})$ controlled by \dot{u}

$$\Delta \dot{u} + \varepsilon |\hat{\phi}|^2 e^u \dot{u} = -2\varepsilon e^u h(\hat{\phi}, \dot{\hat{\phi}})$$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\hat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ WLOG $\langle i\hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ Then $g_0(v, v) = \|\dot{\hat{\phi}}(0)\|_{L^2}^2$
- ▶ $(\dot{\phi}, \dot{A})$ controlled by \dot{u}

$$\Delta \dot{u} + \varepsilon |\hat{\phi}|^2 e^u \dot{u} = -2\varepsilon e^u h(\hat{\phi}, \dot{\hat{\phi}})$$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \hat{\phi}(t) e^{u(t)/2}, \hat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\hat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ WLOG $\langle i\hat{\phi}, \dot{\hat{\phi}} \rangle_{L^2} = 0$
- ▶ Then $g_0(v, v) = \|\dot{\hat{\phi}}(0)\|_{L^2}^2$
- ▶ $(\dot{\phi}, \dot{A})$ controlled by \dot{u}

$$\Delta \dot{u} + \varepsilon |\hat{\phi}|^2 e^u \dot{u} = -2\varepsilon e^u h(\hat{\phi}, \dot{\hat{\phi}})$$

- ▶ “SEE for $\Delta + \varepsilon |\hat{\phi}|^2 e^u$ ” $\Rightarrow \|\dot{u}\|_{H^2} \leq C\varepsilon \|\dot{\hat{\phi}}\|_{L^2}$

The proof (cont)

- ▶ Take a curve of vortex solutions

$$(\phi(t), A(t)) = (\sqrt{\varepsilon} \widehat{\phi}(t) e^{u(t)/2}, \widehat{A} - \frac{1}{2} * du(t))$$

- ▶ $\|\widehat{\phi}(t)\|_{L^2}^2 \equiv 1 \Rightarrow \langle \widehat{\phi}, \dot{\widehat{\phi}} \rangle_{L^2} = 0$
- ▶ WLOG $\langle i\widehat{\phi}, \dot{\widehat{\phi}} \rangle_{L^2} = 0$
- ▶ Then $g_0(v, v) = \|\dot{\widehat{\phi}}(0)\|_{L^2}^2$
- ▶ $(\dot{\widehat{\phi}}, \dot{\widehat{A}})$ controlled by \dot{u}

$$\Delta \dot{u} + \varepsilon |\widehat{\phi}|^2 e^u \dot{u} = -2\varepsilon e^u h(\widehat{\phi}, \dot{\widehat{\phi}})$$

- ▶ “SEE for $\Delta + \varepsilon |\widehat{\phi}|^2 e^u$ ” $\Rightarrow \|\dot{u}\|_{H^2} \leq C\varepsilon \|\dot{\widehat{\phi}}\|_{L^2}$
- ▶ Suffices to get bound on $|g_\varepsilon - g_0|$.

Convergence of spec Δ

$$\Delta = -g^{ij} \left(\frac{\partial^2}{\partial x_i \partial x_j} + \Gamma_{ij}^k \frac{\partial}{\partial x_k} \right)$$

Convergence of spec Δ

$$\Delta = -g^{ij} \left(\frac{\partial^2}{\partial x_i \partial x_j} + \Gamma_{ij}^k \frac{\partial}{\partial x_k} \right)$$

- Spectrum of Δ on (M, g)

$$0 = \lambda_0(g) < \lambda_1(g) \leq \lambda_2(g) \leq \lambda_3(g) \leq \dots$$

Convergence of spec Δ

$$\Delta = -g^{ij} \left(\frac{\partial^2}{\partial x_i \partial x_j} + \Gamma_{ij}^k \frac{\partial}{\partial x_k} \right)$$

- Spectrum of Δ on (M, g)

$$0 = \lambda_0(g) < \lambda_1(g) \leq \lambda_2(g) \leq \lambda_3(g) \leq \dots$$

- **Corollary** (JMS,RGL): There exists $C, \varepsilon_* > 0$ such that, for all $k \in \mathbb{Z}^+$ and all $\varepsilon \in (0, \varepsilon_*)$,

$$\left| \frac{\lambda_k(g_\varepsilon)}{\lambda_k(g_0)} - 1 \right| \leq C\varepsilon$$

- Spectrum of M_n converges uniformly to spectrum of FS!
- Surprising this follows from only C^0 convergence!

Open questions

- Urakawa-Bando (1983): for any finite dimensional subspace $V \subset C^\infty(M)$

$$\Lambda_g(V) := \sup \left\{ \frac{\|df\|_{L^2}^2}{\|f\|_{L^2}^2} : f \in V \setminus \{0\} \right\}.$$

Then

$$\lambda_k(g) = \inf \{ \Lambda_g(V) : \dim V = k + 1 \}$$

Open questions

- ▶ Urakawa-Bando (1983): for any finite dimensional subspace $V \subset C^\infty(M)$

$$\Lambda_g(V) := \sup \left\{ \frac{\|df\|_{L^2}^2}{\|f\|_{L^2}^2} : f \in V \setminus \{0\} \right\}.$$

Then

$$\lambda_k(g) = \inf \{ \Lambda_g(V) : \dim V = k + 1 \}$$

- ▶ Corollary easily follows

Open questions

- ▶ Convergence of geodesics? Need $g_\varepsilon \rightarrow g_0$ in C^1
- ▶ Convergence of curvature? Need $g_\varepsilon \rightarrow g_0$ in C^2
- ▶ n -dependence of the bounds?
- ▶ Leading correction to g_0 ?
- ▶ Higher genus? Much more subtle (Manton, Romao)