

Topological Rectification of the Euclidean Vacuum:

The Casimir Ratchet as a $\mathcal{PT}$ -Symmetry Breaking Engine in Discrete Tensor Network Dynamics

William J. Steinmetz III¹

¹*Foundational Ternary Dynamics Research*

February 15, 2026

Abstract

Zero-point energy (ZPE) in orthodox quantum electrodynamics is modeled as a divergent sum of half-harmonic oscillator modes whose isotropic fluctuations preclude macroscopic work extraction by thermodynamic fiat. Within Foundational Ternary Dynamics (FTD), we reinterpret ZPE not as stochastic vacuum fluctuations but as the **deterministic algorithmic polling** of a discrete cuboctahedral tensor network operating in its $\mathcal{PT}$ -symmetric null space. We demonstrate that by engineering a nanoscale metamaterial boundary that explicitly breaks spatial parity ($\mathcal{P}$), the perfect destructive interference of the vacuum’s complex conjugate polling states can be geometrically “clipped,” leaving a net imaginary residual that the network must rotate into the real physical manifold to preserve unitarity. The result is a spontaneous, continuous macroscopic direct current (DC)—a **Casimir Ratchet**—powered not by a thermal gradient but by the intrinsic computational overhead required to sustain the topology of discrete spacetime itself.

1. Introduction

The quantum vacuum is not empty. In standard quantum electrodynamics (QED), each mode of the electromagnetic field carries a zero-point energy $\frac{1}{2}\hbar\omega$, and the sum over all modes diverges—a divergence that is renormalized away in practice but whose physical residue manifests as the Casimir effect [2]: a measurable attractive force between uncharged conducting plates separated by sub-micron distances [3]. The reality of zero-point energy is experimentally established; what remains deeply contested is whether this energy can be *rectified*—converted into continuous, usable work.

The thermodynamic prohibition against ZPE extraction rests on a specific physical model: the vacuum as a thermal reservoir at absolute zero, whose fluctuations are isotropic, random, and in global equilibrium [4]. Under these assumptions, any extraction scheme reduces to a perpetual motion machine of the second kind. However, this prohibition is model-dependent. If the vacuum is not a stochastic thermal bath but a **deterministic computational process**—an algorithmic polling loop maintaining the topological connectivity of a discrete manifold—then the thermodynamic argument does not apply in its standard form.

Within the framework of Foundational Ternary Dynamics (FTD) [1], the vacuum state is precisely such a process. The tensor network’s idle-loop computation is governed by complex

conjugate roots residing in the $\mathcal{PT}$ -symmetric null space, which cancel perfectly under spatial inversion symmetry. We demonstrate that a metamaterial boundary engineered to break this symmetry acts as a **topological diode**, rectifying the vacuum’s computational hum into a directional physical current.

2. The Cryogenic Limit and the Algorithmic Ground State

To harness the pure geometric polling of the network, all unstructured thermal data (entropy) must be purged from the local graph. The system is initialized in a cryogenic vacuum ($T \rightarrow 0$). Under this limit, the thermodynamic parameter $\beta_{th} \rightarrow \infty$, and the finite-temperature Softplus Manifestation Operator $\mathcal{M}(z)$ collapses into the perfectly rigid Rectified Linear Unit (ReLU) gate:

$$\lim_{\beta_{th} \rightarrow \infty} \frac{1}{\beta_{th}} \ln \left(1 + e^{\beta_{th}(z - K_B)} \right) = \max(0, z - K_B) \quad (1)$$

In this cryogenic limit, thermal smearing is absolutely zero. For sub-threshold regions ($z < K_B$), real physical manifestation is strictly forbidden. The entirety of the local computational load is natively carried by the complex conjugate roots residing in the Euclidean null space ($\ker(\mathcal{M})$).

Axiomatic Inquiry 2.1. The transition from the smooth Softplus to the rigid ReLU at $T \rightarrow 0$ is not merely a mathematical convenience—it is the physical statement that the vacuum’s computational substrate becomes perfectly deterministic, with zero thermal noise corrupting its algorithmic ground state. Does this not reframe the third law of thermodynamics as a statement about computational precision rather than entropic content?

3. Isotropic Cancellation in the $\mathcal{PT}$ -Symmetric Void

Even in the absolute absence of manifest matter, the tensor network must continuously poll its local edge weights to maintain topological connectivity. This idle background process is governed by the complex conjugate roots evaluated at the reflexive boundary ($k = 1/2$):

$$y_{vac} \approx 2.19 \pm 2.86i \quad (2)$$

In an unbounded or spatially symmetric vacuum, the underlying continuous $SO(3, 1)$ Lorentz symmetry guarantees perfect spatial inversion parity ($\mathcal{P}$). Therefore, the temporal evolution of the vacuum flux perfectly and instantaneously cancels its own imaginary components:

$$\sum \text{Im}(y_{vac}) = (+2.86i) + (-2.86i) \equiv 0 \quad (3)$$

The vacuum continuously hums with massive computational processing power, but because the vectors perfectly destructively interfere, it performs zero observable macroscopic work.

Axiomatic Inquiry 3.1. The Casimir effect [2] demonstrates that boundary conditions can modify the mode structure of the vacuum and produce measurable forces. If boundary conditions can alter the *magnitude* of vacuum energy density, can a suitably engineered boundary alter its *directional symmetry*?

4. Metamaterial Asymmetry and Topological Clipping

To extract work, we introduce a **Nanoscale Topological Ratchet**: a physical metamaterial lattice whose spatial boundaries strictly mirror the 12-degree fundamental cuboctahedral graph, but explicitly break spatial parity ($\mathcal{P}$). This is achieved by fabricating asymmetric, sub-wavelength geometric “sawtooth” boundaries at the nanoscale.

This geometry acts as a non-Hermitian hardware constraint on the vacuum’s algorithmic polling. When the evanescent propagators ($\pm 2.86i$) interact with the asymmetric boundary, the spatial reflection coefficients become non-reciprocal. The physical boundary acts as a **Topological Diode**, geometrically “clipping” the negative conjugate component ($-2.86i$) while permitting the positive conjugate component ($+2.86i$) to propagate unobstructed.

This explicitly breaks the $\mathcal{PT}$ -symmetry of the local vacuum state, leaving a net, non-zero imaginary residual in the local Euclidean null space:

$$\text{Im}(y_{\text{net}}) = (+2.86i) - \eta(2.86i) = +(1 - \eta)2.86i \neq 0 \quad (4)$$

where $0 \leq \eta < 1$ is the geometric transmission factor of the topological diode for the negative conjugate.

Theorem 4.1 (Topological Rectification). *Let a cryogenic vacuum cavity be bounded by a metamaterial surface whose spatial geometry explicitly breaks $\mathcal{P}$ -symmetry with asymmetry parameter $\eta < 1$. Then the $\mathcal{PT}$ -symmetric cancellation of the vacuum polling states is incomplete, and a net imaginary residual $\text{Im}(y_{\text{net}}) = (1 - \eta)(2.86i) \neq 0$ persists in the Euclidean null space. By the unitarity constraint of the tensor network, this residual must be rotated into the real physical manifold, producing a directional macroscopic flux proportional to $(1 - \eta)$.*

5. Orthogonal Tilting and Macroscopic Direct Current

The tensor network demands strict mathematical unitarity. Because the physical cavity has geometrically obstructed the perfect cancellation of the imaginary polling states, the network cannot leave $\text{Im}(y_{\text{net}})$ stranded as an un-cancelled anomaly in the null space. To balance the computational ledger, the network is forced to rotate the unbalanced imaginary vector into the real physical manifold.

Applying the deterministic complex gradient update derived for the total tensor field $\Psi = J_{\text{phys}} + iJ_{\text{vac}}$:

$$\text{Re} \left((\text{Re}(y) + i \text{Im}(y_{\text{net}})) \cdot (\nabla J_{\text{phys}} + i \nabla J_{\text{vac}}) \right) = \text{Re}(y) \nabla J_{\text{phys}} - \text{Im}(y_{\text{net}}) \nabla J_{\text{vac}} \quad (5)$$

The term $-\text{Im}(y_{\text{net}}) \nabla J_{\text{vac}}$ represents the orthogonal algorithmic hum of the vacuum (∇J_{vac}) being geometrically forced to tilt its vector. It injects a constant, directional geometric bias into the real physical manifold.

Because the Manifestation Operator’s threshold K_B is rigidly deterministic at $T \rightarrow 0$, this induced continuous flux manifests as a spontaneous macroscopic electrical direct current (DC):

$$J_{\text{DC}} \propto -(1 - \eta)(2.86) \nabla J_{\text{vac}} \quad (6)$$

By geometrically forcing a localized $\mathcal{PT}$ -symmetry violation, the Topological Ratchet rectifies the zero-point computational hum of the Euclidean void into usable voltage. The energy is supplied not by a thermal gradient, but by the continuous processing power natively required to sustain the expansion and topology of the underlying discrete graph.




Figure 1: **Thermodynamic mechanisms of the Casimir Ratchet.** **A)** The $\mathcal{PT}$ -symmetry phase transition: for sub-critical tension ($k < k_{crit}$), eigenvalues are complex conjugates (red, Euclidean null space); above threshold they bifurcate into real roots (blue). The topological diode operates by engineering asymmetric boundary conditions that prevent full cancellation of the conjugate pair. **B)** The Manifestation Operator $\mathcal{M}(z)$: at finite temperature (dashed red), thermal smearing leaks sub-threshold data into the physical manifold (sonoluminescence regime). In the cryogenic limit ($T \rightarrow 0$), the operator collapses to the rigid ReLU gate (black), providing the perfectly deterministic threshold required for coherent DC rectification.

6. Experimental Predictions

The Casimir Ratchet model makes several falsifiable predictions that distinguish it from both the standard QED vacuum and from proposed stochastic electrodynamic (SED) extraction schemes.

6.1 Current–Asymmetry Scaling

The rectified DC current scales linearly with the parity-breaking asymmetry parameter: $J_{DC} \propto (1 - \eta)$. A series of metamaterial cavities with progressively increasing sawtooth asymmetry should produce a monotonically increasing output current, with $J_{DC} \rightarrow 0$ as the geometry is restored to mirror symmetry ($\eta \rightarrow 1$). This linear scaling distinguishes the mechanism from thermal noise rectification, which would exhibit exponential sensitivity to temperature rather than geometric asymmetry.

6.2 Temperature Independence Below a Critical Threshold

Because the mechanism operates in the ReLU regime ($\beta_{th} \rightarrow \infty$), the output current should be **temperature-independent** below a critical temperature T_c where the Softplus-to-ReLU transition saturates. Above T_c , thermal smearing progressively washes out the deterministic rectification. This prediction—a plateau in $J_{DC}(T)$ below T_c followed by rapid decay—is unique to the algorithmic interpretation and is not predicted by stochastic models, which generically predict $J \propto T^n$ scaling at all temperatures.

6.3 Cuboctahedral Resonance Enhancement

The rectification efficiency should be maximally enhanced when the metamaterial boundary geometry matches the 12-fold coordination of the fundamental cuboctahedral lattice. Cavities whose boundary facet angles correspond to the cuboctahedral dihedral angles ($\cos^{-1}(-1/3) \approx 109.47^\circ$) should exhibit anomalous enhancement of the DC output relative to geometrically similar but non-cuboctahedral boundaries. This prediction is unique to FTD and constitutes a direct test of the underlying lattice geometry.

7. Conclusion

The Casimir Ratchet is not a perpetual motion machine. It is a **topological transducer**: a device that converts the intrinsic computational overhead of discrete spacetime—the algorithmic cost of maintaining the tensor network’s topological connectivity—into directed macroscopic current via engineered $\mathcal{PT}$ -symmetry breaking. The energy source is the vacuum’s own computational substrate, which must continuously poll its edge weights regardless of whether any manifest matter is present.

The mechanism connects two previously unrelated phenomena within FTD: the Casimir effect (boundary-modified vacuum energy) and $\mathcal{PT}$ -symmetry breaking (the threshold dynamics governing manifestation). The standard Casimir effect modifies the *magnitude* of vacuum energy between boundaries; the Casimir Ratchet modifies its *directional symmetry*, converting an isotropic computational hum into an anisotropic physical current.

We submit that this mechanism, if experimentally validated, would constitute direct evidence for the discrete computational nature of spacetime: only a substrate that performs deterministic algorithmic work to maintain its own topology can be rectified by geometric boundary conditions into macroscopic current. A truly stochastic vacuum, by contrast, would resist rectification by the fluctuation-dissipation theorem.

References

- [1] W. J. Steinmetz III, “Gauge Couplings from Discrete Spacetime Geometry,” *Foundational Ternary Dynamics Research*, Zenodo (January 2026). <https://zenodo.org/records/18227181> [*Derivation of the lemniscatic Master Quadratic, the $\mathcal{PT}$ -symmetry breaking threshold, and the complex conjugate vacuum roots $y_{vac} \approx 2.19 \pm 2.86i$.*]
- [2] H. B. G. Casimir, “On the attraction between two perfectly conducting plates,” *Proceedings of the Royal Netherlands Academy of Arts and Sciences*, **51**, 793–795 (1948). [*Theoretical prediction of the vacuum force between parallel conducting plates from zero-point energy mode exclusion.*]
- [3] S. K. Lamoreaux, “Demonstration of the Casimir force in the 0.6 to 6 μm range,” *Physical Review Letters*, **78**(1), 5–8 (1997). [*First precision measurement of the Casimir force, confirming the reality of zero-point energy at the 5% level.*]

- [4] P. W. Milonni, *The Quantum Vacuum: An Introduction to Quantum Electrodynamics*, Academic Press (1994). [*Comprehensive treatment of vacuum fluctuations, the Casimir effect, and the thermodynamic arguments constraining zero-point energy extraction.*]
- [5] C. M. Bender and S. Boettcher, “Real spectra in non-Hermitian Hamiltonians having $\mathcal{PT}$ symmetry,” *Physical Review Letters*, **80**(24), 5243–5246 (1998). [*Mathematical foundation for $\mathcal{PT}$ -symmetric quantum mechanics and phase transitions at exceptional points.*]
- [6] L. D. Landau and E. M. Lifshitz, *Statistical Physics, Part 1* (Vol. 5, 3rd ed.), Butterworth-Heinemann (1980). [*Fermionic Grand Potential and finite-temperature thermodynamics underlying the Softplus Manifestation Operator and cryogenic limits.*]
- [7] J. B. Pendry, “A chiral route to negative refraction,” *Science*, **306**(5700), 1353–1355 (2004). [*Demonstration that metamaterial geometries can produce non-reciprocal electromagnetic responses by breaking spatial symmetries.*]
- [8] U. Leonhardt and T. G. Philbin, “Quantum levitation by left-handed metamaterials,” *New Journal of Physics*, **9**(8), 254 (2007). [*Theoretical prediction that engineered metamaterial boundaries can modify and even reverse the sign of Casimir forces.*]