

THE MASTER QUADRATIC P_{G^*} AS A MATH-PHYSICS BRIDGE: χ_{-4} , THE TERNARY SUBSTRATE, AND THE FTD IDENTIFICATION

ABSTRACT. The companion paper [1] establishes that the Kronecker character χ_{-4} of $K = \mathbb{Q}(i)$ generates the entire algebraic-analytic identity ring of the lemniscatic constant $G^* = \Gamma(1/4)/\Gamma(3/4)$ through four functorial projections (lattice, Chowla-Selberg, Hecke, Dirichlet), and that $\chi_{-4}(n) = \text{Im}(i^n)$ with value set $\{-1, 0, +1\}$. This companion note draws the consequences of those results for the math-physics bridge of the Foundational Ternary Dynamics (FTD) framework [2], where the same set $\{-1, 0, +1\}$ appears as the ternary voxel alphabet of a discrete physical substrate. Three structural observations are recorded, all explicitly tagged as synthesis rather than theorem: (i) the FTD voxel alphabet coincides as a set with the value set of χ_{-4} , sourcing a candidate *Gaussian-integer-module ontology*; (ii) the Born rule $\psi \mapsto |\psi|^2$ and the character $\chi_{-4} : i^n \mapsto \text{Im}(i^n)$ are the same arrow — a real-coordinate projection of a complex object — applied at different scopes; (iii) the FTD identification $(x_+, x_-) \approx (\alpha^{-1}, N_c)$ of the roots of the master quadratic $P_{G^*}(x) = x^2 - 16G^{*2}x + 16G^{*3}$ with the fine-structure constant and the number of QCD colours requires exactly one bit of physical input — the exponent-pair selector $(2, 3)$ — which is over-determined by joint root-matching. All mathematical content is taken from [1]; this companion contributes only the bridge interpretation, presented under the project's [SYNTHESIS] tag.

1. SETTING AND SCOPE

We adopt the entire mathematical content of [1] without re-derivation. The constants in use are:

- $G^* = \Gamma(1/4)/\Gamma(3/4) \approx 2.9587$;
- $G_G = 1/\text{AGM}(1, \sqrt{2}) \approx 0.8346$, with the bridge identity $G^* = 2\sqrt{\pi} G_G$;
- $\chi_{-4} : (\mathbb{Z}/4\mathbb{Z})^\times \rightarrow \{\pm 1\}$, the unique non-trivial Dirichlet character mod 4;
- the master quadratic $P_{G^*}(x) = x^2 - 16G^{*2}x + 16G^{*3}$, with integer coefficient $16 = |\text{Aut}_{\mathbb{C}}(E_{\text{lemn}})|^2$ and roots $x_{\pm}$ satisfying Vieta's identities $x_+ + x_- = 16G^{*2}$, $x_+x_- = 16G^{*3}$, $1/x_+ + 1/x_- = 1/G^*$.

The mathematical paper [1] proves that χ_{-4} admits the explicit form $\chi_{-4}(n) = \text{Im}(i^n) = \sin(\pi n/2)$ and that its value set is

$$\{\chi_{-4}(n) : n \in \mathbb{Z}\} = \{i^2, 0, |i^2|\} = \{-1, 0, +1\}.$$

Numerical observation: the roots of P_{G^*} are $x_+ \approx 137.036 \dots$ (matching the inverse fine-structure constant α^{-1} to 1.26 ppm) and $x_- \approx 3.024 \dots$ (matching the number of QCD colours $N_c = 3$ to 0.8%). The Foundational Ternary Dynamics framework [2] posits the identification $(x_+, x_-) = (\alpha^{-1}, N_c)$ as a strongly motivated conjecture.

This note is the math-physics bridge: it draws structural consequences of the χ_{-4} -unification of [1] for the FTD identification, without proving the identification itself. All load-bearing claims are tagged [SYNTHESIS] or [STRONGLY MOTIVATED CONJECTURE] as appropriate; no new mathematical theorem is claimed in this note.

2. THE VALUE-SET COINCIDENCE AND THE GAUSSIAN-INTEGER-MODULE ONTOLOGY

Observation (Value-set coincidence). [**Synthesis.**] Three independent starting points terminate at the same three-element set:

- (1) the FTD physical primitive — each lattice site v takes a state $s_v \in \{-1, 0, +1\}$ encoding *negative state, void, positive state* [2];

- (2) the value set of the unique non-trivial real Dirichlet character of conductor 4 [1, §16];
- (3) the closure of the imaginary-unit orbit $\{i^n : n \in \mathbb{Z}\}$ under the imaginary-part map: $\{\text{Im}(i^0), \text{Im}(i^1), \text{Im}(i^2), \text{Im}(i^3)\} = \{0, 1, 0, -1\}$.

That these three sets coincide is a set-level identity, not yet a *functorial* identification. Whether the FTD voxel alphabet *arises from* the character χ_{-4} , or only happens to share its value set, is a physical-modelling question that lies outside the scope of the mathematical paper.

Remark (Candidate ontology: $\mathbb{Z}[i]$. -substrate realism) [**Synthesis.**] The value-set coincidence of Observation 2 suggests a working ontology in which the empirical universe is the real-coordinate readout of a $\mathbb{Z}[i]$ -module structure. Layering the established FTD framework against this ontology:

- voxels are positions in a $\mathbb{Z}[i]^d$ -style lattice ($d = 3$ in current FTD);
- voxel states $s_v \in \{-1, 0, +1\}$ are the values $\text{Im}(i^\bullet)$ readouts of the unit group $\mathbb{Z}[i]^\times$;
- the flux field $J \in \mathbb{R}^3$ is the continuous real part of an underlying complex object;
- time is the order parameter of the cyclic action $\langle i \rangle$ at long wavelength;
- the mass-length calibration $a_{\text{phys}} \equiv \ell_P$ is the choice of unit on the $\mathbb{Z}[i]$ -module;
- dimensionless physical constants (α , N_c , mass ratios) are intrinsic invariants of the $\mathbb{Z}[i]$ -module structure.

The ontological position is most accurately described as *Eleatic structural realism, specialized to the Gaussian-integer module*: Eleatic because the substrate is a single algebraic object; structural realist in the sense of Ladyman–Ross [4], where the entities are nodes in the structural network rather than independent objects; specialized because the particular structure is the maximal order of $\mathbb{Q}(i)$ rather than an arbitrary mathematical structure. This is a sharper position than Tegmark’s broad mathematical-universe hypothesis [3]: not “all mathematical structures are physical” but “the physical is the $\mathbb{Z}[i]$ -module slice of the mathematical.” Wigner’s “unreasonable effectiveness” [5] becomes, in this picture, eminently reasonable.

Remark (Falsifiability). [**Synthesis.**] The $\mathbb{Z}[i]$ -substrate ontology is empirically refutable: if experiment ever produced an observable that requires a $\mathbb{Z}[\rho]$ -module structure (Eisenstein, sixfold) that cannot be embedded in a $\mathbb{Z}[i]$ ambient, or that requires $\mathbb{Z}[\sqrt{-2}]$ or $\mathbb{Z}[\sqrt{-7}]$ (other class-number-1 quadratic orders), the framework breaks. The position has therefore the falsifiability profile of a substantive empirical hypothesis, not a metaphysical claim.

3. THE BORN RULE AND χ_{-4} AS THE SAME ARROW

The deepest single structural connection that the χ_{-4} -unification of [1] makes available is the identification of two distinct operations as instances of the same arrow.

Observation (Born rule and χ_{-4} as the same arrow). [**Synthesis.**] The Born rule of quantum mechanics and the character χ_{-4} are formally the same projection applied to different complex objects:

- **Born rule:** $\psi \in \mathbb{C} \mapsto |\psi|^2 \in \mathbb{R}_{\geq 0}$, projecting a complex amplitude onto its real modulus.
- **Character** χ_{-4} : $i^n \in \langle i \rangle \subset \mathbb{C} \mapsto \text{Im}(i^n) \in \{-1, 0, +1\}$, projecting a complex unit onto its signed imaginary coordinate.

Both are functions of the form (complex structure) $\xrightarrow{\text{(single real coordinate)}} \mathbb{R}$. The Born rule reads the modulus; χ_{-4} reads the imaginary part. They are dual projections from the same complex structure, both producing ternary or near-ternary outcomes precisely because the imaginary-part projection of the cyclic group $\langle i \rangle$ takes exactly three distinct values.

Remark (Reflexive projection at three scopes). [**Synthesis.**] Under the FTD framework’s identification of the voxel state field with the substrate’s reflexive projection [2], the same arrow applies

at three nested scopes:

$$\underbrace{\text{reflexive projection of substrate}}_{\text{FTD voxel state} = \text{Im}(i^\bullet)} \sim \underbrace{\text{measurement event of amplitude}}_{\text{Born rule outcome} = |\psi|^2} \sim \underbrace{\text{imaginary-part trace of unit orbit}}_{\chi_{-4}(n) = \text{Im}(i^n)}.$$

The first and third are mathematically equivalent (the FTD voxel state field, viewed as a function on lattice sites with values in $\mathbb{Z}[i]^\times \cup \{0\}$ read off through the imaginary-part map, is literally the character χ_{-4} on the angular index). The second is conjectured-structurally equivalent to the first via the FTD reflexivity ontology; this is the load-bearing physical conjecture and is tagged [STRONGLY MOTIVATED CONJECTURE] accordingly.

4. THE MASTER QUADRATIC AS THE BRIDGE

The FTD bridge from χ_{-4} -derived mathematics to physical observables runs through the master quadratic P_{G^*} . We recall (from [1, Theorem 16.x]) that χ_{-4} generates the integer coefficient $16 = |\mathbb{Z}[i]^\times|^2$ (lattice projection) and the constant G^* (Chowla–Selberg projection), but does *not* generate the specific polynomial form $x^2 - 16G^{*2}x + 16G^{*3}$ from classical CM/Weber/Hecke machinery (the three negative tests of [1, §16.6]).

The exponent pair $(2, 3)$ on G^* in the polynomial coefficients is nevertheless *uniquely picked out* by joint root-matching:

Observation (Joint-matching uniqueness, restated from [1, §17.2]). **[Synthesis.]** Among polynomial forms $y^2 - 16R^p y + 16R^q$ at $R = G^*$ with $(p, q) \in \mathbb{Z}^2$, $|p|, |q| \leq 5$, only the pair $(p, q) = (2, 3)$ produces an ordered root pair (y_+, y_-) that simultaneously matches the FTD pair (α^{-1}, N_c) to sub-percent precision. The match precision at $(2, 3)$ is 1.26 ppm for $y_+ = 137.036$ and 0.8% for $y_- = 3.024$. Other exponent pairs match one root but not both, or neither.

Remark (Information-theoretic minimality of the FTD bridge). **[Synthesis.]** The FTD math-physics bridge requires exactly *one bit* of input from the physical side — the exponent-pair selector $(p, q) \in \{(2, 3), (3, 4), (4, 5), \dots, (1, 2), (2, 4), \dots\}$ — and that bit is over-determined by joint matching to two distinct physical constants (α^{-1}, N_c) . Equivalently: the FTD master quadratic is the unique polynomial of the form $y^2 - 16R^p y + 16R^q$ at $R = G^*$ whose ordered root pair (y_+, y_-) jointly matches (α^{-1}, N_c) to sub-percent precision.

This is not a derivation of the bridge — the joint match itself is the input that selects (p, q) — but it is the cleanest available *minimality statement*: the math-physics bridge in the FTD framework requires no parameters beyond the choice of one exponent pair, and the choice is over-determined (hence robust) against the joint match to two distinct physical constants.

Observation (The locus of the math-physics gap). Combining [1, §16.6] (no CM-internal source for $(2, 3)$) with Observation 4 (joint match uniquely fixes $(2, 3)$): the gap between χ_{-4} -derived mathematical structure and the FTD physical identification is localised precisely at the polynomial *assembly* step. The integer coefficient 16 and the constant G^* are both forced by χ_{-4} . The exponent pair $(2, 3)$ is forced only by the joint physical-matching requirement. Closing this last gap — if a clean derivation of $(2, 3)$ from χ_{-4} -internal data exists — would convert the FTD bridge from a two-arrow conjecture into a single-arrow theorem.

5. THE FULL DESCENT CHAIN

The cumulative content of [1] and this companion can be displayed as a single descent chain from physics to ontological bedrock:

$$\begin{array}{c}
 \underbrace{(\alpha^{-1}, N_c)}_{\text{measured physical constants}} \\
 \uparrow \text{ [STRONGLY MOTIVATED CONJECTURE] (FTD bridge)} \\
 \underbrace{P_{G^*}(x) = x^2 - 16G^{*2}x + 16G^{*3}}_{\text{master quadratic}} \\
 \uparrow (2, 3) \text{ exponents: uniquely forced by joint root match [1, §17.2]} \\
 \underbrace{\{16, G^*\}}_{\text{building blocks}} \\
 \uparrow \text{ four functorial projections [1, Thm 16.x]} \\
 \underbrace{\chi_{-4}}_{\text{Kronecker character of } \mathbb{Q}(i)} \\
 \uparrow \text{ imaginary-part trace } \chi_{-4}(n) = \text{Im}(i^n) = \sin(\pi n/2) \\
 \underbrace{\{-1, 0, +1\}}_{\text{value set = FTD voxel alphabet (set-level)}} \\
 \uparrow \text{ closure of } \langle i \rangle\text{-orbit under squaring \& modulus} \\
 \underbrace{i^2 = -1}_{\text{ontological zero-point}}
 \end{array}$$

Every arrow above the FTD bridge is established (theorem, observation, or honest conjecture) in [1]. The FTD bridge itself is the physical conjecture that this companion records under [STRONGLY MOTIVATED CONJECTURE]. Everything below the bridge is pure mathematics; the only place that requires physical input is the single bit of the exponent-pair selector.

6. OUTLOOK

Two productive lines of work are identified by the descent chain.

Closing the bridge. If a derivation of the exponent pair $(2, 3)$ from χ_{-4} -internal data can be made rigorous — the candidate framing in [1, §16.6, Conjecture (Sym²–Sym³)] is the most natural starting point — the FTD bridge would become a single-arrow theorem rather than a two-arrow conjecture. This would not prove the empirical match to (α^{-1}, N_c) (that remains an experimental fact about the universe), but it would localise the conjectural content to a single physical assertion: that the universe’s electroweak and strong sectors are built from a $\mathbb{Z}[i]$ -substrate.

Extending the bridge. The current FTD identifications cover α and N_c . The Standard Model has additional dimensionless constants ($\sin^2 \theta_W$, m_μ/m_e , m_τ/m_e , PMNS angles, etc.) which the FTD framework currently parameterises without first-principles derivation. The χ_{-4} -unification suggests that further dimensionless invariants may be discoverable as roots of higher-rank master polynomials living in $\text{Sym}^\bullet(H^1(E_{\text{lemn}}))$, or as Hecke L-values of related CM elliptic curves. This is the natural program for extending the bridge from (α^{-1}, N_c) to the full SM.

A second-tier program suggested by the polymath synthesis of [1, appendix] is the identification of the χ_{-3} -side (equianharmonic, $|\text{Aut}| = 6$) as the structural home of the QCD gauge structure, with χ_{-4} (lemniscatic) as the home of the electroweak structure. The Moore Layer Theorem of FTD [2] produces $U(1) \times SU(2) \times SU(3)$ from a 27-block polyhedral decomposition; whether this can be reformulated as the joint action of χ_{-4} (giving $U(1)$ and part of $SU(2)$) and χ_{-3} (giving

$SU(3)$ and the rest of $SU(2)$ via the Hurwitz-quaternion intersection $\mathbb{H} \supset \mathbb{Z}[i], \mathbb{Z}[\rho]$) is an open speculative direction worth investigating but not pursued here.

REFERENCES

- [1] C. Pacitanimulli, *The Kronecker character χ_{-4} as the joint source of the lemniscatic identity algebra*, manuscript, 2026.
- [2] C. Pacitanimulli, *Foundational Ternary Dynamics: specification and algebraic spine*, project documentation, 2026. Available at the project repository; companion manuscript in preparation.
- [3] M. Tegmark, *The Mathematical Universe*, Foundations of Physics **38** (2008), 101–150.
- [4] J. Ladyman, D. Ross, D. Spurrett, and J. Collier, *Every Thing Must Go: Metaphysics Naturalized*, Oxford University Press, 2007.
- [5] E. P. Wigner, *The Unreasonable Effectiveness of Mathematics in the Natural Sciences*, Communications on Pure and Applied Mathematics **13** (1960), 1–14.
- [6] S. Chowla and A. Selberg, *On Epstein’s zeta-function*, Journal für die reine und angewandte Mathematik **227** (1967), 86–110.
- [7] G. V. Chudnovsky, *Algebraic independence of constants connected with the exponential and elliptic functions*, Doklady Akademii Nauk SSSR **4** (1976), 698–701.
- [8] J. Coates and A. Wiles, *On the conjecture of Birch and Swinnerton-Dyer*, Inventiones Mathematicae **39** (1977), 223–251.