

A Precision Formula for the Fine-Structure Constant from Discrete Lattice Combinatorics

William J. Steinmetz III¹

¹*Independent Researcher*

February 2026 — Pre-print Draft

Epistemic Status: [CONJECTURE]

Abstract

We present a closed-form analytic formula for the fine-structure constant α that achieves sub-parts-per-trillion agreement with the CODATA 2022 recommended value. The formula extends the Foundational Ternary Dynamics (FTD) master quadratic—whose larger root $x_+ \approx 137.036$ already approximates α^{-1} to 1.26 ppm—by applying a four-term correction series in the small parameter $\varepsilon = e^\pi - \pi - 20 \approx -9.0 \times 10^{-4}$. Each correction coefficient is an exact rational number constructed from the FTD framework integers $\{3, 4, 7, 13\}$ and the constrained vacuum mode count $D = 47$. The resulting formula yields $\alpha^{-1} = 137.035\,999\,177\,000\,04\dots$, matching the CODATA 2022 central value of $137.035\,999\,177(21)$ to a residual of 4.1×10^{-14} , well within experimental uncertainty. We present this as a conjectured numerical identity, discuss the combinatorial origins of the correction coefficients within the FTD framework, and derive a falsifiable prediction for future precision measurements of α .

Keywords: fine-structure constant, discrete spacetime, lemniscatic constant, FTD, precision formula, lattice combinatorics

PACS: 06.20.Jr, 12.20.Ds, 11.15.Ha

Contents

1	Introduction	2
2	The Correction Formula	3
2.1	The Expansion Parameter	3
2.2	The Correction Coefficients	3
2.3	The Precision Formula	4
2.4	Numerical Evaluation	4

3	Combinatorial Structure of the Coefficients	5
3.1	The Constrained Vacuum Denominator ($D = 47$)	5
3.2	Coefficient Patterns	5
3.3	On the Sign Pattern	6
4	On the Number of Correction Terms	6
4.1	Empirical Saturation	6
4.2	Topological Motivation	6
4.3	Practical vs. Formal Truncation	6
5	Falsifiable Prediction	7
6	Discussion	7
6.1	What This Paper Claims	7
6.2	What This Paper Does Not Claim	7
6.3	Degrees of Freedom Analysis	7
6.4	Relationship to QED	8
6.5	Relationship to Lattice Gauge Theory	8
7	Conclusion	9
A	Computational Verification	9
B	Summary of FTD Framework Integers	10
C	Betti Number Calculation	10

1. Introduction

The fine-structure constant $\alpha \approx 1/137.036$ governs the strength of electromagnetic interactions and is one of the most precisely measured quantities in physics. The CODATA 2022 recommended value is [1]:

$$\alpha^{-1} = 137.035\,999\,177(21), \quad (1)$$

with a relative uncertainty of 1.5×10^{-10} . Despite this extraordinary experimental precision, no fundamental theory predicts the *value* of α from first principles. In the Standard Model, α is a free parameter determined by measurement.

Foundational Ternary Dynamics (FTD) is a discrete computational framework that postulates a three-dimensional cubic lattice with ternary states $\{-1, 0, +1\}$ and a continuous flux field $\mathbf{J} \in \mathbb{R}^3$ [2]. Within this framework, a master quadratic equation derived from the lemniscatic

constant

$$G^* = \frac{\sqrt{2} \Gamma(1/4)^2}{2\pi} \approx 2.9587, \quad (2)$$

yields two roots:

$$x^2 - 16(G^*)^2 x + 16(G^*)^3 = 0. \quad (3)$$

The solutions are:

$$x_+ = 137.036\,171\,458\,155\dots, \quad x_- = 3.023\,963\,916\,339\dots \quad (4)$$

The identification $x_+ \leftrightarrow \alpha^{-1}$ achieves 1.26 ppm accuracy. In this paper, we present a four-term correction formula that closes the remaining gap of $\delta = x_+ - \alpha_{\text{CODATA}}^{-1} \approx 1.72 \times 10^{-4}$, achieving sub-ppt precision.

Remark 1 (Epistemic status). This formula is presented as a [CONJECTURE]—a mathematical observation with suggestive combinatorial structure, not a rigorous derivation from quantum field theory. The physical interpretation of the correction terms remains [OPEN].

2. The Correction Formula

2.1 The Expansion Parameter

We define the dimensionless expansion parameter:

$$\boxed{\varepsilon = e^\pi - \pi - 20 \approx -9.000\,208\,105 \times 10^{-4}} \quad (5)$$

Mathematical context. The near-integer relation $e^\pi \approx \pi + 20$ is documented in number theory [6]. Within FTD, the connection to the framework is twofold:

- (i) $e^\pi = 1/q_L$ where $q_L = e^{-\pi}$ is the nome of the lemniscate elliptic curve underlying the master quadratic;
- (ii) the integer 20 decomposes as $b_3 + N_{\text{eff}} = 7 + 13$, the sum of the topological coefficient and effective mode count from the FTD integer set.

Whether this decomposition has deeper significance or is coincidental remains [OPEN].

2.2 The Correction Coefficients

We define four rational correction coefficients constructed from the FTD framework integers $N_c = 3$, $N_{\text{base}} = 4$, $b_3 = 7$, $N_{\text{eff}} = 13$, and the constrained vacuum denominator $D = N_{\text{base}}^2 \times N_c - 1 = 47$:

$$c_1 = \frac{N_c^2}{D} = \frac{9}{47} \approx 0.19149, \quad (6)$$

$$c_2 = \frac{N_{\text{eff}} - 2 N_{\text{base}}}{N_{\text{base}}^3} = \frac{5}{64} = 0.078125, \quad (7)$$

$$c_3 = \frac{N_{\text{base}}}{N_c \cdot D} = \frac{4}{141} \approx 0.02837, \quad (8)$$

$$c_4 = \frac{N_c \cdot D}{b_3 + N_{\text{base}}} = \frac{141}{11} \approx 12.818. \quad (9)$$

Each coefficient is an exact rational fraction whose numerator and denominator are specific combinatorial functions of the integer set $\{3, 4, 7, 13\}$.

2.3 The Precision Formula

The complete formula is:

$$\boxed{\alpha^{-1} = x_+ - c_1|\varepsilon| + c_2|\varepsilon|^2 - c_3|\varepsilon|^3 - c_4|\varepsilon|^4} \quad (10)$$

with sign pattern $(-, +, -, -)$.

2.4 Numerical Evaluation

Using $G^* = \sqrt{2}\Gamma(1/4)^2/(2\pi)$ computed to 60 decimal places via `mpmath` [7], the successive corrections converge rapidly (see Table 1).

Table 1: Successive correction terms and convergence to the CODATA 2022 value. Each term provides approximately three additional digits of precision.

Stage	Correction	Cumulative α^{-1}	Residual from CODATA
Bare x_+	—	137.036 171 458 155...	$+1.72 \times 10^{-4}$
$-c_1 \varepsilon $	-1.723×10^{-4}	137.035 999 114...	-6.33×10^{-8}
$+c_2 \varepsilon ^2$	$+6.328 \times 10^{-8}$	137.035 999 177 029...	$+2.91 \times 10^{-11}$
$-c_3 \varepsilon ^3$	-2.068×10^{-11}	137.035 999 177 008...	$+8.45 \times 10^{-12}$
$-c_4 \varepsilon ^4$	-8.411×10^{-12}	137.035 999 177 000 04...	$+4.14 \times 10^{-14}$

Final result:

$$\alpha^{-1} = 137.035 999 177 000 041 4... \quad (11)$$

The residual of 4.1×10^{-14} is three orders of magnitude below the CODATA uncertainty of 2.1×10^{-8} .

3. Combinatorial Structure of the Coefficients

In this section we describe the algebraic patterns connecting each coefficient to the discrete phase space of the FTD lattice. We emphasize that these are **structural observations within the FTD framework** [CONJECTURE], not derivations from quantum field theory.

3.1 The Constrained Vacuum Denominator ($D = 47$)

The minimal FTD lattice cell ($2 \times 2 \times 2$) contains 8 vertices, each carrying a 3-component flux vector $\mathbf{J} \in \mathbb{R}^3$, yielding $8 \times 3 = 24$ total degrees of freedom. The Gauss constraint $\nabla_L \cdot \mathbf{J} = \rho$ eliminates 7 longitudinal modes (one per interior face plus one global), and charge conservation removes 1 additional global degree of freedom, leaving

$$24 - 7 - 1 = 16 \quad \text{physical degrees of freedom per cell.} \quad (12)$$

Multiplexed across $N_c = 3$ color sectors with one global constraint subtracted:

$$D = 16 \times N_c - 1 = 47. \quad (13)$$

[SELECTION]: The Gauss constraint counting follows from the FTD action [2]. The color multiplexing and global constraint subtraction are argued from the framework structure but are not independently derived.

3.2 Coefficient Patterns

Each coefficient c_n is a ratio of FTD combinatorial factors. We observe:

- $c_1 = N_c^2/D$: ratio of $U(N_c)$ generator count to the constrained vacuum phase space.
- $c_2 = (N_{\text{eff}} - 2N_{\text{base}})/N_{\text{base}}^3$: ratio of internal interaction modes to volumetric phase space.
- $c_3 = N_{\text{base}}/(N_c \cdot D)$: the first coefficient suppressed by one additional color factor.
- $c_4 = (N_c \cdot D)/(b_3 + N_{\text{base}})$: hierarchical inversion of c_3 's component factors.

We note the algebraic relationship:

$$c_3 \times c_4 = \frac{4}{141} \times \frac{141}{11} = \frac{4}{11} = \frac{N_{\text{base}}}{b_3 + N_{\text{base}}}. \quad (14)$$

3.3 On the Sign Pattern

The sign pattern $(-, +, -, -)$ is empirically determined. In standard QED, the one-loop vacuum polarization screens the bare charge (negative correction to α^{-1}), which is consistent with our c_1 sign. Higher-order corrections involve both screening and anti-screening contributions from vertex corrections and light-by-light scattering.

[IMPOSED]: We do not claim that our sign pattern is derived from QED diagram counting.

4. On the Number of Correction Terms

4.1 Empirical Saturation

The four-term formula saturates the available experimental precision. A hypothetical fifth term of the form $c_5 |\varepsilon|^5$ would contribute at most $\sim c_5 \times 6 \times 10^{-16}$ for c_5 of order unity, which is below any foreseeable experimental sensitivity.

4.2 Topological Motivation

The first Betti number of the cube skeleton (the 1-skeleton of the $2 \times 2 \times 2$ lattice cell with $V = 8$ vertices and $E = 12$ edges) is:

$$b_1 = E - V + 1 = 12 - 8 + 1 = 5. \quad (15)$$

This counts the number of independent cycles in the graph. The Gauss constraint removes one longitudinal (non-physical) cycle, suggesting that the number of *physical* independent topological cycles may be $b_1 - 1 = 4$.

[OPEN]: Whether this topological observation provides a rigorous truncation mechanism, or whether the agreement at four terms is coincidental, remains to be determined.

4.3 Practical vs. Formal Truncation

Even if the series does not formally truncate, the geometric suppression factor $|\varepsilon|^n$ ensures rapid convergence. The ratio of successive terms is:

$$\frac{c_{n+1} |\varepsilon|^{n+1}}{c_n |\varepsilon|^n} \sim |\varepsilon| \times \frac{c_{n+1}}{c_n} \sim 10^{-3} \times O(1) \sim 10^{-3}. \quad (16)$$

Each term provides roughly three additional digits of precision. Four terms exhaust the precision of current experimental data ($\sim 10^{-12}$), rendering the question of formal truncation experimentally moot for the foreseeable future.

5. Falsifiable Prediction

Because the formula produces specific digits beyond current experimental precision, it makes a concrete, falsifiable prediction.

Conjecture 1 (Digit prediction). *The true value of α^{-1} lies within 10^{-12} of*

$$\alpha^{-1} = 137.035\,999\,177\,000\,04(1). \quad (17)$$

Specifically, the digits following the currently measured 137.035 999 177 are predicted to be 000 041..., not (for example) 100 or 200.

Falsification criterion: If a future measurement determines α^{-1} with uncertainty below 10^{-12} and the central value deviates from 137.035 999 177 000 by more than 10^{-11} , this formula is strictly falsified.

Current experimental context. The most precise determination of α comes from the electron $g-2$ measurement combined with five-loop QED calculations [3, 4]. The CODATA 2022 uncertainty of ± 21 in the last two digits ($\times 10^{-9}$) leaves substantial room for this prediction to be tested.

6. Discussion

6.1 What This Paper Claims

1. A closed-form formula for α^{-1} matching CODATA to 4×10^{-14} .
2. All coefficients expressible as exact rationals from FTD integers $\{3, 4, 7, 13\}$.
3. A specific, falsifiable prediction for future precision measurements.

6.2 What This Paper Does Not Claim

1. That the coefficients are QED vacuum polarization loop integrals—they are combinatorial ratios within the FTD integer framework.
2. That the perturbation series is proven to truncate—truncation is motivated but unproven.
3. That the asymptotic divergence of QED is resolved—UV regularization and series truncation are distinct issues.
4. That renormalization is eliminated—the lattice provides a natural cutoff, but the relationship to continuum renormalization is not established.

6.3 Degrees of Freedom Analysis

A legitimate concern is that the formula contains sufficient discrete freedom to fit the CODATA value:

- One expansion parameter expression (chosen, not derived from the lattice action);
- Four rational coefficients (constructed from a four-element integer set);
- Four sign choices ($2^4 = 16$ possibilities).

Against this, we note:

1. The coefficients are not arbitrary—they are constrained to ratios of specific FTD combinatorial quantities.
2. The expansion parameter is not a free real number—it is a fixed mathematical constant.
3. The convergence pattern is remarkably clean, with each term providing ~ 3 additional digits.
4. The formula predicts digits beyond current measurement that can be independently tested.

Whether the formula represents a deep structural truth or an elaborate numerical coincidence is a question that future experiment can decisively answer.

6.4 Relationship to QED

The standard QED approach computes measurable quantities (anomalous magnetic moment, Lamb shift, etc.) as perturbative series in α/π , taking α as input. The coefficients of these series are transcendental numbers involving π , $\zeta(3)$, $\zeta(5)$, etc., and require enormous computational effort at high loop order [4].

Our formula operates in a fundamentally different register: it proposes a *value* for α from discrete combinatorics, rather than using α to predict scattering amplitudes. The two approaches are complementary, not competing. If our formula is correct, it would constrain α more tightly than current QED + experiment, but it does not replace the QED calculations that relate α to observables.

6.5 Relationship to Lattice Gauge Theory

Standard lattice gauge theory (Wilson, 1974) provides a non-perturbative formulation of QFT on a discrete lattice [5]. The FTD lattice shares the UV cutoff property (momentum restricted to the Brillouin zone $[-\pi, \pi]^3$) but differs in crucial respects:

- FTD postulates a ternary state ontology ($\{-1, 0, +1\}$) rather than continuous gauge fields.
- The flux field $\mathbf{J}$ replaces the gauge potential A_μ .
- Manifestation dynamics (threshold crossing) replaces path integral quantization.

The correction coefficients (6)–(9) are *not* derived from lattice Feynman diagram computations. They arise from combinatorial counting of lattice degrees of freedom, not from loop momentum integrals. Establishing a formal connection between this combinatorial structure and standard lattice QED remains an important open problem.

7. Conclusion

We have presented a four-term analytic formula for the inverse fine-structure constant that achieves sub-parts-per-trillion agreement with the CODATA 2022 value. The formula combines the FTD master quadratic root x_+ with correction terms whose coefficients are exact rational numbers built from the framework integers $\{3, 4, 7, 13\}$, expanded in the small parameter $\varepsilon = e^\pi - \pi - 20$.

The result is presented as a mathematical conjecture, not as a QED calculation. Its ultimate significance depends on whether future measurements confirm or refute the predicted digits ...0000414 beyond current experimental precision.

Should this prediction survive future experimental tests, it would constitute strong evidence that the fine-structure constant is determined by the discrete combinatorial structure of spacetime—an extraordinary claim requiring extraordinary evidence, which only precision measurement can provide.

A. Computational Verification

All calculations were performed using `mpmath` [7] at 60-digit precision. The complete computation is reproduced below for independent verification:

```
from mpmath import mp, mpf, gamma, sqrt, pi, exp
mp.dps = 60

G_star = sqrt(2) * gamma(mpf('1')/4)**2 / (2*pi)
disc    = (16*G_star**2)**2 - 4*16*G_star**3
x_plus  = (16*G_star**2 + sqrt(disc)) / 2
eps      = abs(exp(pi) - pi - 20)

c = [mpf(9)/47, mpf(5)/64, mpf(4)/141, mpf(141)/11]
alpha_inv = (x_plus
             - c[0]*eps
             + c[1]*eps**2
             - c[2]*eps**3
             - c[3]*eps**4)
# Result: 137.035999177000041405833862...
```

Output: $\alpha^{-1} = 137.035\,999\,177\,000\,041\,405\,833\,862\,67\dots$

Difference from CODATA: $+4.14 \times 10^{-14}$ (within $\pm 2.1 \times 10^{-8}$ uncertainty).

B. Summary of FTD Framework Integers

Table 2: The four FTD framework integers and their physical interpretations within the lattice framework.

Symbol	Value	Interpretation
N_c	3	Color charges / spatial dimensions
N_{base}	4	Base lattice dimension (2^2)
b_3	7	Topological coefficient (Fibonacci $F_5 + F_3$)
N_{eff}	13	Effective modes ($b_3 + 2N_c = 7 + 6$; also Fibonacci F_7)
D	47	Constrained vacuum: $N_{\text{base}}^2 \times N_c - 1$

C. Betti Number Calculation

For a connected graph G with V vertices and E edges, the first Betti number (cycle rank) is:

$$b_1(G) = E - V + 1. \quad (18)$$

For the 1-skeleton of the $2 \times 2 \times 2$ cube ($V = 8$, $E = 12$):

$$b_1 = 12 - 8 + 1 = 5. \quad (19)$$

This counts five independent topological cycles. If the Gauss constraint eliminates one non-physical (longitudinal) cycle, the number of physical independent cycles is 4, suggesting a possible connection to the four correction terms. This remains [OPEN].

References

- [1] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor, “CODATA recommended values of the fundamental physical constants: 2022,” *Rev. Mod. Phys.* **96**, 025010 (2024).
- [2] W. J. Steinmetz III, “Foundational Ternary Dynamics: A Discrete Ontology for Computational Physics,” Pre-print (2026).
- [3] L. Morel, Z. Yao, P. Cladé, and S. Guellati-Khélifa, “Determination of the fine-structure constant with an accuracy of 81 parts per trillion,” *Nature* **588**, 61–65 (2020).
- [4] T. Aoyama, T. Kinoshita, and M. Nio, “Theory of the anomalous magnetic moment of the electron,” *Atoms* **7**(1), 28 (2019).

- [5] K. G. Wilson, “Confinement of quarks,” *Phys. Rev. D* **10**, 2445 (1974).
- [6] E. W. Weisstein, “Almost integer,” *MathWorld*—A Wolfram Web Resource.
- [7] F. Johansson et al., “mpmath: a Python library for arbitrary-precision floating-point arithmetic (version 1.3),” <https://mpmath.org/> (2023).
- [8] F. J. Dyson, “Divergence of perturbation theory in quantum electrodynamics,” *Phys. Rev.* **85**, 631 (1952).