

# Cross-relaxation in dense ensembles of NV centers and application to magnetometry

Clément Pellet-Mary

PhD Defense

# Outline

## Sensing with quantum mechanics

- Quantum sensing and magnetometry

- NV center magnetometry

## Diamond and NV centers in practice

- Diamond properties

- NV center energy levels

- Basic experiment with NV centers

## Low field depolarization magnetometry (LFDM)

- Principle

- Characterization

- Applications

## Depolarization mechanisms in dense NV ensemble

# Outline

## Sensing with quantum mechanics

### Quantum sensing and magnetometry

#### NV center magnetometry

## Diamond and NV centers in practice

### Diamond properties

### NV center energy levels

### Basic experiment with NV centers

## Low field depolarization magnetometry (LFDM)

### Principle

### Characterization

### Applications

## Depolarization mechanisms in dense NV ensemble


# Quantum sensing and metrology

## Quantum metrology:

Using quantum\* properties to create more sensitive measurement protocols.

\* quantum  $\equiv$  discrete energy levels

Time measurement




Atomic clock

Medical imaging



MRI

Magnetometry



SQUIDs

# Magnetometry key parameters


Magnetoencephalography



$$|\vec{B}| \sim 10 \text{ fT}$$


$$l \sim 1 \text{ cm}$$

Earth's magnetic field anomaly




$$|\vec{B}| \sim 10 \text{ nT}$$

$$l \geq 1 \text{ km}$$




Paleomagnetism



$$|\vec{B}| \sim 100 \text{ nT}$$

$$l \sim 10 \mu\text{m}$$

Surface magnetism




$$|\vec{B}| \sim 100 \mu\text{T}$$

$$l \sim 50 \text{ nm}$$

# Sate of the art magnetometers

Mitchell, M. W., & Alvarez, S. P. *Reviews of Modern Physics*, 92(2), 021001.


Higher spatial resolution



# Sate of the art magnetometers

Mitchell, M. W., & Alvarez, S. P. *Reviews of Modern Physics*, 92(2), 021001.

Higher spatial resolution



Optically pumped magnetometers (OPM)


Optically detected spin precession  
of alkali vapor cell




# Sate of the art magnetometers

Mitchell, M. W., & Alvarez, S. P. *Reviews of Modern Physics*, 92(2), 021001.

Higher spatial resolution



Higher sensitivity




Superconducting quantum interference device (SQUID)

Magnetic field screening due to quantized magnetic flux

# Sate of the art magnetometers

Mitchell, M. W., & Alvarez, S. P. *Reviews of Modern Physics*, 92(2), 021001.

Higher spatial resolution




Nitrogen-Vacancy center in diamond (NV center)

Optically detected spin interferometry


# NV center overview

Point-like defect  
in diamond



Electronic spin 1:  
→ 2 spin transitions

Rondin, L. et al. Reports on progress in physics, 77(5), 056503.




Zeeman effect:  
→ linear dependence of the spin  
transition frequency with  $\vec{B}$

- Works at room temperature
  - Can be put arbitrarily close to the magnetic source  
(low toxicity, chemically inert)
- Potential replacement for SQUID and OPM ?

# NV center magnetometry sensitivity


Ideal (DC) sensitivity for N independent NV centers:

$$\eta[\text{T}/\sqrt{\text{Hz}}] \approx \frac{\hbar\sqrt{\Delta\nu}}{g\mu_B C\sqrt{N}}$$




- $\hbar$  : Planck constant
  - $\mu_B$  : Bohr magneton
  - $g$  : NV electron Landé factor
  - $C$  : Spin readout contrast
  - $N$  : Number of NV centers
  - $\Delta\nu = \frac{1}{T_2^*}$  : Spectral linewidth
- Constants

Experimental parameters
- Sample parameters





# NV center magnetometry: the interaction limit



Going beyond the “Interaction limit” ( $[\text{N}_s] > 10 \text{ ppm}$ ):

- Decoupling interaction (Hamiltonian engineering)
- Exploiting interactions



# Outline

Sensing with quantum mechanics

Quantum sensing and magnetometry

NV center magnetometry

**Diamond and NV centers in practice**

Diamond properties

NV center energy levels

Basic experiment with NV centers

Low field depolarization magnetometry (LFDM)

Principle


Characterization

Applications


Depolarization mechanisms in dense NV ensemble

# Colored centers in diamond


Diamond crystal lattice



Point-like defects





Vacancy



Substitution



Colored center fluorescence





Natural diamonds fluorescence under UV light

# Synthetic diamond and NV centers

High Pressure High Temperature (HPHT)





$N_s$  (P1 center)




Adamas 15/150  $\mu\text{m}$   
 $[N_s] \approx 100 \text{ ppm}$   
 $[NV] \approx 3 \text{ ppm}$

Chemical Vapour Deposition (CVD)





IRCP-LSPM  
 $[N_s] \approx 25 \text{ ppm}$   
 $[NV] \approx 4.5 \text{ ppm}$

# The NV center energy levels




# The NV center energy levels




- Population accumulation in the  $|0\rangle$  state
  - ↳ Initialization of the spin state
- $|0\rangle$  state brighter than  $|\pm 1\rangle$  states
  - ↳ Optical readout of the spin state


# Experimental setup






# Optically detected magnetic resonance (ODMR)



# ODMR with NV ensemble



4 different projections of  $\vec{B}$   
over the 4 possible NV axes  
→ 4 classes of resonances




Position of the 8 lines:  
→ 3D reconstruction of  $\vec{B}$

# Transverse magnetic field effect

$$\hat{\mathcal{H}}_s = D S_z^2 + \gamma \mathbf{B} \cdot \hat{\mathbf{S}}$$

$$D = 2.87 \text{ GHz}, \gamma = 2.8 \text{ MHz/G}$$




$$\mathcal{H}_s = \begin{pmatrix} | -1 \rangle & | 0 \rangle & | +1 \rangle \\ D - \gamma B_z & \gamma B_{\perp} & 0 \\ \gamma B_{\perp} & 0 & \gamma B_{\perp} \\ 0 & \gamma B_{\perp} & D + \gamma B_z \end{pmatrix}$$

When  $D \gg \gamma B_{\perp}$  :

- = Zeeman shift
- = State mixing

$$B_z = |\vec{B}| \cos \theta \quad B_{\perp} = |\vec{B}| \sin \theta / \sqrt{2}$$



# Outline

Sensing with quantum mechanics

Quantum sensing and magnetometry

NV center magnetometry

Diamond and NV centers in practice

Diamond properties

NV center energy levels

Basic experiment with NV centers

**Low field depolarization magnetometry (LFDM)**



**Principle**

**Characterization**


**Applications**

Depolarization mechanisms in dense NV ensemble

# Depolarization of dense NV ensemble at low magnetic field



**Low NV density**  
 $[NV] \leq 100 \text{ ppb}$




**High NV density**  
 $[NV] \geq 1 \text{ ppm}$


- Better understand the dipole-dipole interaction in dense NV ensembles
- Exploit the PL feature for magnetometry

# LFDM experimental setup

## Experimental setup





Samples



Experimental parameters


- $f_{\text{mod}} \sim 1$  kHz
- $|B_{\text{mod}}| \sim 10$  G
- $I_{\text{las}} \sim 1$  mW
- $\text{PL} \sim 1$   $\mu$ W

# Sensitivity of LFDM



Low pass filter  $\tau = 3 \text{ ms}$        $\sqrt{\langle \delta B^2 \rangle} \approx 1.2 \mu\text{T}$

$$\rightarrow \text{sensitivity } \eta = \sqrt{2\tau \langle \delta B^2 \rangle} \approx 116 \text{ nT}/\sqrt{\text{Hz}}$$



Shot noise limited  
for  $f_{\text{mod}} > 500 \text{ Hz}$

# Comparison with the state of the art

Sensitivity comparison



	GSLAC [1]	ODMR [2]	LFDM
$\eta$ (nT/ $\sqrt{\text{Hz}}$ )	0.3*	0.015	116
$V$ ( $\mu\text{m}^3$ )	??	$5.2 \cdot 10^6$	$3.3 \cdot 10^3$
$\eta_v$ (nT $\mu\text{m}^{3/2}\text{Hz}^{-1/2}$ )	??	34	6700

[1] Zheng, H.[...] Budker, D. (2020). Physical Review Applied, 13(4), 044023.

[2] Barry, J. F. [...] Walsworth, R. L (2016). PNAS, 113(49), 14133-14138.

	ODMR	GSLAC	LFDM
Microwave free	✗	✓	✓
Low magnetic field (<10 G)	✓	✗	✓
Robust to T° and B-field inhomogeneities	✗	✗	✓
Orientation free (polycrystalline, powder)	✗	✗	✓



# Comparison with CW ODMR



Adding a fixed microwave tone



$$\text{MW slope sensitivity: } \eta \approx 40 \text{ nT}/\sqrt{\text{Hz}}$$
$$B=0 \text{ sensitivity: } \eta \approx 120 \text{ nT}/\sqrt{\text{Hz}}$$

## Temporal stability



# Application: wide-field magnetometry on irregular surfaces

(Commercially available  $1 \mu\text{m}$  diamonds)



Glenn, D. R. [...] Walsworth, R. L. (2017)  
Geochemistry, Geophysics, Geosystems, 18(8), 3254-3267.

Area normalized sensitivity:  
 $\eta_S \approx 6 \mu\text{T} \cdot \mu\text{m}/\sqrt{\text{Hz}}$

Area normalized sensitivity:  
 $\eta_S \approx 20 \mu\text{T} \cdot \mu\text{m}/\sqrt{\text{Hz}}$

# Outline

Sensing with quantum mechanics

Quantum sensing and magnetometry

NV center magnetometry

Diamond and NV centers in practice

Diamond properties

NV center energy levels

Basic experiment with NV centers

Low field depolarization magnetometry (LFDM)



Principle

Characterization



Applications

Depolarization mechanisms in dense NV ensemble


## Principle of cross-relaxation with NV centers





# Example: Cross-relaxation between NV centers and VH<sup>-</sup>



NV center




VH center






Optical detection of paramagnetic defects in diamond grown by chemical vapor deposition

C. Pellet-Mary, P. Huillery, M. Perdrat, A. Tallaire, and G. Hétet  
Phys. Rev. B **103**, L100411 – Published 24 March 2021


# Cross-relaxation between NV centers and NV centers



samples : Adamas 15/150  $\mu\text{m}$   
 fluorescent microdiamond  
 $[\text{N}] = 100\sim200 \text{ PPM}$   
 $[\text{NV}] \sim 3 \text{ PPM}$



# Presentation of the fluctuator model



Fluctuators are NV centers with a fast intrinsic depolarization mechanism



Localized noise sources with the spectral response of an NV center

Precedents in:

- P-doped Si
- solid-state NMR
- FRET



Possible microscopic explanation:

- charge tunneling
- modulation of J-coupling

Up to 1/3 of all NV centers could be fluctuators



Choi, Joonhee, et al. Physical review letters 118.9 (2017): 093601.

# Stretched exponential decay profile



Low NV density

- Exponential profile
- $T_1 \sim 5$  ms




High NV density

- Stretched exp. profile
- $T_1 \sim 0.5$  ms

## Zero field depolarization sources (theory)

$$\Gamma_1 = \Gamma_0^{\text{th}}$$




$$\Gamma_1 = 102 \Gamma_0^{\text{th}}$$

- 4-classes degeneracy  
 $\Gamma_1 = \Gamma_0^{\text{th}} \rightarrow 43\Gamma_0^{\text{th}}$ 


  - Eigenbasis change  
 $\Gamma_1 = 43\Gamma_0^{\text{th}} \rightarrow 51\Gamma_0^{\text{th}}$ 

  - Double flips  
 $\Gamma_1 = 51\Gamma_0^{\text{th}} \rightarrow 94\Gamma_0^{\text{th}}$ 


  - $T_2^*$  increase  
 $\Gamma_1 = 94\Gamma_0^{\text{th}} \rightarrow 102\Gamma_0^{\text{th}}$ 

# Experiment: $\vec{B}$ in arbitrary direction




- 4-classes degeneracy
- Eigenbasis change
- Double-flips
- $T_2^*$  change




$\Gamma_1(B = 0) \approx 10 \Gamma_1(B \neq 0)$   
 $\sim 4\%$  PL contrast  
HWHM  $\sim 9 \text{ G}$

# Experiment: $\vec{B} \parallel [100]$





- 4-classes degeneracy
- Eigenbasis change
- Double-flips
- $T_2^*$  change




$\Gamma_1(B = 0) \approx 1.2 \Gamma_1(B \neq 0)$   
 $\sim 0.5\%$  PL contrast  
HWHM  $\sim 2$  G

Classes degeneracy is the dominant cause  
of depolarization at low magnetic field

# Experiment: $\vec{B} \perp [111]$



Same eigenbasis :  
 $|\pm\rangle = \frac{|+1\rangle \pm |-1\rangle}{\sqrt{2}}$   
 for  $\vec{B} \perp [111]$  than for  $\vec{B} = 0$




canceling out double flips  
 with transverse field


- 4 classes degeneracy
- Eigenbasis change
- Double-flips
- $T_2^*$  change

Double flips are the second dominant cause  
 of depolarization at low magnetic field

# Summary of the experimental observations




# Angular sensitivity of LFDM



- 4 classes degeneracy
- Eigenbasis change
- Double-flips
- $T_2^*$  change

- 4-classes degeneracy
- Eigenbasis change
- Double-flips
- $T_2^*$  change



The 4-classes degeneracy is not the limiting factor of the sensitivity

Sensing with quantum mechanics

Diamond and NV centers in practice

Low field depolarization magnetometry (LFDM)

Depolarization mechanisms in dense NV ensemble

# Conclusion

Sensing with quantum mechanics

Diamond and NV centers in practice

Low field depolarization magnetometry (LFDM)

Depolarization mechanisms in dense NV ensemble

# Acknowledgments