


Unit 4.3 Properties of the Laplace Transform

Contents

- [Agenda](#)
- [Some Selected Properties of the Laplace transform](#)
- [Table of Laplace Transform Properties](#)
- [Examples 10: Laplace transforms of common waveforms](#)
- [Homework](#)
- [Summary](#)
- [Next time](#)
- [References](#)
- [Answers to Examples 10](#)

The preparatory reading for this section is Chapter 2.3 of [\[Karris, 2012\]](#) and Chapter 3.3 of [\[Hsu, 2020\]](#).

Follow along at cpjobling.github.io/eg-150-textbook/laplace_transform/3/laplace_properties



Agenda

- [Some Selected Properties of the Laplace transform](#)
- [Table of Laplace Transform Properties](#)
- [Examples 10: Laplace transforms of common waveforms](#)

Some Selected Properties of the Laplace transform

Basic properties of the Laplace transform are presented in the following.

Verification of these properties is given in Worked Problems 3.8 to 3.16 in [\[Hsu, 2020\]](#) and in Section 2.2 in [\[Karris, 2012\]](#).

We will verify some of the properties in MATLAB Lab 4, but will not examine them further in this course.

You will, however, be expected to use **Tables of Properties**, such as that given in [Table of Laplace Transform Properties](#) below, to solve problems.

Linearity

$$c_1 f_1(t) + c_2 f_2(t) + \dots + c_n f_n(t) \Leftrightarrow c_1 F_1(s) + c_2 F_2(s) + \dots + c_n F_n(s)$$

Time shift

$$f(t - a)u_0(t - a) \Leftrightarrow e^{-as}F(s)$$

Frequency shift

$$e^{-at}f(t) \Leftrightarrow F(s + a)$$

Scaling

$$f(at) \Leftrightarrow \frac{1}{a}F\left(\frac{s}{a}\right)$$

Differentiation in the time domain

$$f'(t) = \frac{d}{dt}f(t) \Leftrightarrow sF(s) - f(0^-)$$

This property facilitates the solution of differential equations

The differentiation property can be extended to higher-orders as follows \$

$$f''(t) = \frac{d^2}{dt^2} f(t) \Leftrightarrow s^2 F(s) - s f(0^-) - f'(0^-)$$

$$f''(t) = \frac{d^3}{dt^3} f(t) \Leftrightarrow s^3 F(s) - s^2 f(0^-) - s f'(0^-) - f''(0^-)$$

and in general

$$f^{(n)}(t) = \frac{d^n}{dt^n} f(t) \Leftrightarrow s^n F(s) - s^{n-1} f(0^-) - s^{n-2} f'(0^-) - \dots - f^{(n-1)}(0^-)$$

Differentiation in the complex frequency domain

$$t f(t) \Leftrightarrow -\frac{d}{ds} F(s)$$

and in general

$$t^n f(t) \Leftrightarrow (-1)^n \frac{d^n}{ds^n} F(s)$$

Integration in the time domain

$$\int_{-\infty}^t f(\tau) d\tau \Leftrightarrow \frac{F(s)}{s} + \frac{f(0^-)}{s}$$

This property is important because it provides a way to model the solution of a differential equation using op-amp integrators in so-called [Analogue Computers](#) and is now the basis for numerical integration systems like Simulink.

Integration in the complex frequency domain

Providing that

$$\lim_{t \rightarrow 0} \frac{f(t)}{t}$$

exists

$$\frac{f(t)}{t} \Leftrightarrow \int_s^\infty F(s) ds$$

Time periodicity property

If $f(t)$ is a periodic function with period T such that $f(t) = f(t + nT)$ for $n = 1, 2, 3, \dots$ then

$$f(t + nT) \Leftrightarrow \frac{\int_0^T f(t) e^{-st} dt}{1 - e^{-sT}}$$

Initial value theorem

$$\lim_{t \rightarrow 0} f(t) \Leftrightarrow \lim_{s \rightarrow \infty} sF(s) = f(0^-)$$

Final value theorem

$$\lim_{t \rightarrow \infty} f(t) \Leftrightarrow \lim_{s \rightarrow 0} sF(s) = f(\infty)$$

Convolution in the time domain

$$f_1(t) * f_2(t) = \int_0^t f_1(\tau) f_2(t - \tau) d\tau \Leftrightarrow F_1(s) F_2(s)$$

This is another important result as it allows us to compute the response of a system by simply multiplying the Laplace transforms of the system and the signal and then inverse Laplace transforming the result.

This is usually much simpler than computing the convolution integral in the time domain – an operation we met in [C. Convolution Integral](#).

Convolution in the complex frequency domain

Multiplying two signals together in the time domain is the same as performing convolution in the complex frequency domain.

$$f_1(t) f_2(t) \Leftrightarrow \frac{1}{2\pi j} F_1(s) * F_2(s) = \frac{1}{2\pi j} \lim_{T \rightarrow \infty} \int_{c-jT}^{c+jT} F_1(\sigma) F_2(s - \sigma) d\sigma$$

Convolution in the complex frequency domain is nasty – multiplication in the time domain is relatively painless.

Table of Laplace Transform Properties

No.	Name	Time Domain $f(t)$	Complex Frequency Domain $F(s)$
1.	Linearity	$a_1 f_1(t) + a_2 f_2(t) + \dots + a_n f_n(t)$	$a_1 F_1(s) + a_2 F_2(s) + \dots + a_n F_n(s)$
2.	Time shifting	$f(t - a)u_0(t - a)$	$e^{-as}F(s)$
3.	Frequency shifting	$e^{-as}f(t)$	$F(s + a)$
4.	Time scaling	$f(at)$	$\frac{1}{a}F\left(\frac{s}{a}\right)$
5.	Time differentiation	$\frac{d}{dt}f(t)$	$F(s) - f(0^-)$
7.	Frequency differentiation	$tf(t)$	$-\frac{d}{ds}F(s)$
8.	Time integration	$\int_{-\infty}^t f(\tau)d\tau$	$\frac{F(s)}{s} + \frac{F(0^-)}{s}$
9.	Frequency integration	$\frac{f(t)}{t}$	$\int_s^\infty F(s)ds$
10.	Time Periodicity	$f(t + nT)$	$\frac{\int_0^T f(t)e^{-st}dt}{1 - e^{-sT}}$
11.	Initial value theorem	$\lim_{t \rightarrow 0} f(t)$	$\lim_{s \rightarrow \infty} sF(s) = f(0^-)$
12.	Final value theorem	$\lim_{t \rightarrow \infty} f(t)$	$\lim_{s \rightarrow 0} sF(s) = f(\infty)$
13.	Time convolution	$f_1(t) * f_2(t)$	$F_1(s)F_2(s)$
14.	Frequency convolution	$f_1(t)f_2(t)$	$\frac{1}{j2\pi}F_1(s) * F_2(s)$

See also: [Wikibooks: Engineering Tables/Laplace Transform Properties](#) and [Laplace Transform—WolframMathworld](#) for more complete references.

Examples 10: Laplace transforms of common waveforms

We will work through a few of the following on the board in class

- [Example 10.1: Pulse](#)
- [Example 10.2: Line segment](#)
- [Example 10.3: Triangular Pulse](#)
- [Example 10.4: Square Wave](#)
- [Example 10.5: Half-rectified Sinewave](#)

Example 10.1: Pulse

Use the tables of Laplace transforms and properties as appropriate to compute the Laplace transform of the pulse shown in [ig:ex10.1](#).




Fig. 45 A rectangular pulse.

For full solution see [Example 2.4.1](#) in Karris.



Example 10.2: Line segment

Use the tables of Laplace transforms and properties as appropriate to compute the Laplace transform of the line segment shown in [ig:ex10.2](#).




Fig. 46 A line segment.

For full solution see [Example 2.4.2](#) in Karris.



Example 10.3: Triangular Pulse

Use the tables of Laplace transforms and properties as appropriate to compute the Laplace transform of the triangular pulse shown in **ig:ex10.3**.




Fig. 47 A triangular pulse.

For full solution see [Examples in 2.4.3](#) in Karris.



Example 10.4: Square Wave

Use the tables of Laplace transforms and properties as appropriate to compute the Laplace transform of the of the periodic function shown in [ig:ex10.4](#).





Fig. 48 A square wave.

For full solution see [Example 2.4.4](#) in Karris.



Example 10.5: Half-rectified Sinewave

Use the tables of Laplace transforms and properties as appropriate to compute the Laplace transform of the half-rectified sine wave shown in [ig:ex10.5](#).




Fig. 49 Half-rectified sine wave.

For full solution see [Example 2.4.5](#) in Karris.



Homework

Attempt at least one of the end-of-chapter exercises from each question 1–7 of [Section 2.7](#) of [\[Karris, 2012\]](#). Don't look at the answers until you have attempted the problems.

If we have time, I will work through one or two of these in class.

Summary

In this section we have presented some of the most useful and commonly used properties of the Laplace transform, provided a table of Laplace Transform properties, and given examples of how properties and transform tables can be used to determine the Laplace transform of some useful aperiodic and periodic signals.

Take Aways

Still to come

The use of the *derivative property* in the complete solution of continuous-time LTI systems will be covered in [Unit 4.6: Transfer Functions](#)!

Next time

We move on to consider

- [Unit 4.4 The Inverse Laplace Transform](#)

References

- [Hsu20](1,2) Hwei P. Hsu. *Schaums outlines signals and systems*. McGraw-Hill, New York, NY, 2020. ISBN 9780071634724. Available as an eBook. URL: <https://www.accessengineeringlibrary.com/content/book/9781260454246>.
- [Kar12](1,2,3) Steven T. Karris. *Signals and systems with MATLAB computing and Simulink modeling*. Orchard Publishing, Fremont, CA., 2012. ISBN 9781934404232. Library call number: TK5102.9 K37 2012. URL: <https://ebookcentral.proquest.com/lib/swansea-ebooks/reader.action?docID=3384197>.

Answers to Examples 10

[Example 10.1: Pulse](#)

$$Au_0(t) - Au_0(t - a) \Leftrightarrow \frac{A(1 - e^{-as})}{s}.$$

[Example 10.2: Line segment](#)

$$(t - 1)u_0(t - 1) \Leftrightarrow \frac{e^{-s}}{s}.$$

[Example 10.3: Triangular Pulse](#)

$$f_T(t) \Leftrightarrow \frac{(1 - e^{-s})^2}{s^2}.$$

[Example 10.4: Square Wave](#)

$$f_R(t) \Leftrightarrow \frac{A \tanh\left(\frac{As}{2}\right)}{s}.$$

[Example 10.5: Half-rectified Sinewave](#)

$$f_{HW}(t) \Leftrightarrow \frac{1}{(s^2 + 1)(1 - e^{\pi s})}.$$

By Dr Chris P. Jobling

© Copyright Swansea University (2023).

This page was created by [Dr Chris P. Jobling](#) for [Swansea University](#).