

# Elementary Signals

The preparatory reading for this section is [Chapter 1](#) of [\[Kar12\]](#) which


- begins with a discussion of the elementary signals that may be applied to electrical circuits
- introduces the unit step, unit ramp and dirac delta functions
- presents the sampling and sifting properties of the delta function and
- concludes with examples of how other useful signals can be synthesised from these elementary signals.

## Colophon

An annotatable worksheet for this presentation is available as [Worksheet 3](#).

- The source code for this page is [elementary\\_signals/index.md](#).
- You can view the notes for this presentation as a webpage ([HTML](#)).
- This page is downloadable as a [PDF](#) file.

Consider the network shown in below where the switch is closed at time  $t = T$  and all components are ideal.




Express the output voltage  $V_{\text{out}}$  as a function of the unit step function, and sketch the appropriate waveform.

### Solution

Before the switch is closed at  $t < T$ :  $V_{\text{out}} = 0$ .

After the switch is closed for  $t > T$ :  $V_{\text{out}} = V_s$ .

We imagine that the voltage jumps instantaneously from 0 to  $V_s$  volts at  $t = T$  seconds as shown below.



We call this type of signal a step function.

## The Unit Step Function

$$u_0(t) = \begin{cases} 0 & t < 0 \\ 1 & t > 0 \end{cases}$$

[Colophon](#)

[The Unit Step Function](#)

[In Matlab](#)

[Simple Signal Operations](#)

[Amplitude Scaling](#)

[Time Reversal](#)

[Time Delay and Advance](#)

[Examples](#)

[Synthesis of Signals from the Unit Step](#)

[The Ramp Function](#)

[The Dirac Delta Function](#)

[The delta function](#)

[Sketch of the delta function](#)

[MATLAB Confirmation](#)

[Important properties of the delta function](#)

[Sampling Property](#)

[Sifting Property](#)

[Higher Order Delta Functions](#)

[Summary](#)

[Takeaways](#)

[Examples](#)

[Homework](#)


[References](#)

<

🔍

📄

Print to PDF



## In Matlab

In Matlab, we use the `heaviside` function (named after [Oliver Heaviside](#)).


File `plot_heaviside.m`

```
syms t
ezplot(heaviside(t), [-1,1])
heaviside(0)
```

```
imatlabs_export_fig('print-svg') % Static svg figures.
```

```
plot_heaviside
```

```
ans =
    0.5000
```



Note that, so that it can be plotted, Matlab defines the *heaviside function* slightly differently from the mathematically ideal unit step:


$$\text{heaviside}(t) = \begin{cases} 0 & t < 0 \\ 1/2 & t = 0 \\ 1 & t > 0 \end{cases}$$

## Simple Signal Operations

### Amplitude Scaling


Sketch  $Au_0(t)$  and  $-Au_0(t)$

```
syms t;
u0(t) = heaviside(t); % rename heaviside function for ease of use
A = 2; % so signal can be plotted
ezplot(A*u0(t), [-1,1]), grid, title('Amplitude scaling')
%%$Au_0(t)$','$', 'interpreter', 'latex')
```



Note that the signal is scaled in the  $y$  direction.

```
ezplot(-A*u0(t), [-1,1]),grid,title('Amplitude scaling and mirroring  $-Au_0(t)$ ','interpreter','latex')
```




Note that, because of the sign, the signal is mirrored about the  $x$  axis as well as being scaled by 2.

# Time Reversal

Sketch  $u_0(-t)$

```
ezplot(A*u0(-t), [-1,1]),grid,title('Time reversal  $Au_0(-t)$ ','interpreter','latex')
```




The sign on the function argument  $-t$  causes the whole signal to be reversed in time. Note that another way of looking at this is that the signal is mirrored about the  $y$  axis.

## Time Delay and Advance


Sketch  $u_0(t - T)$  and  $u_0(t + T)$

```
T = 1; % again to make the signal plottable.
ezplot(u0(t - T), [-1,2]),grid,title('Time delay  $u_0(t - T)$ ','interpreter','latex')
```



This is a *time delay* ... note for  $u_0(t - T)$  the step change occurs  $T$  seconds **later** than it does for  $u_0(t)$ .

```
ezplot(u0(t + T), [-2,1]),grid,title('Time advance  $u_0(t + T)$ ','interpreter','latex')
```



This is a *time advance* ... note for  $u_0(t+T)$  the step change occurs  $T$  seconds **earlier** than it does for  $u_0(t)$ .


## Examples

We will work through some examples in class. See [Worksheet 3](#).

## Synthesis of Signals from the Unit Step

Unit step functions can be used to represent other time-varying functions such as rectangular pulses, square waves and triangular pulses. See [Worksheet 3](#) for the examples that we will look at in class.

## The Ramp Function



In the circuit shown above  $i_s$  is a constant current source and the switch is closed at time  $t = 0$ .

When the current through the capacitor  $i_c(t) = i_s$  is a constant and the voltage across the capacitor is

$$v_c(t) = \frac{1}{C} \int_{-\infty}^t i_c(\tau) d\tau$$

where  $\tau$  is a dummy variable.

Since the switch closes at  $t = 0$ , we can express the current  $i_c(t)$  as

$$i_c(t) = i_s u_0(t)$$

and if  $v_c(t) = 0$  for  $t < 0$  we have

$$v_c(t) = \frac{i_s}{C} \int_{-\infty}^t u_0(\tau) d\tau = \underbrace{\frac{i_s}{C} \int_{-\infty}^0 0 d\tau}_0 + \frac{i_s}{C} \int_0^t 1 d\tau$$


So, the voltage across the capacitor can be represented as

$$v_C(t) = \frac{i_s}{C} t u_0(t)$$

**Note** that in this as in other examples throughout these notes, and in published tables of transforms, the inclusion of  $u_0(t)$  in  $v_C(t)$  acts as a “gating function” that limits the definition of the signal to the causal range  $0 \leq t < \infty$ .

To sketch the wave form, let’s arbitrarily let  $C$  and  $i_s$  be one and then plot with MATLAB.

```
C = 1; is = 1;
vc(t)=(is/C)*t*u0(t);
ezplot(vc(t), [-1,4]),grid,title('A ramp function')
```



This type of signal is called a **ramp function**. Note that it is the *integral* of the step function (the resistor-capacitor circuit implements a simple integrator circuit).

The unit ramp function is defined as

$$u_1(t) = \int_{-\infty}^t u_0(\tau) d\tau$$

so

$$u_1(t) = \begin{cases} 0 & t < 0 \\ t & t \geq 0 \end{cases}$$

and

$$u_0(t) = \frac{d}{dt} u_1(t)$$

**Note**


Higher order functions of  $t$  can be generated by the repeated integration of the unit step function.

For future reference, you should determine  $u_2(t)$ ,  $u_3(t)$  and  $u_n(t)$  for yourself and make a note of the general rule:

$$u_{n-1} = \frac{1}{n} \frac{d}{dt} u_n(t)$$

Details are given in equations 1.26—1.29 in Karris.

## The Dirac Delta Function



In the circuit shown above, the switch is closed at time  $t = 0$  and  $i_L(t) = 0$  for  $t < 0$ . Express the inductor current  $i_L(t)$  in terms of the unit step function and hence derive an expression for  $v_L(t)$ .

**Solution**

$$v_L(t) = L \frac{di_L}{dt}$$

Because the switch closes instantaneously at  $t = 0$

$$i_L(t) = i_s u_0(t)$$

Thus

$$v_L(t) = i_s L \frac{d}{dt} u_0(t).$$

To solve this problem we need to invent a function that represents the derivative of the unit step function. This function is called  $\delta(t)$  or the *dirac delta* function (named after [Paul Dirac](#)).

## The delta function

The unit impulse or the delta function, denoted as  $\delta(t)$ , is the derivative of the unit step.

This function is tricky because  $u_0(t)$  is discontinuous at  $t = 0$  but it must have the properties

$$\int_{-\infty}^t \delta(\tau) d\tau = u_0(t)$$

and

$$\delta(t) = 0 \ \forall \ t \neq 0.$$

## Sketch of the delta function



## MATLAB Confirmation

```
syms is L;
vL(t) = is * L * diff(u0(t))
```

```
vL(t) =
```

```
L*is*dirac(t)
```

Note that we can't plot `dirac(t)` in MATLAB with `ezplot`.

# Important properties of the delta function

## Sampling Property

The *sampling property* of the delta function states that

$$f(t)\delta(t - a) = f(a)\delta(t - a)$$

or, when  $a = 0$ ,

$$f(t)\delta(t) = f(0)\delta(t)$$

Multiplication of any function  $f(t)$  by the delta function  $\delta(t)$  results in sampling the function at the time instants for which the delta function is not zero.

The study of discrete-time (sampled) systems is based on this property.

*You should work through the proof for yourself.*

## Sifting Property

The *sifting property* of the delta function states that

$$\int_{-\infty}^{\infty} f(t)\delta(t - \alpha)dt = f(\alpha)$$

That is, if multiply any function  $f(t)$  by  $\delta(t - \alpha)$ , and integrate from  $-\infty$  to  $+\infty$ , we will get the value of  $f(t)$  evaluated at  $t = \alpha$ .

*You should also work through the proof for yourself.*

## Higher Order Delta Functions

the  $n$ th-order *delta function* is defined as the  $n$ th derivative of  $u_0(t)$ , that is

$$\delta^n(t) = \frac{d^n}{dt^n}[u_0(t)]$$

The function  $\delta'(t)$  is called the *doublet*,  $\delta''(t)$  is called the *triplet* and so on.

By a procedure similar to the derivation of the sampling property we can show that

$$f(t)\delta'(t - a) = f(a)\delta'(t - a) - f'(t)\delta(t - a)$$

Also, derivation of the sifting property can be extended to show that

$$\int_{-\infty}^{\infty} f(t)\delta^n(t - \alpha)dt = (-1)^n \frac{d^n}{dt^n}[f(t)] \Big|_{t=\alpha}$$

## Summary

In this chapter we have looked at some elementary signals and the theoretical circuits that can be used to generate them.

## Takeaways

- You should note that the unit step is the *heaviside function*  $u_0(t)$ .
- Many useful signals can be synthesized by use of the unit step as a “gating function” in combination with other signals
- That unit ramp function  $u_1(t)$  is the integral of the step function.

<

🔍

📄



- The *Dirac delta* function  $\delta(t)$  is the derivative of the unit step function. We sometimes refer to it as the *unit impulse function*.
- The delta function has sampling and sifting properties that will be useful in the development of *time convolution* and *sampling theory*.

## Examples

We will do some of these in class. See [Worksheet 3](#).

## Homework

These are for you to do later for further practice. See [Homework 1](#).

## References

See [Bibliography](#).

By Dr Chris P. Jobling  
© Copyright Swansea University (2019-2022).

This page was created by [Dr Chris P. Jobling](#) for [Swansea University](#).

