

# The Inverse Z-Transform

## Colophon

An annotatable worksheet for this presentation is available as [Worksheet 16](https://cpjobling.github.io/eg-247-textbook/dt_systems/3/worksheet16.html) ([https://cpjobling.github.io/eg-247-textbook/dt\\_systems/3/worksheet16.html](https://cpjobling.github.io/eg-247-textbook/dt_systems/3/worksheet16.html)).

- The source code for this page is [content/dt\\_systems/3/i\\_z\\_transform.ipynb](https://github.com/cpjobling/eg-247-textbook/blob/master/content/dt_systems/3/i_z_transform.ipynb) ([https://github.com/cpjobling/eg-247-textbook/blob/master/content/dt\\_systems/3/i\\_z\\_transform.ipynb](https://github.com/cpjobling/eg-247-textbook/blob/master/content/dt_systems/3/i_z_transform.ipynb)).
- You can view the notes for this presentation as a webpage ([HTML](https://cpjobling.github.io/eg-247-textbook/dt_systems/3/i_z_transform.html) ([https://cpjobling.github.io/eg-247-textbook/dt\\_systems/3/i\\_z\\_transform.html](https://cpjobling.github.io/eg-247-textbook/dt_systems/3/i_z_transform.html))).
- This page is downloadable as a [PDF](https://cpjobling.github.io/eg-247-textbook/dt_systems/3/i_z_transform.pdf) ([https://cpjobling.github.io/eg-247-textbook/dt\\_systems/3/i\\_z\\_transform.pdf](https://cpjobling.github.io/eg-247-textbook/dt_systems/3/i_z_transform.pdf)) file.

## Scope and Background Reading

This session we will talk about the Inverse Z-Transform and illustrate its use through an examples class.

The material in this presentation and notes is based on Chapter 9 (Starting at [Section 9.6](https://ebookcentral.proquest.com/lib/swansea-ebooks/reader.action?docID=3384197&ppg=351) (<https://ebookcentral.proquest.com/lib/swansea-ebooks/reader.action?docID=3384197&ppg=351>)) of {% cite karris %}.

## Agenda

- Inverse Z-Transform
- Examples using PFE
- Examples using Long Division

- Analysis in MATLAB

## The Inverse Z-Transform

The inverse Z-Transform enables us to extract a sequence  $f[n]$  from  $F(z)$ . It can be found by any of the following methods:

- Partial fraction expansion
- The inversion integral
- Long division of polynomials

### Partial fraction expansion

We expand  $F(z)$  into a summation of terms whose inverse is known. These terms have the form:

$$k, \frac{r_1 z}{z - p_1}, \frac{r_1 z}{(z - p_1)^2}, \frac{r_3 z}{z - p_2}, \dots$$

where  $k$  is a constant, and  $r_i$  and  $p_i$  represent the residues and poles respectively, and can be real or complex<sup>1</sup>.

### Notes

1. If complex, the poles and residues will be in complex conjugate pairs

$$\frac{r_i z}{z - p_i} + \frac{r_i^* z}{z - p_i^*}$$

### Step 1: Make Fractions Proper

- Before we expand  $F(z)$  into partial fraction expansions, we must first express it as a *proper* rational function.
- This is done by expanding  $F(z)/z$  instead of  $F(z)$
- That is we expand

$$\frac{F(z)}{z} = \frac{k}{z} + \frac{r_1}{z - p_1} + \frac{r_2}{z - p_2} + \dots$$

## Step 2: Find residues

- Find residues from

$$r_k = \lim_{z \rightarrow p_k} (z - p_k) \frac{F(z)}{z} = (z - p_k) \frac{F(z)}{z} \Big|_{z=p_k}$$

## Step 3: Map back to transform tables form

- Rewrite  $F(z)/z$ :

$$z \frac{F(z)}{z} = F(z) = k + \frac{r_1 z}{s - p_1} + \frac{r_2 z}{s - p_2} + \dots$$

We will work through an example in class.

[Skip next slide in Pre-Lecture]

## Example 1

Karris Example 9.4: use the partial fraction expansion to compute the inverse z-transform of

$$F(z) = \frac{1}{(1 - 0.5z^{-1})(1 - 0.75z^{-1})(1 - z^{-1})}$$



## MATLAB solution

See [example1.mlx \(matlab/example1.mlx\)](#). (Also available as [example1.m \(matlab/example1.m\)](#).)

Uses MATLAB functions:

- `collect` – expands a polynomial
- `sym2poly` – converts a polynomial into a numeric polynomial (vector of coefficients in descending order of exponents)
- `residue` – calculates poles and zeros of a polynomial
- `ztrans` – symbolic z-transform
- `iztrans` – symbolic inverse z-transform
- `stem` – plots sequence as a "lollipop" diagram

In [1]:

```
clear all
cd matlab
format compact
```

ans =

```
 '/Users/eechris/dev/eg-247-textbook/content/dt
_systems/3/matlab'
```

In [2]:

```
syms z n
```

The denominator of  $F(z)$

In [3]:

```
Dz = (z - 0.5)*(z - 0.75)*(z - 1);
```

Multiply the three factors of Dz to obtain a polynomial

In [4]:

```
Dz_poly = collect(Dz)
```

```
Dz_poly =
z^3 - (9*z^2)/4 + (13*z)/8 - 3/8
```

## Make into a rational polynomial

$$z^2$$

In [5]:

```
num = [0, 1, 0, 0];
```

$$z^3 - 9/4z^2 - 13/8z - 3/8$$

In [6]:

```
den = sym2poly(Dz_poly)
```

```
den =  
1.0000    -2.2500    1.6250   -0.3750
```

## Compute residues and poles

In [7]:

```
[r,p,k] = residue(num,den);
```

## Print results

- fprintf works like the c-language function

In [8]:

```
fprintf('\n')  
fprintf('r1 = %4.2f\t', r(1)); fprintf('p1 = %4.2f\n', p(1));.  
..  
fprintf('r2 = %4.2f\t', r(2)); fprintf('p2 = %4.2f\n', p(2));.  
..  
fprintf('r3 = %4.2f\t', r(3)); fprintf('p3 = %4.2f\n', p(3));
```

```
r1 = 8.00      p1 = 1.00  
r2 = -9.00    p2 = 0.75  
r3 = 2.00     p3 = 0.50
```

## Symbolic proof

$$f[n] = 2\left(\frac{1}{2}\right)^n - 9\left(\frac{3}{4}\right)^n + 8$$

In [9]:

```
% z-transform  
fn = 2*(1/2)^n-9*(3/4)^n + 8;  
Fz = ztrans(fn)
```

```
Fz =  
(8*z)/(z - 1) + (2*z)/(z - 1/2) - (9*z)/(z - 3/4)
```

In [10]:


```
% inverse z-transform  
iztrans(Fz)
```

```
ans =  
2*(1/2)^n - 9*(3/4)^n + 8
```

## Sequence

In [11]:

```
n = 1:15;  
sequence = subs(fn,n);  
stem(n,sequence)  
title('Discrete Time Sequence f[n] = 2*(1/2)^n-9*(3/4)^n + 8')  
;  
ylabel('f[n]')  
xlabel('Sequence number n')
```



## Example 2

Karris example 9.5: use the partial fraction expansion method to compute the inverse z-transform of

$$F(z) = \frac{12z}{(z+1)(z-1)^2}$$



## MATLAB solution

See [example2.mlx \(matlab/example2.mlx\)](#). (Also available as [example2.m \(matlab/example2.m\)](#).)



Uses additional MATLAB functions:


- `dimpulse` – computes and plots a sequence  $f[n]$  for any range of values of  $n$

In [ ]:

```
open example2
```


## Results

### 'Lollipop' Plot



### 'Staircase' Plot


Simulates output of Zero-Order-Hold (ZOH) or Digital Analogue Converter (DAC)



### Example 3

Karris example 9.6: use the partial fraction expansion method to compute the inverse z-transform of

$$F(z) = \frac{z + 1}{(z - 1)(z^2 + 2z + 2)}$$



## MATLAB solution


See [example3.mlx \(matlab/example3.mlx\)](#). (Also available as [example3.m \(matlab/example3.m\)](#).)

In [ ]:


```
open example3
```

## Results

Lollipop Plot



## Staircase Plot



## Inverse Z-Transform by the Inversion Integral

The inversion integral states that:

$$f[n] = \frac{1}{j2\pi} \oint_C F(z) z^{n-1} dz$$

where  $C$  is a closed curve that encloses all poles of the integrand.

This can (*apparently*) be solved by Cauchy's residue theorem!!

Fortunately (:-), this is beyond the scope of this module!

See Karris Section 9.6.2 (pp 9-29—9-33) if you want to find out more.

## Inverse Z-Transform by the Long Division

To apply this method,  $F(z)$  must be a rational polynomial function, and the numerator and denominator must be polynomials arranged in descending powers of  $z$ .


We will work through an example in class.

[Skip next slide in Pre-Lecture]

## Example 4

Karris example 9.9: use the long division method to determine  $f[n]$  for  $n = 0, 1$ , and  $2$ , given that

$$F(z) = \frac{1 + z^{-1} + 2z^{-2} + 3z^{-3}}{(1 - 0.25z^{-1})(1 - 0.5z^{-1})(1 - 0.75z^{-1})}$$



## MATLAB

See [example4.mlx \(matlab/example4.mlx\)](#). (also available as [example4.m \(matlab/example4.m\)](#).)

In [ ]:

open example4

## Results


sym\_den =

$$z^3 - (3*z^2)/2 + (11*z)/16 - 3/32$$

fn =

1.0000  
2.5000  
5.0625  
.....

## Combined Staircase/Lollipop Plot



# Methods of Evaluation of the Inverse Z-Transform

## Partial Fraction Expansion

### *Advantages*

- Most familiar.
- Can use MATLAB `residue` function.

### *Disadvantages*

- Requires that  $F(z)$  is a proper rational function.

## Invserion Integral

### *Advantage*

- Can be used whether  $F(z)$  is rational or not

### *Disadvantages*

- Requires familiarity with the *Residues theorem* of complex variable analysis.

## Long Division

### *Advantages*

- Practical when only a small sequence of numbers is desired.
- Useful when z-transform has no closed-form solution.

### *Disadvantages*

- Can use MATLAB `dimpulse` function to compute a large sequence of numbers.
- Requires that  $F(z)$  is a proper rational function.
- Division may be endless.



# Summary

- Inverse Z-Transform
- Examples using PFE
- Examples using Long Division
- Analysis in MATLAB

## *Coming Next*

- DT transfer functions, continuous system equivalents, and modelling DT systems in Matlab and Simulink.

# Reference

{% bibliography --cited %}

# Answers to Examples

## Answer to Example 1

$$f[n] = 2\left(\frac{1}{2}\right)^n - 9\left(\frac{3}{4}\right)^n + 8$$

## Answer to Example 2

$$f[n] = 3(-1)^n + 6n - 3$$

## Answer to Example 3

$$f[n] = -0.5\delta[n] + 0.4 + \frac{(\sqrt{2})^n}{10} \cos \frac{3n\pi}{4} - \frac{3(\sqrt{2})^n}{10} \sin \frac{3n\pi}{4}$$

## Answer to Example 4

$$f[0] = 1, f[1] = 5/2, f[2] = 81/16, \dots$$