

The gravitational wave energy spectrum of a parabolic encounter

Christopher P. L. Berry* and Jonathan R. Gair†

Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, United Kingdom

(Dated: August 25, 2010)

We derive an analytic expression for the energy spectrum of gravitational waves from a parabolic Keplerian binary by taking the limit of the Peters and Matthews spectrum for eccentric orbits. This demonstrates that the location of the peak of the energy spectrum depends primarily on the orbital periape rather than the eccentricity. We compare this weak-field result to strong-field calculations and find it is reasonably accurate ($\sim 10\%$) provided that the azimuthal and radial orbital frequencies do not differ by more than $\sim 10\%$. For equatorial orbits in the Kerr spacetime, this corresponds to periape radii of $r_p \gtrsim 20M$. These results will be of use to model radiation bursts from compact objects on highly eccentric orbits about massive black holes in the local Universe, which could be detected by LISA.

PACS numbers: 04.30.-w, 04.25.Nx, 97.60.Lf, 98.35.Jk

I. INTRODUCTION

An important source of gravitational waves for the proposed space-based gravitational wave detector, the Laser Interferometer Space Antenna (LISA) [1, 2], are the inspirals of stellar-mass compact objects into massive black holes in the centres of galaxies. During the last few years of inspiral these systems generate continuous gravitational waves in the LISA band, which will allow the detection of as many as several hundred systems out to redshift $z \sim 1$ [3]. However, prior to this phase, the inspiral object spends many years on a highly eccentric orbit, generating bursts of gravitational radiation at each periape passage. LISA could resolve individual bursts from sources in the nearby Universe. Initial estimates [4] suggested a LISA event rate of 18 yr^{-1} , including 15 yr^{-1} from the centre of the Milky Way, although this was subsequently revised downwards to 1 yr^{-1} with negligible contribution from extragalactic sources [5]. If even a single burst from the Galactic centre is detected during the LISA mission, this will provide an unparalleled probe of the structure of spacetime there.

The spectrum of radiation from these bursts will be well approximated by the spectrum of a parabolic orbit.¹ In this note we derive an analytic approximation to this spectrum by taking the limit of the Peters and Matthews [6, 7] (PM) energy spectrum for eccentric Keplerian binaries. We show that the peak of the spectrum depends primarily on the orbital periape and only weakly on the eccentricity. We also estimate the range of validity of the approximation (in Sec. III) by comparing to numerical Teukolsky data, finding that it is a good approximation for equatorial orbits in Kerr with periape $r_p \gtrsim 20M$. The parabolic spectrum takes a neat analytic

form that has not previously appeared in the literature. We therefore hope this note will be a useful resource for future work on gravitational radiation from high eccentricity orbits.

II. PARABOLIC LIMIT

A. Energy Spectrum

For an orbit of eccentricity e with periape radius r_p , Peters and Matthews [6] give the power radiated into the n th harmonic of the orbital angular frequency as

$$P(n) = \frac{32}{5} \frac{G^4}{c^5} \frac{M_1^2 M_2^2 (M_1 + M_2) (1 - e)^5}{r_p^5} g(n, e) \quad (1)$$

where the function $g(n, e)$ is defined in terms of Bessel functions of the first kind

$$g(n, e) = \frac{n^4}{32} \left\{ \left[J_{n-2}(ne) - 2eJ_{n-1}(ne) + \frac{2}{n}J_n(ne) + 2eJ_{n+1}(ne) - J_{n+2}(ne) \right]^2 + (1 - e^2) [J_{n-2}(ne) - 2J_n(ne) + J_{n+2}(ne)]^2 + \frac{4}{3n^2} [J_n(ne)]^2 \right\}. \quad (2)$$

The Keplerian orbital frequency is

$$\omega_1^2 = \frac{G(M_1 + M_2)(1 - e)^3}{r_p^3} = (1 - e)^3 \omega_c^2, \quad (3)$$

where ω_c is defined as the angular frequency of a circular orbit of radius r_p . The energy radiated per orbit into the n th harmonic, that is at frequency $\omega_n = n\omega_1$, is

$$E(n) = \frac{2\pi}{\omega_1} P(n); \quad (4)$$

as $e \rightarrow 1$ for a parabolic orbit, $\omega_1 \rightarrow 0$ as the orbital period becomes infinite. The energy radiated per orbit

* cplb2@ast.cam.ac.uk

† jgair@ast.cam.ac.uk

¹ We use “parabolic” to refer to marginally bound orbits. Marginally bound Keplerian orbits are parabola; in curved spacetimes they do not retain such a simple shape.

is the total energy radiated. The spacing of harmonics is $\Delta\omega = \omega_1$, giving energy spectrum

$$\left. \frac{dE}{d\omega} \right|_{\omega_n} \omega_1 = E(n). \quad (5)$$

Changing to linear frequency $2\pi f = \omega$,

$$\left. \frac{dE}{df} \right|_{f_n} = \frac{128\pi^2}{5} \frac{G^3}{c^5} \frac{M_1^2 M_2^2}{r_p^2} (1-e)^2 g(n, e) \quad (6)$$

$$= \frac{4\pi^2}{5} \frac{G^3}{c^5} \frac{M_1^2 M_2^2}{r_p^2} \ell(n, e), \quad (7)$$

where the function $\ell(n, e)$ is defined in the last line. For a parabolic orbit, we must take the limit of $\ell(n, e)$ as $e \rightarrow 1$.

We simplify $\ell(n, e)$ using the recurrence formulae (Watson [8] 2.12)

$$J_{\nu-1}(z) + J_{\nu+1}(z) = \frac{2\nu}{z} J_\nu(z) \quad (8)$$

$$J_{\nu-1}(z) - J_{\nu+1}(z) = 2J'_\nu(z), \quad (9)$$

and eliminate n using

$$n = \frac{\omega_n}{\omega_1} = (1-e)^{-3/2} \tilde{f}, \quad (10)$$

where $\tilde{f} = \omega_n/\omega_c = f_n/f_c$ is a dimensionless frequency. To find the limit we define two new functions

$$A(\tilde{f}) = \lim_{e \rightarrow 1} \left\{ \frac{J_n(ne)}{(1-e)^{1/2}} \right\}; \quad B(\tilde{f}) = \lim_{e \rightarrow 1} \left\{ \frac{J'_n(ne)}{1-e} \right\}. \quad (11)$$

To give a well defined energy spectrum, both of these must be finite.

The Bessel function has an integral representation

$$J_\nu(z) = \frac{1}{\pi} \int_0^\pi \cos(\nu\vartheta - z \sin \vartheta) d\vartheta; \quad (12)$$

we want the limit of this for $\nu \rightarrow \infty$, $z \rightarrow \infty$, with $z \leq \nu$. Using the stationary phase approximation, the dominant contribution to the integral comes from the regime in which the argument of the cosine is approximately zero (Watson [8] 8.2, 8.43), for small ϑ

$$J_\nu(z) \sim \frac{1}{\pi} \int_0^\pi \cos\left(\nu\vartheta - z\vartheta + \frac{z}{6}\vartheta^3\right) d\vartheta \quad (13)$$

$$\sim \frac{1}{\pi} \int_0^\infty \cos\left(\nu\vartheta - z\vartheta + \frac{z}{6}\vartheta^3\right) d\vartheta; \quad (14)$$

this last expression is an Airy integral and has a standard form (Watson [8] 6.4)

$$\int_0^\infty \cos(t^3 + xt) dt = \frac{\sqrt{x}}{3} K_{1/3}\left(\frac{2x^{3/2}}{3^{3/2}}\right), \quad (15)$$

where $K_\nu(z)$ is a modified Bessel function of the second kind. Using this to evaluate the limit gives

$$J_\nu(z) \sim \frac{1}{\pi} \sqrt{\frac{2(\nu-z)}{3z}} K_{1/3}\left(\frac{2^{3/2}}{3} \sqrt{\frac{(\nu-z)^3}{z}}\right). \quad (16)$$

For our case

$$J_n(ne) \sim \frac{1}{\pi} \sqrt{\frac{2}{3}} (1-e)^{1/2} K_{1/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right), \quad (17)$$

and the first limiting function is well defined,

$$A(\tilde{f}) = \frac{1}{\pi} \sqrt{\frac{2}{3}} K_{1/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right). \quad (18)$$

To find the derivative we combine (9) and (16), and expand to lowest order yielding

$$J'_n(ne) \sim -\frac{1}{2\pi} \sqrt{\frac{2}{3}} (1-e) \left[2^{3/2} K'_{1/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) + \frac{1}{\tilde{f}} K_{1/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) \right]. \quad (19)$$

We may re-express the derivative using the recurrence formula (Watson [8] 3.71)

$$K_{\nu-1}(z) - K_{\nu+1}(z) = -2K'_\nu(z) \quad (20)$$

to give

$$J'_n(ne) \sim \frac{1-e}{\sqrt{3}\pi} \left[K_{-2/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) + K_{4/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) - \frac{1}{\sqrt{2}\tilde{f}} K_{1/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) \right]. \quad (21)$$

And so finally,

$$B(\tilde{f}) = \frac{1}{\sqrt{3}\pi} \left[K_{-2/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) + K_{4/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) - \frac{1}{\sqrt{2}\tilde{f}} K_{1/3}\left(\frac{2^{3/2}\tilde{f}}{3}\right) \right], \quad (22)$$

which is also well defined.

Having obtained expressions for $A(\tilde{f})$ and $B(\tilde{f})$ in terms of standard functions, we can now calculate the energy spectrum for a parabolic orbit. From (7)

$$\frac{dE}{df} = \frac{4\pi^2}{5} \frac{G^3}{c^5} \frac{M_1^2 M_2^2}{r_p^2} \ell\left(\frac{f}{f_c}\right), \quad (23)$$

where we have used the limit

$$\ell(\tilde{f}) = \left[8\tilde{f}^2 B(\tilde{f}) - 2\tilde{f} A(\tilde{f}) \right]^2 + \left(128\tilde{f}^4 + \frac{4\tilde{f}^2}{3} \right) \left[A(\tilde{f}) \right]^2. \quad (24)$$

Figure 1 shows how $\ell(n, e)$ changes with eccentricity including our result for a parabolic encounter (cf. Fig. 3 of Peters and Matthews [6]). Although more power is radiated into higher harmonics, the peak of the spectrum does not move much: it is always between $f = f_c$ and $f = 2f_c$, with $f = 2f_c$ for $e = 0$ and $f \simeq 1.637f_c$ for $e = 1$.




FIG. 1. The relative energy (per orbit) spectrum $\ell(n, e)$ for $e = 0.2$ (heavy line), $e = 0.5$ (medium line), $e = 0.7$ (light line), and our limiting result for $e = 1$ (dashed line) versus frequency.

B. Total Energy

To check the validity of this limit we can calculate the total energy radiated by integrating (23) over all frequencies, or by summing the energy radiated into each harmonic. These must yield the same result. Summing over harmonics

$$E_{\text{sum}} = \frac{64\pi}{5} \frac{G^3}{c^5} \frac{M_1^2 M_2^2}{r_p^2} \omega_c (1-e)^{7/2} \sum_n g(n, e), \quad (25)$$

where we have used equations (1), (3) and (4). Peters and Matthews [6] provide the result

$$\sum_n g(n, e) = \frac{1 + (73/24)e^2 + (37/96)e^4}{(1 - e^2)^{7/2}}. \quad (26)$$

Using this,

$$E_{\text{sum}} = \frac{64\pi}{5} \frac{G^3}{c^5} \frac{M_1^2 M_2^2}{r_p^2} \omega_c \frac{1 + (73/24)e^2 + (37/96)e^4}{(1 + e)^{7/2}}, \quad (27)$$

which is perfectly well behaved as $e \rightarrow 1$,

$$E_{\text{sum}} = \frac{85\pi}{2^{5/2} 3} \frac{G^3}{c^5} \frac{M_1^2 M_2^2}{r_p^2} \omega_c. \quad (28)$$

Integrating the energy spectrum (23) gives

$$E_{\text{int}} = \frac{2\pi}{5} \frac{G^3}{c^5} \frac{M_1^2 M_2^2}{r_p^2} \omega_c \int_0^\infty \ell(\tilde{f}) d\tilde{f}. \quad (29)$$

The integral can be evaluated numerically as

$$\int_0^\infty \ell(\tilde{f}) d\tilde{f} = 12.5216858 \dots = \frac{425}{2^{7/2} 3}. \quad (30)$$

The two total energies are consistent, $E_{\text{int}} = E_{\text{sum}}$.




FIG. 2. Ratio of the total energy radiated as calculated using the Peters and Matthews [6] approach to that calculated by Martel [9] using black hole perturbation theory (solid line), and the reciprocal of the number of rotations $1/N$ (dashed line) versus periape radius r_p . The latter approach should give more accurate results. The Keplerian limit corresponds to $N = 1$.

III. APPLICABILITY

A. Limit of Approximation

The approach of Peters and Matthews assumes Keplerian orbits in flat spacetime. This should be a valid approximation in the weak-field regime far from a massive body. To find the limit of this approximation, we can compare the PM results with those from more accurate techniques. Energy spectra for parabolic orbits do not seem to be available in the literature yet, so we will make do with the total energy fluxes calculated by Martel [9], who uses time-domain black hole perturbation theory for a Schwarzschild black hole of mass M . Figure 2 shows the ratio of the two energies as a function of periape distance. As expected the PM result is more accurate for larger periapees. The agreement worsens as the periape distance decreases. At $r_p = 4M$, corresponding to the radius of the innermost stable circular orbit (ISCO), the energy flux calculated by Martel diverges, so the ratio tends to zero. This divergence is because in Schwarzschild (or Kerr) spacetime a parabolic orbit may have a zoom-whirl structure where it undergoes a number of near circular rotations (whirls) about the black hole. As the radius of the ISCO is approached, the number of whirls tends to infinity (in the absence of radiation reaction), so an infinite amount of energy is radiated. Figure 2 shows how the ratio of energies follows the number of rotations, defined as $N = \Delta\phi/2\pi$, where $\Delta\phi$ is the total change in the azimuthal angle over an orbit. As N increases the PM approximation worsens because the Keplerian orbit does not include this extra rotation. The accuracy of the PM result deteriorates rapidly once the orbit transitions to a zoom-whirl trajectory and is therefore far from parabolic in shape.

The PM result is accurate to $\sim 10\%$ for orbits with $N \lesssim 1.1$. We will adopt this as a cutoff point. For an equatorial orbit in Kerr spacetime

$$N = \frac{1}{\pi} \int_{r_p}^{\infty} \frac{d\phi}{dr} dr = \frac{L_z}{\pi\sqrt{2M}} \int_{r_p}^{\infty} \frac{r^2 - 2M(1 - a/L_z)r}{(r^2 - 2Mr + a^2)w} dr, \quad (31)$$

where

$$w^2 = r^3 - \frac{L_z^2}{2M}r^2 + (L_z - a)^2r; \quad (32)$$

L_z is the specific angular momentum about the z -axis; a is the spin parameter, and we have adopted units with $G = c = 1$. We will find it useful to define

$$r_{\pm} = M \pm \sqrt{M^2 - a^2}, \quad (33)$$

and the two non-zero roots of the cubic w^2

$$r_{p,1} = \frac{L_z^2}{4M} \pm \sqrt{\frac{L_z^4}{16M^2} - (L_z - a)^2}; \quad (34)$$

the periapsis is the larger root $r_p > r_1$. The integral may be rewritten as

$$N = \frac{L_z}{\pi\sqrt{2M}} \int_{r_p}^{\infty} \frac{1}{w} \left(1 + \frac{\alpha_+}{r - r_+} + \frac{\alpha_-}{r - r_-} \right) dr, \quad (35)$$

where

$$\alpha_{\pm} = \pm \frac{2Mar_{\pm} - a^2L_z}{2L_z\sqrt{M^2 - a^2}}. \quad (36)$$

This may be evaluated using elliptic integrals (Gradshteyn and Ryzhik [10] 3.131.8, 3.137.8)

$$N = \frac{L_z}{\pi} \sqrt{\frac{2}{r_p M}} \left[\frac{\alpha_+}{r_+} \Pi \left(\frac{r_+}{r_p} \middle| \frac{r_1}{r_p} \right) + \frac{\alpha_-}{r_-} \Pi \left(\frac{r_-}{r_p} \middle| \frac{r_1}{r_p} \right) \right], \quad (37)$$

where $\Pi(n|m) = \int_0^{\pi/2} d\vartheta / (1 - n \sin^2 \vartheta) \sqrt{1 - m \sin^2 \vartheta}$ is the complete elliptic integral of the third kind. In the limit of $a \rightarrow 0$ we recover the Schwarzschild result [11]

$$N = \frac{L_z}{\pi} \sqrt{\frac{2}{r_p M}} K \left(\frac{r_1}{r_p} \right), \quad (38)$$

where $K(m) = \int_0^{\pi/2} d\vartheta / \sqrt{1 - m \sin^2 \vartheta}$ is the complete elliptic integral of the first kind. Figure 3 shows the periapsis for which $N = 1.1$ for a range of spins. Equatorial

orbits with larger periapses should be reasonably approximated by the PM result.

Non-equatorial orbits are more complicated because of the additional precession of the orbital plane. This extra rotation will mean that the PM approach is less accurate; however, this should be sub-dominant to the perihelion precession effect and so the cutoff periapsis should not be much larger than for the equatorial case.




FIG. 3. Periapse radius corresponding to $N = 1.1$ as a function of spin parameter a (solid line). The curve may be approximated by a straight line $r_p = -3.91a + 17.36M$ (dashed line).

B. Astrophysical Implications

Considering bursts from the Galactic centre, orbits with periapses of $r_p \lesssim 120M$ could generate bursts that would be detectable with LISA [4, 5]. It is therefore likely that any such burst that was detected would be in the regime or validity of the Peters and Matthews approach, $r_p \gtrsim 20M$ for equatorial orbits. The results described in this note will therefore have application in that context and it should be possible to explore the majority of parameter space using this approximation. The most interesting orbits, those which come deep within the strong-field region of the black hole's spacetime, will be beyond the range of validity of this approximation, but these will be a small subset of all the events.

ACKNOWLEDGMENTS

CPLB is supported by an STFC studentship. JRG's work is supported by the Royal Society.

[1] P. Bender, A. Brillet, I. Ciufolini, A. M. Cruise, C. Cutler, K. Danzmann, F. Fidecaro, W. M. Folkner, J. Hough, P. McNamara, M. Peterseim, D. Robertson,

M. Rodrigues, A. Rüdiger, M. Sandford, G. Schäfer, R. Schilling, B. Schutz, C. Speake, R. T. Stebbins, T. Sumner, P. Touboul, J. Vinet, S. Vitale, H. Ward, and

- W. Winkler, *LISA Pre-Phase A Report*, Tech. Rep. (Max-Planck-Institut für Quantenoptik, Garching, 1998).
- [2] K. Danzmann and A. Rüdiger, *Classical Quantum Gravity*, **20**, S1 (2003).
 - [3] J. R. Gair, *Classical Quantum Gravity*, **26**, 094034 (2009).
 - [4] L. J. Rubbo, K. Holley-Bockelmann, and L. S. Finn, *Astrophys. J.*, **649**, L25 (2006).
 - [5] C. Hopman, M. Freitag, and S. L. Larson, *Mon. Not. R. Astron. Soc.*, **378**, 129 (2007).
 - [6] P. C. Peters and J. Mathews, *Phys. Rev.*, **131**, 435 (1963).
 - [7] P. C. Peters, *Phys. Rev.*, **136**, B1224 (1964).
 - [8] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd ed., Cambridge Mathematical Library (Cambridge University Press, Cambridge, 1995).
 - [9] K. Martel, *Phys. Rev. D*, **69**, 044025 (2004).
 - [10] I. Gradshteyn and I. Ryzhik, *Table of Integrals, Series, and Products*, sixth ed., edited by A. Jeffrey (Academic Press, London, 2000).
 - [11] C. Cutler, D. Kennefick, and E. Poisson, *Phys. Rev. D*, **50**, 3816 (1994).