



Unweighted legacy fit | OLD matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390


Distributions and residual correlations; raw residuals, no display shifts; statistics include tails
















Unweighted legacy fit | OLD matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390
xptar residual distributions by x sieve column; empty columns shown explicitly


Sieve column -5 (-12.5 cm)




Sieve column -4 (-10 cm)




Sieve column -3 (-7.5 cm)




Sieve column -2 (-5 cm)




Sieve column -1 (-2.5 cm)




Sieve column +0 (0 cm)




Sieve column +1 (2.5 cm)




Sieve column +2 (5 cm)




Sieve column +3 (7.5 cm)



Sieve column +4 (10 cm)




Sieve column +5 (12.5 cm)




Unweighted legacy fit | OLD matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390
ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Unweighted legacy fit | OLD matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390
yptar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)



Sieve column +5 (8.2 cm)



Unweighted legacy fit | OLD matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay

