



Supplied weights | NEW matrix | foil 0, z=+10 cm | -15 < delta < -6% | N=6684


Distributions and residual correlations; raw residuals, no display shifts; statistics include tails
















Supplied weights | NEW matrix | foil 0, z=+10 cm | -15 < delta < -6% | N=6684

xptar residual distributions by x sieve column; empty columns shown explicitly


Sieve column -5 (-12.5 cm)




Sieve column -4 (-10 cm)




Sieve column -3 (-7.5 cm)




Sieve column -2 (-5 cm)




Sieve column -1 (-2.5 cm)




Sieve column +0 (0 cm)




Sieve column +1 (2.5 cm)




Sieve column +2 (5 cm)




Sieve column +3 (7.5 cm)



Sieve column +4 (10 cm)




Sieve column +5 (12.5 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -15 < delta < -6% | N=6684

ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -15 < delta < -6% | N=6684

yptar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -15 < delta < -6% | N=6684

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay


Reconstructed sieve




xptar residual versus delta



ytar residual versus delta




yptar residual versus delta




Supplied weights | NEW matrix | foil 0, z=+10 cm | -6 <= delta < -2% | N=5211

Distributions and residual correlations; raw residuals, no display shifts; statistics include tails


xptar residual




ytar residual




yptar residual




xptar residual versus target



ytar residual versus target




yptar residual versus target




Supplied weights | NEW matrix | foil 0, z=+10 cm | -6 <= delta < -2% | N=5211

xptar residual distributions by x sieve column; empty columns shown explicitly


Sieve column -5 (-12.5 cm)




Sieve column -4 (-10 cm)




Sieve column -3 (-7.5 cm)




Sieve column -2 (-5 cm)




Sieve column -1 (-2.5 cm)




Sieve column +0 (0 cm)




Sieve column +1 (2.5 cm)




Sieve column +2 (5 cm)




Sieve column +3 (7.5 cm)



Sieve column +4 (10 cm)




Sieve column +5 (12.5 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -6 <= delta < -2% | N=5211
ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -6 <= delta < -2% | N=5211

yptar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)





Sieve column +5 (8.2 cm)





Supplied weights | NEW matrix | foil 0, z=+10 cm | -6 <= delta < -2% | N=5211

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay













Supplied weights | NEW matrix | foil 0, z=+10 cm | -2 <= delta < 2% | N=4564


Distributions and residual correlations; raw residuals, no display shifts; statistics include tails
















Supplied weights | NEW matrix | foil 0, z=+10 cm | -2 <= delta < 2% | N=4564

xptar residual distributions by x sieve column; empty columns shown explicitly


Sieve column -5 (-12.5 cm)




Sieve column -4 (-10 cm)




Sieve column -3 (-7.5 cm)




Sieve column -2 (-5 cm)




Sieve column -1 (-2.5 cm)




Sieve column +0 (0 cm)




Sieve column +1 (2.5 cm)




Sieve column +2 (5 cm)




Sieve column +3 (7.5 cm)



Sieve column +4 (10 cm)




Sieve column +5 (12.5 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -2 <= delta < 2% | N=4564

ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -2 <= delta < 2% | N=4564

yptar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | -2 <= delta < 2% | N=4564

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay


Reconstructed sieve




xptar residual versus delta



ytar residual versus delta




yptar residual versus delta




Supplied weights | NEW matrix | foil 0, z=+10 cm | 2 <= delta < 6% | N=3400

Distributions and residual correlations; raw residuals, no display shifts; statistics include tails


xptar residual




ytar residual




yptar residual




xptar residual versus target



ytar residual versus target




yptar residual versus target




Supplied weights | NEW matrix | foil 0, z=+10 cm | 2 <= delta < 6% | N=3400

xptar residual distributions by x sieve column; empty columns shown explicitly


Sieve column -5 (-12.5 cm)




Sieve column -4 (-10 cm)




Sieve column -3 (-7.5 cm)




Sieve column -2 (-5 cm)




Sieve column -1 (-2.5 cm)




Sieve column +0 (0 cm)




Sieve column +1 (2.5 cm)




Sieve column +2 (5 cm)




Sieve column +3 (7.5 cm)



Sieve column +4 (10 cm)




Sieve column +5 (12.5 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | 2 <= delta < 6% | N=3400

ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | 2 <= delta < 6% | N=3400

yptar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | 2 <= delta < 6% | N=3400

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay


Reconstructed sieve




xptar residual versus delta



ytar residual versus delta





yptar residual versus delta





Supplied weights | NEW matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390


Distributions and residual correlations; raw residuals, no display shifts; statistics include tails
















Supplied weights | NEW matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390

xptar residual distributions by x sieve column; empty columns shown explicitly


Sieve column -5 (-12.5 cm)




Sieve column -4 (-10 cm)




Sieve column -3 (-7.5 cm)




Sieve column -2 (-5 cm)




Sieve column -1 (-2.5 cm)




Sieve column +0 (0 cm)




Sieve column +1 (2.5 cm)




Sieve column +2 (5 cm)




Sieve column +3 (7.5 cm)



Sieve column +4 (10 cm)




Sieve column +5 (12.5 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390

ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)




Sieve column +4 (6.56 cm)





Sieve column +5 (8.2 cm)





Supplied weights | NEW matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390
yptar residual distributions by y sieve column; empty columns shown explicitly

































Supplied weights | NEW matrix | foil 0, z=+10 cm | 6 <= delta < 10% | N=2390

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay













Supplied weights | NEW matrix | foil 0, z=+10 cm | 10 <= delta < 22% | N=2147


Distributions and residual correlations; raw residuals, no display shifts; statistics include tails



















Supplied weights | NEW matrix | foil 0, z=+10 cm | 10 <= delta < 22% | N=2147
xptar residual distributions by x sieve column; empty columns shown explicitly































Supplied weights | NEW matrix | foil 0, z=+10 cm | 10 <= delta < 22% | N=2147
ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | 10 <= delta < 22% | N=2147
yptar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | NEW matrix | foil 0, z=+10 cm | 10 <= delta < 22% | N=2147

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay


Reconstructed sieve



xptar residual versus delta



ytar residual versus delta



yptar residual versus delta

