



Supplied weights | OLD matrix | foil 1, z=+0 cm | -15 < delta < -6% | N=6899


Distributions and residual correlations; raw residuals, no display shifts; statistics include tails
















Supplied weights | OLD matrix | foil 1, z=+0 cm | -15 < delta < -6% | N=6899

xptar residual distributions by x sieve column; empty columns shown explicitly


Sieve column -5 (-12.5 cm)




Sieve column -4 (-10 cm)




Sieve column -3 (-7.5 cm)




Sieve column -2 (-5 cm)




Sieve column -1 (-2.5 cm)




Sieve column +0 (0 cm)




Sieve column +1 (2.5 cm)




Sieve column +2 (5 cm)




Sieve column +3 (7.5 cm)



Sieve column +4 (10 cm)




Sieve column +5 (12.5 cm)




Supplied weights | OLD matrix | foil 1, z=+0 cm | -15 < delta < -6% | N=6899
ytar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | OLD matrix | foil 1, z=+0 cm | -15 < delta < -6% | N=6899

yptar residual distributions by y sieve column; empty columns shown explicitly


Sieve column -5 (-8.2 cm)




Sieve column -4 (-6.56 cm)




Sieve column -3 (-4.92 cm)




Sieve column -2 (-3.28 cm)




Sieve column -1 (-1.64 cm)




Sieve column +0 (0 cm)




Sieve column +1 (1.64 cm)




Sieve column +2 (3.28 cm)




Sieve column +3 (4.92 cm)



Sieve column +4 (6.56 cm)




Sieve column +5 (8.2 cm)




Supplied weights | OLD matrix | foil 1, z=+0 cm | -15 < delta < -6% | N=6899

Sieve reconstruction and residuals versus delta; fixed geometric xtar, not an iterative replay


Reconstructed sieve



xptar residual versus delta



ytar residual versus delta



yptar residual versus delta

