


Test 6.1: Faraday's Law

According to Faraday's law, an electromotive force V_{emf} is induced:



- (a) whenever a magnetic field \mathbf{B} crosses through the loop cross section.
- (b) only if \mathbf{B} is time-varying.
- (c) only if the loop is rotating.
- (d) Either \mathbf{B} is changing with time, the loop is rotating and \mathbf{B} is static, or \mathbf{B} is changing in time and the loop is rotating.



Test 6.2: Lenz's Law

The purpose of Lenz's law is to:

- (a) Compute the magnetic flux through the loop surface.
- (b) Compute the induced field \mathbf{B}_{ind} .
- (c) Determine the direction of I .
- (d) Determine the internal resistance R_i .

(a) Loop in a changing \mathbf{B} field

(b) Equivalent circuit

Test 6.3: Ideal Transformer

In an ideal transformer, the secondary to primary turns ratio is $N_2/N_1 = 10$. Consequently, only one of the following statements is true, which one?

- (a) $\frac{V_2}{V_1} = 0.1$, $\frac{I_2}{I_1} = 10$, $\frac{P_2}{P_1} = 100$
- (b) $\frac{V_2}{V_1} = 10$, $\frac{I_2}{I_1} = 0.1$, $\frac{P_2}{P_1} = 1$
- (c) $\frac{V_2}{V_1} = 10$, $\frac{I_2}{I_1} = 10$, $\frac{P_2}{P_1} = 100$
- (d) $\frac{V_2}{V_1} = 0.1$, $\frac{I_2}{I_1} = 10$, $\frac{P_2}{P_1} = 1$

Test 6.4: Piezoelectricity


Piezoelectric crystals are used in:

- (a) Ultrasound transducers, microphones, and speakers.
- (b) Thermocouples to measure temperature.
- (c) Accelerometers.

- (d) Proximity sensors.

Test 6.5: TV Antenna

A circular-loop TV antenna with 0.02-m^2 area is in the presence of a uniform-amplitude 300 MHz signal. When oriented for maximum response, the loop develops an emf with a peak value of 15 (mV). What is the peak magnitude B_0 of the incident wave?




- (a) $B_0 = 2$ (nT)
- (b) $B_0 = 0.4$ (nT)
- (c) $B_0 = 0.1$ (nT)
- (d) $B_0 = 2$ (μT)

Test 6.6: Square Loop

The square loop shown in the figure is coplanar with (but not touching) a long, straight wire carrying a current

$$I(t) = 5 \cos(2\pi \times 10^4 t) \quad (\text{A}).$$

What is the emf induced across a small gap created in the loop?




- (a) $V_{\text{emf}} = 7 \cos(2\pi \times 10^4 t)$ (mV)
- (b) $V_{\text{emf}} = 3.5 \sin(2\pi \times 10^4 t)$ (mV)
- (c) $V_{\text{emf}} = 0$
- (d) $V_{\text{emf}} = 7 \sin(2\pi \times 10^4 t)$ (mV)

Test 6.7: EM Generator

The rectangular conducting loop shown in the figure rotates at 6,000 revolutions per minute in a uniform magnetic flux density given by

$$\mathbf{B} = \hat{\mathbf{z}} 50 \quad (\text{mT}).$$




When an EE student connected a voltmeter across a small gap in the loop, the

voltmeter registered a voltage of zero. How should the configuration be changed so as to generate the maximum possible emf?

- (a) Reverse direction of \mathbf{B} .
- (b) Reverse direction of loop rotation.
- (c) Increase the magnitude of \mathbf{B} .
- (d) Change \mathbf{B} so it points along any direction in the x - y plane.

Test 6.8: EM Generator

The electromagnetic generator shown in the figure is connected to an electric bulb with a resistance of $150\ \Omega$. If the loop area is $0.1\ \text{m}^2$ and it rotates at 1800 revolutions per minute in a uniform magnetic flux density $B_0 = 1.6\ \text{T}$, determine the amplitude of the current generated in the light bulb.



- (a) $I \approx 200\ (\text{mA})$
- (b) $I \approx 10\ (\text{mA})$
- (c) $I \approx 100\ (\text{mA})$
- (d) $I \approx 1\ (\text{mA})$

Test 6.9: EM Potentials

The magnetic field in a dielectric material with $\epsilon = 9\epsilon_0$, $\mu = \mu_0$, and $\sigma = 0$ is

given by

$$\mathbf{H}(y,t) = \hat{\mathbf{x}}5 \cos(2\pi \times 10^7 t + 0.2\pi y) \quad (\text{A/m}).$$

Find the associated electric field \mathbf{E} .

- (a) $\hat{\mathbf{y}}30 \cos(2\pi \times 10^7 t + 0.2\pi y)$ (V/m)
- (b) $-\hat{\mathbf{z}}630 \cos(2\pi \times 10^7 t + 0.2\pi y)$ (V/m)
- (c) $\hat{\mathbf{z}}30 \cos(2\pi \times 10^7 t + 0.2\pi y)$ (V/m)
- (d) $\hat{\mathbf{x}}60 \cos(2\pi \times 10^7 t + 0.2\pi y)$ (V/m)

Test 6.10: EM Potentials

The electric field of an electromagnetic wave propagating in air is given by


$$\mathbf{E}(z,t) = \hat{\mathbf{x}}4 \cos(3 \times 10^8 t - z) + \hat{\mathbf{y}}3 \sin(3 \times 10^8 t - z) \quad (\text{V/m}).$$

Find the associated magnetic field $\mathbf{H}(z,t)$.

- (a) $\mathbf{H}(z,t) = \hat{\mathbf{x}}12 \cos(3 \times 10^8 t - z) + \hat{\mathbf{y}}8 \cos(3 \times 10^8 t - z)$ (A/m)
- (b) $\mathbf{H}(z,t) = -\hat{\mathbf{x}}8 \cos(3 \times 10^8 t - z) + \hat{\mathbf{y}}10.6 \cos(3 \times 10^8 t - z)$ (mA/m)
- (c) $\mathbf{H}(z,t) = -\hat{\mathbf{x}}8 \sin(3 \times 10^8 t - z) + \hat{\mathbf{y}}10.6 \cos(3 \times 10^8 t - z)$ (mA/m)
- (d) $\mathbf{H}(z,t) = \hat{\mathbf{x}}16 \cos(3 \times 10^8 t - z) + 2\hat{\mathbf{y}} \cos(3 \times 10^8 t - z)$ (A/m)

Test 6.11: Lenz's Law

Based on your understanding of Lenz's law, which of currents I_1 through I_4 is/are pointed in the wrong direction?



- (a) I_2 and I_3 are printed in the wrong direction.
- (b) All four currents are printed in the wrong direction.
- (c) I_1 and I_4 are printed in the wrong direction.
- (d) All four currents are printed correctly.

Test 6.12: Charge Dissipation

Suppose a certain amount of electric charge is placed at a point inside a block of carbon. Given that for carbon, $\epsilon_r = 20$ and $\sigma = 3 \times 10^4 \text{ (S/m)}$, how long does it take the charge density at that point to dissipate to 0.1% of its original value?

- (a) $t \approx 4 \text{ ms}$
- (b) $t \approx 4 \text{ ns}$
- (c) $t \approx 40 \text{ ps}$
- (d) $t \approx 40 \text{ fs}$

Test 6.13: Boundary Conditions

At the interface between two different materials, the boundary conditions for electric and magnetic fields:

- (a) are the same for static and dynamic (time-varying) conditions.
- (b) are not the same for static and dynamic (time-varying) conditions.
- (c) may or may not be the same depending on the relative conductivities of the two materials.
- (d) may or may not be the same depending on the charge relaxation time constants for the two materials.