

## SIEO 4150: REVIEW HMWK

### 1 Review: Not to be turned in for credit (solutions will be posted)

Start Reading Chapter 3 of the textbook.

1. Given a constant  $\lambda > 0$ , let

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } x \geq 0; \\ 0, & \text{otherwise.} \end{cases}$$

Compute the following:

- (a)  $\int_0^x f(u)du$  and  $\int_0^\infty f(x)dx$
- (b)  $\int_0^\infty xf(x)dx$
- (c)  $\int_0^\infty x^2 f(x)dx$
- (d) Let  $F(x) = \int_0^x f(u)du$ . Graph the function  $\bar{F}(x) \stackrel{\text{def}}{=} 1 - F(x)$ . Show that  $\int_0^\infty xf(x)dx = \int_0^\infty \bar{F}(x)dx$ .

2. Let

$$f(x) = \begin{cases} \frac{1}{2}, & \text{if } 2 < x < 4; \\ 0, & \text{otherwise.} \end{cases}$$

Graph  $f(x)$  and then compute the following:

- (a)  $\int_0^3 f(x)dx$  and  $\int_0^\infty f(x)dx$
- (b)  $\int_0^\infty xf(x)dx$
- (c)  $\int_0^\infty x^2 f(x)dx$
- (d) Let  $F(x) = \int_0^x f(u)du$ . Show that

$$F(x) = \begin{cases} 0, & \text{if } x \leq 2; \\ (x - 2)/2, & \text{if } 2 < x < 4; \\ 1, & \text{if } x \geq 4, \end{cases}$$

and that (recall that  $\bar{F}(x) \stackrel{\text{def}}{=} 1 - F(x)$ )

$$\bar{F}(x) = \begin{cases} 1, & \text{if } x \leq 2; \\ (4 - x)/2, & \text{if } 2 < x < 4; \\ 0, & \text{if } x \geq 4, \end{cases}$$

Graph both  $F(x)$  and  $\bar{F}(x)$ .

- (e) Show that  $\int_0^\infty xf(x)dx = \int_0^\infty \bar{F}(x)dx$ .

3. Let

$$f(x, y) = \begin{cases} 2e^{-x}e^{-2y}, & \text{if } x \geq 0, y \geq 0; \\ 0, & \text{otherwise.} \end{cases}$$

Compute

$$\int_0^1 \int_0^4 f(x, y)dx dy \text{ and } \int_0^\infty \int_0^\infty f(x, y)dx dy.$$

4. (a) Show that for any  $x$ ,

$$(1 + x + x^2 + \cdots + x^n) = \frac{1 - x^{n+1}}{1 - x}.$$

- (b) Use (a) to show that if  $|x| < 1$ , then

$$\sum_{k=0}^{\infty} x^k = \frac{1}{1 - x}. \quad (1)$$

- (c) Let  $g(x) = \frac{1}{1-x}$ ,  $0 < x < 1$ . Noting that  $g'(x) = \frac{1}{(1-x)^2}$ , argue from (b) that if  $0 < x < 1$ , then

$$\sum_{k=0}^{\infty} kx^k = \frac{x}{(1-x)^2}.$$

5. Consider tossing a fair coin until it first lands Heads (H). The sample space (set of all possible outcomes of this experiment) is given by the infinite set

$$\mathcal{S} = \{H, TH, TTH, TTTH, \dots\}.$$

For example,  $TTTTTH$  is the outcome “the coin landed tails the first 5 flips then landed heads on the 6th flip”, and

$$P(TTTTTTH) = \frac{1}{2} \times \frac{1}{2} \cdots \times \frac{1}{2} = \left(\frac{1}{2}\right)^6.$$

For simplicity of notation, let  $T^kH$  = the outcome “the coin landed tails the first  $k$  flips then landed heads on the  $(k+1)$ th flip”,  $k \geq 1$ , and  $T^0H = H$ . Then  $P(T^kH) = (1/2)^{k+1}$ .

Note that

$$P(\mathcal{S}) = \sum_{k=0}^{\infty} P(T^kH) = \sum_{k=1}^{\infty} \left(\frac{1}{2}\right)^k.$$

Use (4b) above to conclude that  $P(\mathcal{S}) = 1$ , and (4c) to conclude that

$$\sum_{k=1}^{\infty} kP(T^{k-1}H) = 2.$$

This last answer “2” is the average number of times the coin needs to be flipped until landing  $H$  for the first time.

Suppose that the coin, instead of being fair, lands  $H$  with “success” probability  $p = 1/4$ , and  $T$  with probability  $q = 3/4$ . Noting that then  $P(T^kH) = q^k p$ , re-do the above computation to obtain the average number of times the coin needs to be flipped until landing  $H$  for the first time. Finally, re-do for a general success probability  $0 < p < 1$ .