

SKIN COLOR SATISFACTION AND BINGE EATING

Although it has been demonstrated that (a) body dissatisfaction and internalization of societal appearance standards contribute to disordered eating and (b) that internalization of societal appearance standards leads to decreased skin color satisfaction among Black women, it has not been established whether skin color dissatisfaction contributes to disordered eating among Black women or girls. The objective of the present study is to determine the influence of skin color satisfaction as a potential predictor for binge eating, and its effect through body image in Black girls during the vulnerable developmental period of adolescence.

Parker, J.E., Enders, C.K., Mujahid, M.S., Laraia, B.A., Epel, E.S., Tomiyama, A.J. (2022). Prospective relationships between skin color satisfaction, body satisfaction, and binge eating in Black girls. *Body Image, 41*, 342-353.



Janet
Tomiyama



Jordan
Parker

KEY VARIABLES



Body Satisfaction

Body satisfaction is the facet of self-concept associated with weight, and includes the attitudes, evaluations, and feelings an individual holds about his or her own body.



Age


The grouping variable was age. Participants were followed longitudinally, with dependent variable measured at ages 10 and 18.

RESEARCH QUESTION

- Question: Do Black girls experience a change in body satisfaction during adolescence from age 10 to 18?
- The explanatory (independent) variable, age, consists of two occasions: ages 10 and 18
- The outcome (dependent) variable, body satisfaction, is a numeric scale derived from several questionnaire items

STATISTICAL ORG CHART

Bold typeface = 250A topic



RACIAL DISPARITIES IN PTSD

People of color in the United States disproportionately bear the burden of trauma and posttraumatic stress disorder (PTSD). Pregnant women of color are at particular risk, as perinatal PTSD is associated with adverse maternal and child health. However, PTSD is a heterogeneous disorder comprising discrete symptom dimensions. In a large sample of Latina, Black, and non-Hispanic White postpartum women in the United States (N = 1663), we examined racial and ethnic differences in the factors of the dysphoric arousal model—a leading dimensional model of PTSD.



Jennifer
Thomas



Jennifer
Sumner

Thomas, J.L., Carter, S.E., Dunkel-Schetter, Sumner, J.A. (2021). Racial and ethnic disparities in posttraumatic psychopathology among postpartum women. *Journal of Psychiatric Research*, 137, 36-40.

KEY VARIABLES



Racial Ethnic Identity

The study classified woman into three racial groups: Black, Latina, and non-Hispanic White



Anxious Arousal (Hyperarousal)


Anxious arousal (hyperarousal) is the abnormally heightened state of anxiety that occurs when thinking about a traumatic event (e.g., hypervigilance, exaggerated startle).

RESEARCH QUESTION

- Question: Do women of color (Black and Latina) differ from White women in their levels of anxious arousal?
- The explanatory (independent) variable, race, consists of three groups: Black, Latina, and White
- The outcome (dependent) variable, PTSD, is a numeric scale derived from several questionnaire items

STATISTICAL ORG CHART

Bold typeface = 250A topic





In small groups of two or three, work together to generate a comparative research question using variables from your domain of interest.

OUTLINE

1

Choosing the correct statistic

2

Comparative research questions

3

Association or trend research questions

4

Study questions

ASSOCIATION RESEARCH QUESTIONS

- Association (trend) research questions ask whether scores on two variables move in the same or opposite directions
- Is an increase in one variable accompanied by an increase or a decrease in the other?
- We focus on cases where the explanatory (independent) and outcome (dependent) variables are numeric scales

DISCRIMINATION AND INTERNALIZING BEHAVIOR

There is a well-documented relationship between perceived discrimination and internalizing symptoms among Latinx adolescents. However, few studies have examined how this psychosocial stressor relates to multiple domains of functioning in rural Latinx adolescents simultaneously. This study tested a spillover model of internalizing symptom development, where the negative effects of perceived discrimination are experienced through peer and family relationships.



Denise
Chavira



Giovanni
Ramos

Ramos, G., Delgadillo, D., Fossum, J., Montoya, A.K., Thamrin, H., Rapp, A., Escovar, E., Chavira, D.A. (2021). Discrimination and internalizing symptoms in rural Latinx adolescents: An ecological model of etiology. *Children and Youth Services Review*, 130, 1-10.

KEY VARIABLES



Perceived Discrimination

Perceived discrimination refers to individuals' perception of negative attitude, judgment, or unfair treatment due to their specific characteristics such as gender, race, ethnicity, and social status.



Internalizing Behavior

Internalizing behaviors are behaviors directed inwardly toward oneself. Examples include anxious and depressive symptoms, social withdrawal, and somatic (physical) complaints.

RESEARCH QUESTION


- Question: Is an increase in perceived discrimination accompanied by a systematic increase or decrease in internalizing behaviors?
- The explanatory (independent) variable, perceived discrimination, is a numeric scale derived from several questionnaire items
- The outcome (dependent) variable, internalizing behaviors, is a numeric scale derived from several questionnaire items



In small groups of two or three, use the statistical org chart to identify the correct statistical procedure for this study.

STATISTICAL ORG CHART

Bold typeface = 250A topic



DAILY DIARY STUDY OF MOOD

This study delivered a daily diary to 61 HIV infected men who have sex with men (MSM) between 16 and 24 years old for 66 days to measure HIV-risk behaviors and other psychosocial variables. The study examined the association between daily life stressors and daily negative mood. The study also examined the person-level association between the average number of daily life stressors across the 66 days and the average level of negative mood across that period.



Manuel
Ramirez



Patrick
Wilson

Ramirez, M, Wilson, P., Mitchell, R., Enders, C., & Woller, M. (in progress). Daily variability in depressed mood among gay and bisexual youth living with HIV. *Manuscript in preparation.*

KEY VARIABLES



Daily Life Stressors

Respondents were presented with list of stressful events (e.g., fights with family or friends, work stress, financial stress), and they checked how many they experienced each day.



Depression


Depression was measured using the Profile of Mood States (POMS). The POMS Depression-Dejection scale is computed by summing responses to five questionnaire items, where higher ratings reflect more negative mood.

RESEARCH QUESTION

- Question: Is there an association or trend between one's number of life stressors and their average depressive mood?
- The explanatory (independent) variable, number of life stressors, is a numeric value derived from a checklist
- The outcome (dependent) variable, depressive mood, is a numeric scale derived from several questionnaire items

STATISTICAL ORG CHART

Bold typeface = 250A topic





In small groups of two or three, work together to generate an association (trend) research question using variables from your domain of interest.

OUTLINE

1

Choosing the correct statistic

2

Comparative research questions

3

Association or trend research questions

4

Study questions

STUDY QUESTIONS

1. Generate a comparative research question using variables from your domain of interest.
2. Generate an association (trend) research question using variables from your domain of interest.